

ON UNIQUENESS FOR TIME HARMONIC ANISOTROPIC MAXWELL'S EQUATIONS WITH PIECEWISE REGULAR COEFFICIENTS

JOHN M. BALL, YVES CAPDEBOSCQ, AND BASANG TSERING XIAO

ABSTRACT. We are interested in the uniqueness of solutions to Maxwell's equations when the magnetic permeability μ and the permittivity ε are symmetric positive definite matrix-valued functions in \mathbb{R}^3 . We show that a unique continuation result for globally $W^{1,\infty}$ coefficients in a smooth, bounded domain, allows one to prove that the solution is unique in the case of coefficients which are piecewise $W^{1,\infty}$ with respect to a suitable countable collection of sub-domains with C^0 boundaries. Such suitable collections include any bounded finite collection. The proof relies on a general argument, not specific to Maxwell's equations. This result is then extended to the case when within these sub-domains the permeability and permittivity are only L^∞ in sets of small measure.

1. INTRODUCTION

Suppose we are given a time-harmonic incident electric field \mathcal{E}^i and magnetic field \mathcal{H}^i , special solutions of the time-harmonic homogeneous linear Maxwell equations of the form $\mathcal{E}^i = \Re(\mathbf{E}^i e^{-i\omega t})$ and magnetic field $\mathcal{H}^i = \Re(\mathbf{H}^i e^{-i\omega t})$, where $\mathbf{E}^i \in H_{\text{loc}}^1(\mathbb{R}^3)^3$ and $\mathbf{H}^i \in H_{\text{loc}}^1(\mathbb{R}^3)^3$ are complex-valued solutions of the homogeneous time-harmonic Maxwell equations

$$\begin{aligned}\nabla \wedge \mathbf{E}^i - i\omega\mu_0\mathbf{H}^i &= 0 \text{ in } \mathbb{R}^3, \\ \nabla \wedge \mathbf{H}^i + i\omega\varepsilon_0\mathbf{E}^i &= 0 \text{ in } \mathbb{R}^3,\end{aligned}$$

where μ_0 and ε_0 are positive constants, representing respectively the magnetic permeability and the electric permittivity of vacuum, and $\omega \in \mathbb{R} \setminus \{0\}$. The full time-harmonic electromagnetic field $(\mathbf{E}, \mathbf{H}) \in H_{\text{loc}}(\text{curl}; \mathbb{R}^3)$, where for any domain W we define

$$H_{\text{loc}}(\text{curl}; W) := \left\{ \mathbf{u} \in L_{\text{loc}}^2(W)^3 \text{ such that } \nabla \wedge \mathbf{u} \in L_{\text{loc}}^2(W)^3 \right\},$$

satisfies Maxwell's equations

$$(1.1) \quad \begin{aligned}\nabla \wedge \mathbf{E} - i\omega\mu_0\mu(x)\mathbf{H} &= 0 \text{ in } \mathbb{R}^3, \\ \nabla \wedge \mathbf{H} + i\omega\varepsilon_0\varepsilon(x)\mathbf{E} &= 0 \text{ in } \mathbb{R}^3,\end{aligned}$$

where ε and μ are real matrix-valued functions in $L^\infty(\mathbb{R}^3)^{3 \times 3}$. Decomposing the full electromagnetic field into its incident part and its scattered part,

$$(1.2) \quad \mathbf{E}^s := \mathbf{E} - \mathbf{E}^i, \text{ and } \mathbf{H}^s := \mathbf{H} - \mathbf{H}^i,$$

we assume that the scattered field satisfies the Silver-Müller radiation condition, uniformly in all directions, that is, if $x := r\theta$ then

$$(1.3) \quad \lim_{r \rightarrow \infty} \sup_{\theta \in S^2} |\mathbf{H}^s(r\theta) \wedge r\theta - r\mathbf{E}^s(r\theta)| = 0,$$

where $S^2 := \{x \in \mathbb{R}^3 \text{ such that } |x| = 1\}$ denotes the unit sphere.

This paper is about the existence of a unique solution to (1.1) satisfying (1.2) and (1.3), under the following additional hypotheses on ε and μ . We assume that both permittivity and permeability are real symmetric, uniformly positive definite and bounded, that is, there exist $0 < \alpha \leq \beta < \infty$ such that for all $\xi \in \mathbb{R}^3$ and almost every $x \in \mathbb{R}^3$,

$$(1.4) \quad \alpha |\xi|^2 \leq \varepsilon(x) \xi \cdot \xi \leq \beta |\xi|^2,$$

$$(1.5) \quad \alpha |\xi|^2 \leq \mu(x) \xi \cdot \xi \leq \beta |\xi|^2.$$

We suppose that ε and μ vary only in an open bounded domain Ω , so that

$$(1.6) \quad \varepsilon = \mu = \mathbf{I}_3 \text{ in } \Omega^c = \mathbb{R}^3 \setminus \Omega,$$

where \mathbf{I}_3 is the identity matrix in $\mathbb{R}^{3 \times 3}$. We assume that Ω is of the form

$$(1.7) \quad \Omega = \text{int} \left(\bigcup_{i \in I} \bar{\Omega}_i \right),$$

where the sub-domains Ω_i , $i \in I \subset \mathbb{N}$ are disjoint and of class C^0 , and int denotes the interior. The permittivity ε and the permeability μ are assumed to be piecewise $W^{1,\infty}$ with respect to the sub-domains Ω_i , so that for each $i \in I$, there exist $\varepsilon_i, \mu_i \in W^{1,\infty}(\mathbb{R}^3)^{3 \times 3}$ satisfying (1.4)-(1.5) and

$$(1.8) \quad \|\varepsilon_i\|_{W^{1,\infty}(\mathbb{R}^3)^{3 \times 3}} + \|\mu_i\|_{W^{1,\infty}(\mathbb{R}^3)^{3 \times 3}} \leq M_i,$$

where $M_i > 0$ is a positive constant, such that

$$(1.9) \quad \varepsilon(x) = \varepsilon_i(x) \text{ and } \mu(x) = \mu_i(x), \quad \text{a.e. } x \in \Omega_i.$$

Given a bounded set $A \subset \mathbb{R}^3$, we write $\mathcal{U}(A)$ as the (unique) unbounded component of \bar{A}^c .

Assumption 1. *For any $J \subset I$, and $\Omega_J = \text{int} \left(\bigcup_{j \in J} \bar{\Omega}_j \right)$, there exists $j_0 \in J$ such that $\partial\mathcal{U}(\Omega_J) \cap \partial\Omega_{j_0}$ admits an interior point relative to $\partial\mathcal{U}(\Omega_J)$. In other words, there exist $j_0 \in J$ and $x_0 \in \partial\mathcal{U}(\Omega_J) \cap \partial\Omega_{j_0}$ such that $B(x_0, \delta) \cap \partial\mathcal{U}(\Omega_J) \subset \partial\Omega_{j_0}$ for some $\delta > 0$.*

Proposition 1. *Assumption 1 holds if for all $J \subset I$, there exist $x_J \in \partial\mathcal{U}(\Omega_J)$ and $\delta_J > 0$ such that $B(x_J, \delta_J) \cap \Omega_j \neq \emptyset$ for only finitely many $j \in J$. In particular, Assumption 1 holds when I is finite.*

Proof. Given J , $x_J \in \partial\mathcal{U}(\Omega_J)$ and δ_J as in the statement of the proposition let $B_J = B(x_J, \delta_J)$ and let J' be the finite subset of J such that $B_J \cap \bigcup_{j \in J} \Omega_j = B_J \cap \bigcup_{j \in J'} \Omega_j$. We first show that

$$(1.10) \quad \partial\mathcal{U}(\Omega_J) \cap B_J = \bigcup_{j \in J} (\partial\mathcal{U}(\Omega_J) \cap B_J) \cap \partial\Omega_j.$$

Indeed, let $x \in \partial\mathcal{U}(\Omega_J) \cap B_J$. Then $x \notin \bigcup_{j \in J} \Omega_j$. We claim that there exists a sequence $x_k \in \bigcup_{j \in J} \Omega_j$ such that x_k tends to x . If not, for some $\eta > 0$ sufficiently




FIGURE 1. Left: an infinite collection of sub-domains satisfying Assumption 1. Centre: an infinite collection of sub-domains excluded by Assumption 1. Right: this collection satisfies Assumption 1.

small, we would have $B(x, \eta) \subset B_J$, and $B(x, \eta) \cap \bigcup_{j \in J} \Omega_j = B(x, \eta) \cap \bigcup_{j \in J'} \Omega_j = \emptyset$. On the other hand, there exists a sequence $y_k \in \mathcal{U}(\Omega_J)$ such that y_k tends to x . But $B(x, \eta)$ is connected and contained in $\bar{\Omega}_J^c$, thus $B(x, \eta) \subset \mathcal{U}(\Omega_J)$. This contradiction proves the claim.

Next, we note that $\partial\mathcal{U}(\Omega_J)$ is closed, thus complete in the subspace topology induced by \mathbb{R}^3 . Its intersection with the open ball B_J is an open subspace of $\partial\mathcal{U}(\Omega_J)$ by definition of the subspace topology. It is therefore a Baire space (see e.g. [11]). If a Baire Space is a countable union of closed sets, then one of the sets has an interior point. Using the identity (1.10), we obtain that there exists j_0 such that $\partial\mathcal{U}(\Omega_J) \cap B_J \cap \partial\Omega_{j_0}$ admits an interior point relative to $\partial\mathcal{U}(\Omega_J) \cap B_J$, that is, there exist $j_0 \in J$, $x_0 \in \partial\mathcal{U}(\Omega_J) \cap B_J$ and $\delta > 0$ such that $B(x_0, \delta) \cap \partial\mathcal{U}(\Omega_J) \cap B_J \subset \partial\Omega_{j_0}$.

Since B_J is open, $B(x_0, \delta) \cap B_J = B(x_0, \delta)$ when δ is sufficiently small, and we have established that Assumption 1 holds. \square

An example of a collection of sub-domains excluded by Assumption 1 is a collection of concentric shells concentrating on an exterior boundary, such as

$$(1.11) \quad \Omega_i = B\left(\mathbf{0}, \frac{i}{i+1}\right) \setminus \bar{B}\left(\mathbf{0}, \frac{i-1}{i}\right), \quad i = 1, 2, 3, \dots$$

In such a case, $\partial\mathcal{U}(\Omega)$ is the unit sphere, which is not the boundary of any of the subsets. On the other hand, Assumption 1 allows the sub-domains Ω_i to concentrate at a point or near an interior boundary. In Figure 1, we represent on the left a non-Lipschitz non-simply connected domain Ω which satisfies Assumption 1. In the centre, the domain given by (1.11) excluded by Assumption 1 is shown. On the right, we sketch a domain inspired by the one described by (1.11) which satisfies Assumption 1: near the accumulating boundary, interior points can be found on the wedge-shaped slit in the domain.

2. MAIN RESULT

Our main result is the following theorem.

Theorem 2. *Assume that (1.4)–(1.9) and Assumption 1 hold. If for a given $\omega \neq 0$, $\mathbf{E} \in H_{\text{loc}}(\text{curl}; \mathbb{R}^3)$ and $\mathbf{H} \in H_{\text{loc}}(\text{curl}; \mathbb{R}^3)$ are solutions of (1.1)–(1.3) corresponding to $\mathbf{E}^i = 0$ and $\mathbf{H}^i = 0$, then $\mathbf{E} = \mathbf{H} = 0$.*

There is a very long history concerning this problem, under various assumptions on the coefficients, see e.g. [1, 2, 3, 6, 7, 10, 13, 12, 14, 16] and the references therein. The improvement provided by the result in this work is that we assume that ε, μ are matrix-valued functions and that the sub-domains Ω_i are only of class C^0 . We do not assume that the sub-domains are Lipschitz as assumed for example in [6] for the isotropic (scalar) case. The authors are not aware of the existence of a general uniqueness result for the above problem when the coefficients are just $C^{0,\alpha}$ Hölder continuous, with $\alpha < 1$. For general elliptic equations, counter-examples to unique continuation, the main technique for proving uniqueness, are known in that case, see [8]. We remind the reader of the definition of a domain of class C^0 .

Definition 3. *A bounded domain Ω of \mathbb{R}^3 is of class C^0 if for any point x_0 on the boundary $\partial\Omega$, there exists a ball $B(x_0, \delta)$ and an orthogonal coordinate system (x_1, x_2, x_3) with origin at x_0 such that there exists a continuous function $f : C^0(\mathbb{R}^2; \mathbb{R})$ that satisfies*

$$\Omega \cap B(x_0, \delta) = \{x \in B(x_0, \delta) : x_3 > f(x_1, x_2)\}.$$

We define B_0 as the smallest open ball containing Ω . Note that the uniqueness of the solution outside B_0 is well known, due to the so-called Rellich's Lemma, see e.g. [2].

Lemma 4 (Rellich's Lemma). *If for a fixed ω , $\mathbf{E} \in H_{\text{loc}}(\text{curl}; \mathbb{R}^3)$ and $\mathbf{H} \in H_{\text{loc}}(\text{curl}; \mathbb{R}^3)$ are solutions of (1.1)-(1.3) corresponding to $\mathbf{E}^i = 0$ and $\mathbf{H}^i = 0$, then $\mathbf{E} = \mathbf{H} = 0$ in \bar{B}_0^c .*

Our proof relies on a recent unique continuation result [13] proved for globally $W^{1,\infty}$ regular coefficients.

Theorem 5 ([13]). *Let V be a connected open set in \mathbb{R}^3 . Assume that ε and μ are two real symmetric matrix valued functions in V satisfying (1.4)-(1.5), and*

$$\|\varepsilon\|_{W^{1,\infty}(V)^{3 \times 3}} + \|\mu\|_{W^{1,\infty}(V)^{3 \times 3}} \leq M,$$

for some constant $M > 0$. Suppose $(\mathbf{E}, \mathbf{H}) \in (L_{\text{loc}}^2(V))^2$ satisfy

$$\nabla \wedge \mathbf{E} - i\omega\mu_0\mu(x)\mathbf{H} = 0 \text{ in } V,$$

$$\nabla \wedge \mathbf{H} + i\omega\varepsilon_0\varepsilon(x)\mathbf{E} = 0 \text{ in } V.$$


Then, there exist $s > 0$ independent of V , \mathbf{E} and \mathbf{H} , such that if for some $x_0 \in V$, and for all $N \in \mathbb{N}$ and all $\delta > 0$ sufficiently small,

$$\int_{B(x_0, \delta)} (|\mathbf{E}|^2 + |\mathbf{H}|^2) dx \leq C_N \exp(-N\delta^{-s})$$

for some constant $C_N > 0$, then $\mathbf{E} = \mathbf{H} = 0$ in V .

The proof of Theorem 2 consists of three steps. The first two steps are given by the two propositions below.

Proposition 6. *Under the hypothesis of Theorem 2, suppose that $A \subset \mathbb{R}^3$ is a bounded open set and that for almost every $x \in \mathcal{U}(A)$ either $\varepsilon(x) = \mu(x) = \mathbf{I}_3$ or $\mathbf{E}(x) = \mathbf{H}(x) = 0$. Then $\mathbf{E} = \mathbf{H} = 0$ in $\mathcal{U}(A)$.*

FIGURE 2. A ball centred at an interior point on the boundary of $\mathcal{U}(\Omega_J)$.

Proof. For any $\mathbf{v}, \mathbf{w} \in L^2(\mathcal{U}(A))^2$, we have

$$(2.1) \quad \int_{\mathcal{U}(A)} \nabla \wedge \mathbf{E} \cdot \mathbf{v} \, dx - i\omega\mu_0 \int_{\mathcal{U}(A)} \mu(x)\mathbf{H} \cdot \mathbf{v} \, dx = 0,$$

$$(2.2) \quad \int_{\mathcal{U}(A)} \nabla \wedge \mathbf{H} \cdot \mathbf{w} \, dx + i\omega\varepsilon_0 \int_{\mathcal{U}(A)} \varepsilon(x)\mathbf{E} \cdot \mathbf{w} \, dx = 0,$$

where the integrals (2.1) and (2.2) are well defined by Rellich's Lemma 4. Since for almost every x in $\mathcal{U}(A)$, either $\varepsilon(x) = \mu(x) = \mathbf{I}_3$ or $\mathbf{E} = \mathbf{H} = 0$, the solutions of the system (2.1)-(2.2) can be written also in the form

$$\begin{aligned} \int_{\mathcal{U}(A)} \nabla \wedge \mathbf{E} \cdot \mathbf{v} \, dx - i\omega\mu_0 \int_{\mathcal{U}(A)} \mathbf{H} \cdot \mathbf{v} \, dx &= 0, \\ \int_{\mathcal{U}(A)} \nabla \wedge \mathbf{H} \cdot \mathbf{w} \, dx + i\omega\varepsilon_0 \int_{\mathcal{U}(A)} \mathbf{E} \cdot \mathbf{w} \, dx &= 0, \end{aligned}$$

which is the weak formulation of

$$\begin{aligned} \nabla \wedge \mathbf{E} - i\omega\mu_0 \mathbf{H} &= 0 \text{ in } \mathcal{U}(A), \\ \nabla \wedge \mathbf{H} + i\omega\varepsilon_0 \mathbf{E} &= 0 \text{ in } \mathcal{U}(A). \end{aligned}$$

Next, since A is bounded, thanks to Rellich's Lemma 4, $\mathbf{E} = \mathbf{H} = 0$ in $\mathcal{U}(A) \cap (\mathbb{R}^3 \setminus \bar{B}(R))$, for R large enough. In particular, \mathbf{E} and \mathbf{H} vanish in a ball contained in $\mathcal{U}(A)$, which is open and connected, and the conclusion follows from Theorem 5, applied with $\varepsilon(x) = \mu(x) = \mathbf{I}_3$, which in this case reduces to a well known result concerning the Helmholtz equation. \square

Proposition 7. *Let*

$$J := \{i \in I : |\mathbf{E}(x)|^2 + |\mathbf{H}(x)|^2 > 0 \text{ on a set of positive measure in } \Omega_i\}.$$

Then $J = \emptyset$.

Proof. Suppose for contradiction that J is nonempty. Then, by Assumption 1 there exists $x_0 \in \partial\mathcal{U}(\Omega_J) \cap \partial\Omega_{j_0}$ such that $B(x_0, \delta) \cap \partial\mathcal{U}(\Omega_J) \subset \partial\Omega_{j_0}$ for some $j_0 \in J$ and $\delta > 0$. To simplify notation, set $j_0 = 1$.

Let us show that there exist a point c on $\partial\mathcal{U}(\Omega_J) \cap \partial\Omega_1$ and a radius $\tilde{\delta} > 0$ such that

$$(2.3) \quad \Omega_J \cap B(c, \tilde{\delta}) = \Omega_1 \cap B(c, \tilde{\delta}).$$

Figure 2 sketches the configuration we have at hand around c .

Since Ω_1 has a C^0 boundary, for some (smaller) $\delta > 0$ there exists a continuous map f and a suitable orientation of axes such that $B(x_0, \delta) \cap \partial\Omega_J \subset \partial\Omega_1$ and

$$\Omega_1 \cap B(x_0, \delta) = \{x \in B(x_0, \delta) : x_3 > f(x_1, x_2)\}.$$

This alone does not prove our claim, since $B(x_0, \delta)$ could still intersect Ω_J when $x_3 \leq f(x_1, x_2)$. Since $x_0 \in \partial\mathcal{U}(\Omega_J)$, there exists a sequence $\{y_j\} \subset \mathcal{U}(\Omega_J) \cap B(x_0, \delta)$ such that y_j tends to x_0 . Consider for a fixed and sufficiently large j the line segment $\{y_j + te_3, t \geq 0\}$, and let $\tau > 0$ be the least value of t such that $y_j + te_3 \in \partial\Omega_J$. Then, $y_j + te_3 \notin \bar{\Omega}_J$, for $t < \tau$, and $y_j + \tau e_3 \in \partial\Omega_1$. Hence $y_j + \tau e_3 \notin \bigcup_{k \in J, k > 1} \Omega_k$. Since the sets Ω_k are disjoint, the line segment does not intersect $\bigcup_{k \in J, k > 1} \Omega_k$ in $B(x_0, \delta)$. The same argument applies to any line segment $\{z + te_3, t \geq 0\}$ for z sufficiently close to y_j . Introducing $c = y_j + \tau e_3$ we have established that there exists a ball $B(c, \tilde{\delta})$ such that $\Omega_1 \cap B(c, \tilde{\delta}) = \Omega_J \cap B(c, \tilde{\delta})$, which is (2.3).

Now, thanks to Proposition 6, and noting that (by Fubini's Theorem) each $\partial\Omega_i$ is of measure zero, $\mathbf{E} = \mathbf{H} = 0$ almost everywhere in $\mathcal{U}(\Omega_J)$. Thus, for almost every $x \in B(c, \tilde{\delta})$, either $\mathbf{E} = \mathbf{H} = 0$ or $\varepsilon(x) = \varepsilon_1(x)$, and $\mu(x) = \mu_1(x)$. Considering the weak formulation of Maxwell's equations, and arguing as in the proof of Proposition 6, we note that \mathbf{E} and \mathbf{H} are weak solutions of

$$\begin{aligned} \nabla \wedge \mathbf{E} - i\omega\mu_0\mu_1(x)\mathbf{H} &= 0 \text{ in } B(c, \tilde{\delta}), \\ \nabla \wedge \mathbf{H} + i\omega\varepsilon_0\varepsilon_1(x)\mathbf{E} &= 0 \text{ in } B(c, \tilde{\delta}), \end{aligned}$$

and vanish on the connected non-empty open set $B(c, \tilde{\delta}) \cap \{x_3 < f(x_1, x_2)\}$. Since ε_1 and μ_1 satisfy (1.8), that is,

$$\|\varepsilon_1\|_{W^{1,\infty}(\mathbb{R}^3)^{3 \times 3}} + \|\mu_1\|_{W^{1,\infty}(\mathbb{R}^3)^{3 \times 3}} \leq M_1,$$

Theorem 5 shows that $\mathbf{E} = \mathbf{H} = 0$ in $B(c, \tilde{\delta})$. This in turn shows that \mathbf{E} and \mathbf{H} vanish on a ball inside Ω_1 , and applying Theorem 5 in Ω_1 we obtain $\mathbf{E} = \mathbf{H} = 0$ almost everywhere in Ω_1 . This contradiction concludes the proof. \square

We now turn to the final step. We have obtained that $\mathbf{E} = \mathbf{H} = 0$ almost everywhere in Ω , and therefore either $\mathbf{E} = \mathbf{H} = 0$ or $\varepsilon(x) = \mu(x) = \mathbf{I}_3$ almost everywhere in \mathbb{R}^3 . Arguing as above, we deduce that (\mathbf{E}, \mathbf{H}) is a weak solution of (1.1) with $\varepsilon(x) = \mu(x) = \mathbf{I}_3$ everywhere and the conclusion of Theorem 2 follows from Rellich's Lemma.

3. THE CASE OF A MEDIUM WITH DEFECTS

We extend our result to the case when defects of small measure are allowed in the medium. One application is to liquid crystals (see [15] for more details). Namely, we assume that the permittivity and permeability are of the form

$$(3.1) \quad \begin{aligned} \varepsilon_D &= (1 - \mathbf{1}_D)\varepsilon + \mathbf{1}_D\tilde{\varepsilon}, \\ \mu_D &= (1 - \mathbf{1}_D)\mu + \mathbf{1}_D\tilde{\mu}, \end{aligned}$$

where ε and μ satisfy (1.4)-(1.9), $\mathbf{1}_D$ is the indicator function of a measurable bounded set D , such that

$$(3.2) \quad D \subset \bigcup_{i \in I} \Omega_i, \quad \overline{D} \cap \bar{\Omega}_i \subset \Omega_i \text{ and } \Omega_i \setminus \overline{D} \cap \bar{\Omega}_i \text{ is connected for each } i \in I,$$

and $\tilde{\varepsilon}$ and $\tilde{\mu}$ are real symmetric positive definite matrices in $L^\infty(\mathbb{R}^3)^{3 \times 3}$ satisfying (1.4)-(1.5).

Theorem 8. *Suppose that the electric and magnetic fields $\mathbf{E}_D \in H_{\text{loc}}(\text{curl}; \mathbb{R}^3)$ and $\mathbf{H}_D \in H_{\text{loc}}(\text{curl}; \mathbb{R}^3)$ are solutions of*

$$(3.3) \quad \begin{aligned} \nabla \wedge \mathbf{E}_D - i\omega\mu_0\mu_D(x)\mathbf{H}_D &= 0 \text{ in } \mathbb{R}^3, \\ \nabla \wedge \mathbf{H}_D + i\omega\varepsilon_0\varepsilon_D(x)\mathbf{E}_D &= 0 \text{ in } \mathbb{R}^3, \end{aligned}$$

together with the Silver-Müller radiation condition (1.3), and that ε_D and μ_D are given by (3.1), with D satisfying (3.2). Suppose Assumption 1 holds. Then, there exists a constant $d_0 > 0$ depending only on the measure $|B_0|$ of B_0 , $|\omega|$ and the lower and upper bounds α and β given in (1.4)-(1.5) such that if the measure of D satisfies $|D| < d_0$, then $\mathbf{E}_D = \mathbf{H}_D = 0$ almost everywhere.

To prove Theorem 8, we use the following variant of Theorem 2.

Proposition 9. *Under the same assumptions as Theorem 2, and assuming that (3.2) holds,*

$$\text{supp } \mathbf{H}_D \cup \text{supp } \mathbf{E}_D \subset \bar{D}.$$

Proof. The proof follows from that of Theorem 2, since by assumption for each $i \in I$, $\bar{D} \cap \Omega_i \subset \Omega_i$, and the boundary of $\Omega \setminus \Omega_i$ is unaltered by the defects. \square

Proof of Theorem 8. Since (3.3) admits a weak formulation, arguing as before we see using Proposition 9 that $\mathbf{E}_D \in H(\text{curl}; B_0)$ and $\mathbf{H}_D \in H(\text{curl}; B_0)$ have compact support in B_0 and are also solutions of

$$\begin{aligned} \nabla \wedge \mathbf{E}_D - i\omega\mu_0\hat{\mu}\mathbf{H}_D &= 0 \text{ in } B_0, \\ \nabla \wedge \mathbf{H}_D + i\omega\varepsilon_0\hat{\varepsilon}\mathbf{E}_D &= 0 \text{ in } B_0, \end{aligned}$$

where $\hat{\varepsilon} = \mathbf{I}_3 + \mathbf{1}_D(\tilde{\varepsilon} - \mathbf{I}_3)$, and $\hat{\mu} = \mathbf{I}_3 + \mathbf{1}_D(\tilde{\mu} - \mathbf{I}_3)$. Note that $i\omega\mu_0\hat{\mu}\mathbf{H}_D$ has compact support and is divergence free. Thus the Helmholtz decomposition (see e.g. [4, 5, 9]) of $i\omega\mu_0\hat{\mu}\mathbf{H}_D$ shows there exists a unique $\mathbf{A}_H \in H^1(B_0)^3$ such that $\mathbf{A}_H \cdot \nu = 0$, on ∂B_0 , $\text{div}(\mathbf{A}_H) = 0$ and such that $i\omega\mu_0\hat{\mu}\mathbf{H}_D = \nabla \wedge \mathbf{A}_H$. Furthermore, \mathbf{A}_H satisfies

$$\begin{aligned} \|\nabla \mathbf{A}_H\|_{L^2(B_0)^{3 \times 3}} &\leq C \left(\|\nabla \wedge \mathbf{A}_H\|_{L^2(B_0)^3} + \|\mathbf{A}_H\|_{L^2(B_0)^3} \right) \\ \text{and } \|\mathbf{A}_H\|_{L^2(B_0)^3} &\leq C|B_0|^{1/3} \|\nabla \wedge \mathbf{A}_H\|_{L^2(B_0)^3}, \end{aligned}$$

where C is a universal constant. Altogether this yields

$$(3.4) \quad \|\nabla \mathbf{A}_H\|_{L^2(B_0)^{3 \times 3}} \leq C\beta\mu_0|\omega|(|B_0| + 1)^{1/3} \|\mathbf{H}_D\|_{L^2(B_0)^3}.$$

Since $\mathbf{E}_D - \mathbf{A}_H$ is curl free, we deduce that there exists $p \in H^1(B_0)$ such that $\mathbf{E}_D = \mathbf{A}_H + \nabla p$, and p is uniquely defined by setting $\int_{B_0} p \, dx = 0$. Noticing that $\hat{\varepsilon}\mathbf{E}_D$ is divergence free, and $\hat{\varepsilon} - \mathbf{I}_3$ is compactly supported in B_0 we have that p is the solution of

$$\begin{aligned} \text{div}(\hat{\varepsilon}\nabla p) &= -\text{div}(\hat{\varepsilon}\mathbf{A}_H) \text{ in } B_0, \\ \nabla p \cdot n &= 0 \text{ on } \partial B_0, \\ \int_{B_0} p \, dx &= 0. \end{aligned}$$

Since \mathbf{A}_H is divergence free, the right-hand side becomes

$$-\text{div}(\hat{\varepsilon}\mathbf{A}_H) = -\text{div}(\mathbf{1}_D(\tilde{\varepsilon} - \mathbf{I}_3)\mathbf{A}_H).$$

To proceed, we compute using the Cauchy-Schwarz inequality the following bound

$$\begin{aligned} \alpha \|\nabla p\|_{L^2(B_0)^3}^2 &\leq \int_{B_0} \hat{\varepsilon} \nabla p \cdot \nabla p \, dx = - \int_{B_0} \mathbf{1}_D (\tilde{\varepsilon} - \mathbf{I}_3) \mathbf{A}_H \cdot \nabla p \, dx \\ &\leq (\beta + 1) \|\mathbf{A}_H\|_{L^2(D)^3} \|\nabla p\|_{L^2(B_0)^3}, \end{aligned}$$

and we have obtained that

$$\|\nabla p\|_{L^2(B_0)^3} \leq \frac{\beta + 1}{\alpha} \|\mathbf{A}_H\|_{L^2(D)^3}.$$

Next note using Proposition 9 that

$$\|\mathbf{E}_D\|_{L^2(B_0)^3} = \|\mathbf{E}_D\|_{L^2(D)^3} \leq \|\nabla p\|_{L^2(B_0)^3} + \|\mathbf{A}_H\|_{L^2(D)^3} \leq \frac{2\beta + 1}{\alpha} \|\mathbf{A}_H\|_{L^2(D)^3}.$$

The Sobolev-Gagliardo-Nirenberg inequality in B_0 shows that

$$\|\mathbf{A}_H\|_{L^6(B_0)^3} \leq C (|B_0| + 1)^{1/3} \|\mathbf{A}_H\|_{H^1(B_0)^3},$$

where C is a universal constant. Therefore, using Hölder's inequality, together with the Poincaré-Friedrichs estimate (3.4), we have

$$\|\mathbf{A}_H\|_{L^2(D)^3} \leq |D|^{1/3} \|\mathbf{A}_H\|_{L^6(B_0)^3} \leq C\beta (|B_0| + 1)^{2/3} \mu_0 |\omega| |D|^{1/3} \|\mathbf{H}_D\|_{L^2(B_0)^3}.$$

Altogether we have obtained

$$(3.5) \quad \|\mathbf{E}_D\|_{L^2(B_0)^3} \leq C \frac{\beta(\beta + 1)}{\alpha} (|B_0| + 1)^{2/3} \mu_0 |\omega| |D|^{1/3} \|\mathbf{H}_D\|_{L^2(B_0)^3}.$$

Repeating the same argument, but starting with \mathbf{H}_D , we obtain also

$$(3.6) \quad \|\mathbf{H}_D\|_{L^2(B_0)^3} \leq C \frac{\beta(\beta + 1)}{\alpha} (|B_0| + 1)^{2/3} \varepsilon_0 |\omega| |D|^{1/3} \|\mathbf{E}_D\|_{L^2(B_0)^3}.$$

The inequalities (3.5) and (3.6) imply that $\mathbf{H}_D = \mathbf{E}_D = 0$ when

$$(3.7) \quad |D| < d_0 := C \frac{\alpha^3}{\beta^3(\beta + 1)^3 (|B_0| + 1)^2 (\sqrt{\varepsilon_0 \mu_0} |\omega|)^3},$$

where C is a universal constant. \square

Remark 10. The dependence of the threshold constant d_0 given by (3.7) on $|\omega|$ and $|B_0|$ shows that for a permeability μ and a permittivity ε satisfying (1.4), (1.5) and (1.6) only, uniqueness for Maxwell's equations holds provided, if ω is fixed, the domain Ω is of small measure and bounded diameter, or, for a given Ω , when the absolute value of the frequency $|\omega|$ is sufficiently small. In such cases, the whole domain Ω can be taken as a defect D (and a fictitious ball containing D plays the role of Ω). We do not claim that the dependence of d_0 in terms of $|\omega|$ or $|B_0|$ in (3.7) is optimal. In contrast, Theorem 2 requires additional regularity assumptions on μ and ε , but does not depend on the frequency or the size of the domain.

ACKNOWLEDGEMENTS

The authors were supported by EPSRC Grant EP/E010288/1 and by the EPSRC Science and Innovation award to the Oxford Centre for Nonlinear PDE (EP/E035027/1).

REFERENCES

- [1] F. Cakoni and D. Colton. A uniqueness theorem for an inverse electromagnetic scattering problem in inhomogeneous anisotropic media. *Proc. Edinb. Math. Soc. (2)*, 46(2):293–314, 2003.
- [2] D. Colton and R. Kress. *Inverse acoustic and electromagnetic scattering theory*, volume 93 of *Applied Mathematical Sciences*. Springer-Verlag, Berlin, second edition, 1998.
- [3] M. M. Eller and M. Yamamoto. A Carleman inequality for the stationary anisotropic Maxwell system. *J. Math. Pures Appl. (9)*, 86(6):449–462, 2006.
- [4] K. O. Friedrichs. Differential forms on Riemannian manifolds. *Comm. Pure Appl. Math.*, 8:551–590, 1955.
- [5] V. Girault and P.-A. Raviart. *Finite element approximation of the Navier-Stokes equations*, volume 749 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1979.
- [6] C. Hazard and M. Lenoir. On the solution of time-harmonic scattering problems for Maxwell's equations. *SIAM J. Math. Anal.*, 27(6):1597–1630, 1996.
- [7] R. Leis. Über die eindeutige Fortsetzbarkeit der Lösungen der Maxwellschen Gleichungen in anisotropen inhomogenen Medien. *Bul. Inst. Politehn. Iași (N.S.)*, 14 (18)(fasc. 3-4):119–124, 1968.
- [8] K. Miller. Nonunique continuation for uniformly parabolic and elliptic equations in self-adjoint divergence form with Hölder continuous coefficients. *Arch. Rational Mech. Anal.*, 54:105–117, 1974.
- [9] Peter Monk. *Finite element methods for Maxwell's equations*. Numerical Mathematics and Scientific Computation. Oxford University Press, New York, 2003.
- [10] C. Müller. *Foundations of the mathematical theory of electromagnetic waves*. Revised and enlarged translation from the German. Die Grundlehren der mathematischen Wissenschaften, Band 155. Springer-Verlag, New York, 1969.
- [11] J. R. Munkres. *Topology: a first course*. Prentice-Hall Inc., Englewood Cliffs, N.J., 1975.
- [12] J.-C. Nédélec. *Acoustic and electromagnetic equations*, volume 144 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 2001. Integral representations for harmonic problems.
- [13] T. Nguyen and J.-N. Wang. Quantitative Uniqueness Estimate for the Maxwell System with Lipschitz Anisotropic Media. *Proc. Amer. Math. Soc.*, 140(2):595–605, 2011.
- [14] T. Ōkaji. Strong unique continuation property for time harmonic Maxwell equations. *J. Math. Soc. Japan*, 54(1):89–122, 2002.
- [15] B. Tsering Xiao. *Electromagnetic inverse problems for nematic liquid crystals*. PhD thesis, University of Oxford, 2011.
- [16] V. Vogelsang. On the strong unique continuation principle for inequalities of Maxwell type. *Math. Ann.*, 289(2):285–295, 1991.