



Shorter Notes: Strongly Continuous Semigroups, Weak Solutions, and the Variation of Constants Formula

J. M. Ball

Proceedings of the American Mathematical Society, Vol. 63, No. 2 (Apr., 1977),
370-373.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9939%28197704%2963%3A2%3C370%3ASNSCSW%3E2.0.CO%3B2-D>

Proceedings of the American Mathematical Society is currently published by American Mathematical Society.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ams.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is an independent not-for-profit organization dedicated to creating and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact support@jstor.org.

STRONGLY CONTINUOUS SEMIGROUPS, WEAK SOLUTIONS, AND THE VARIATION OF CONSTANTS FORMULA

J. M. BALL

ABSTRACT. Let A be a densely defined closed linear operator on a Banach space X , and let $f \in L^1(0, \tau; X)$. A definition of weak solutions of the equation $\dot{u} = Au + f(t)$ is given. It is shown that a necessary and sufficient condition for the existence of unique weak solutions for every initial data in X is that A generate a strongly continuous semigroup on X , and that in this case the solution is given by the variation of constants formula.

Let A be a densely defined closed linear operator on a real or complex Banach space X , let $\tau > 0$ and let $f \in L^1(0, \tau; X)$. Let $D(A) \subseteq X$ denote the domain of A . It is well known (cf. Kato [3, p. 486]) that if A is the generator of a strongly continuous semigroup of bounded linear operators $\{T(t)\}, t \geq 0$, on X , and if $x \in D(A)$, $f \in C([0, \tau]; X)$, then the equation

$$(1) \quad \dot{u}(t) = Au(t) + f(t), \quad t \in (0, \tau],$$

has a unique continuous solution satisfying $u(0) = x$, and that u is given by

$$(2) \quad u(t) = T(t)x + \int_0^t T(t-s)f(s) ds, \quad t \in [0, \tau].$$

When $x \in X$ is arbitrary, then unless $\{T(t)\}$ and f have special properties (e.g. $\{T(t)\}$ holomorphic and f Hölder continuous), $u(t)$ given by (2) will not, in general, belong to $D(A)$ for $t \in (0, \tau]$, so that (1) does not even make sense. The purpose of this note is to establish an abstract equivalence between functions u given by (2) and weak solutions, suitably defined, of (1), and to give a related characterization of strongly continuous semigroups. Although the proof of the theorem is simple, there seems to be no statement of it in the literature even in the case $f = 0$. An application to a class of nonlinear operator equations including certain nonlinear wave equations appears in [1].

Let A^* denote the adjoint of A and $\langle \cdot, \cdot \rangle$ the pairing between X and its dual space X^* .

DEFINITION. A function $u \in C([0, \tau]; X)$ is a *weak solution* of (1) if and only if for every $v \in D(A^*)$ the function $\langle u(t), v \rangle$ is absolutely continuous on $[0, \tau]$ and

Received by the editors June 24, 1976 and, in revised form, October 1, 1976.

AMS (MOS) subject classifications (1970). Primary 47D05; Secondary 34G05, 35D05.

Key words and phrases. Linear semigroups, weak solutions, variation of constants formula.

© American Mathematical Society 1977

$$(3) \quad \frac{d}{dt} \langle u(t), v \rangle = \langle u(t), A^*v \rangle + \langle f(t), v \rangle$$

for almost all $t \in [0, \tau]$.

THEOREM. *There exists for each $x \in X$ a unique weak solution $u(t)$ of (1) satisfying $u(0) = x$ if and only if A is the generator of a strongly continuous semigroup $\{T(t)\}$ of bounded linear operators on X , and in this case $u(t)$ is given by (2).*

We need the following lemma (cf. Goldberg [2, p. 127]).

LEMMA. *Let $x, z \in X$ satisfy $\langle z, v \rangle = \langle x, A^*v \rangle$ for all $v \in D(A^*)$. Then $x \in D(A)$ and $z = Ax$.*

PROOF. Let $G(A) \subseteq X \times X$ denote the graph of A , which is closed by assumption. By the Hahn-Banach theorem there exist $v, v^* \in X^*$, such that $\langle Ax, v \rangle + \langle x, v^* \rangle = 0$ for all $x \in D(A)$, and $\langle z, v \rangle + \langle x, v^* \rangle \neq 0$. Thus $v \in D(A^*)$, $v^* = -A^*v$ and $\langle z, v \rangle \neq \langle x, A^*v \rangle$, which is a contradiction.

PROOF OF THEOREM. Let A generate the strongly continuous semigroup $\{T(t)\}$. There exists a constant M such that $\|T(t)\| \leq M$ for $t \in [0, \tau]$. First note that if $x \in X$ and $v \in D(A^*)$ then $\langle T(t)x, v \rangle$ is differentiable with respect to t with derivative $\langle T(t)x, A^*v \rangle$. This is obvious if $x \in D(A)$, and holds for arbitrary $x \in X$ because $D(A)$ is dense and $\{T(t)\}$ strongly continuous. Let u be given by (2). It is easily shown that $u \in C([0, \tau]; X)$. For every $v \in D(A^*)$ and $t \in [0, \tau]$,

$$\langle u(t), v \rangle = \langle T(t)x, v \rangle + \int_0^t \langle T(t-s)f(s), v \rangle ds.$$

Suppose that $f \in C([0, \tau]; X)$. Since $(t, x) \mapsto T(t)x$ is jointly continuous on $[0, \tau] \times X$ it follows that

$$\frac{d}{dt} \int_0^t \langle T(t-s)f(s), v \rangle ds = \langle f(t), v \rangle + \int_0^t \langle T(t-s)f(s), A^*v \rangle ds,$$

so that $\langle u(t), v \rangle$ is differentiable for $t \in [0, \tau]$ and satisfies (3). If $f \in L^1(0, \tau; X)$, let $f_n \in C([0, \tau]; X)$ for $n = 1, 2, \dots$, with $f_n \rightarrow f$ in $L^1(0, \tau; X)$ and define

$$u_n(t) = T(t)x + \int_0^t T(t-s)f_n(s) ds, \quad s \in [0, \tau].$$

Then

$$\|u_n(t) - u(t)\| \leq M \int_0^t \|f_n(s) - f(s)\| ds,$$

so that $u_n \rightarrow u$ in $C([0, \tau]; X)$. But by the above, for each $v \in D(A^*)$,

$$\langle u_n(t), v \rangle = \langle x, v \rangle + \int_0^t [\langle u_n(s), A^*v \rangle + \langle f_n(s), v \rangle] ds, \quad t \in [0, \tau].$$

Passing to the limit we see that u is a weak solution of (1).

Next we prove that $u(t)$ is the only weak solution of (1) satisfying $u(0) = x$. Let $\bar{u}(t)$ be another such weak solution and set $w = u - \bar{u}$. Then

$$\langle w(t), v \rangle = \left\langle \int_0^t w(s) ds, A^*v \right\rangle$$

for all $v \in D(A^*)$, $t \in [0, \tau]$, so that by the lemma, $z(t) = \text{def} \int_0^t w(s) ds$ belongs to $D(A)$ and $\dot{z} = Az$. By [3, p. 481] $z = 0$ and hence $u = \bar{u}$.

Suppose that A is such that (1) has, for each $x \in X$, a unique weak solution $u(t)$ satisfying $u(0) = x$. For $t \in [0, \tau]$ define $T(t)x = u(t) - u_0(t)$, where u_0 is the weak solution of (1) satisfying $u_0(0) = 0$. If $t \geq 0$ let $t = n\tau + s$, where n is a nonnegative integer and $s \in [0, \tau)$, and define $T(t)x = T(s)T(\tau)^n x$. The map $\theta: X \rightarrow C([0, \tau]; X)$ defined by $\theta(x) = T(\cdot)x$ has closed graph and, hence, $T(\cdot)$ is a strongly continuous semigroup. Let B be the generator of $T(\cdot)$ and let $x \in D(B)$. For any $v \in D(A^*)$,

$$\frac{d}{dt} \langle T(t)x, v \rangle \Big|_{t=0} = \langle Bx, v \rangle = \langle x, A^*v \rangle.$$

It follows from the lemma that $x \in D(A)$ and $Bx = Ax$. In particular, $D(B) \subseteq D(A)$. The proof of the theorem is completed by showing that $D(A) \subseteq D(B)$.

Let $x \in D(A)$. Using the lemma we see that for each $t \in [0, \tau]$ the integrals $\int_0^t T(s)x ds$ and $\int_0^t T(s)Ax ds$ belong to $D(A)$ and

$$(4) \quad T(t)x = x + A \int_0^t T(s)x ds,$$

$$(5) \quad T(t)Ax = Ax + A \int_0^t T(s)Ax ds.$$

Consider the function

$$z(t) = \int_0^t T(s)Ax ds - A \int_0^t T(s)x ds.$$

It follows from (4) that $z \in C([0, \tau]; X)$. Clearly $z(0) = 0$. Let $v \in D(A^*)$. Using (4) and (5) we see that

$$\frac{d}{dt} \langle z(t), v \rangle = \langle z(t), A^*v \rangle, \quad t \in [0, \tau].$$

But it follows from our assumptions that the equation $\dot{z} = Az$, $z(0) = 0$, has only the zero weak solution. Hence

$$\int_0^t T(s)Ax ds = A \int_0^t T(s)x ds, \quad t \in [0, \tau].$$

Therefore by (4),

$$\lim_{t \rightarrow 0^+} \frac{1}{t} [T(t)x - x] = Ax$$

and, hence, $x \in D(B)$.

NOTE ADDED IN PROOF. The 'if' part of the above theorem is stated and proved by a somewhat different method in the recent book by Balakrishnan

[4, Theorem 4.8.3] under the assumption that X is a Hilbert space.

REFERENCES

1. J. M. Ball, *On the asymptotic behaviour of generalized processes, with applications to nonlinear evolution equations*, J. Differential Equations (to appear).
2. S. Goldberg, *Unbounded linear operators: Theory and applications*, McGraw-Hill, New York, 1966. MR 34 # 580.
3. T. Kato, *Perturbation theory for linear operators*, Springer-Verlag, New York, 1966. MR 34 # 3324.
4. A. V. Balakrishnan, *Applied functional analysis*, Applications of Mathematics, vol. 3, Springer, New York, 1976.

DEPARTMENT OF MATHEMATICS, HERIOT-WATT UNIVERSITY, RICcartON, CURRIE, EDINBURGH EH14 4AS, SCOTLAND