

PDE-CDT Core Course

Analysis of Partial Differential Equations-Part III

**EPSRC Centre for Doctoral Training in
Partial Differential Equations
Trinity Term**

27 April – 15 June 2016 (16 hours)

Final Exam: 27 June 2016

Course format: Teaching Course (TT)

**By Prof. Gui-Qiang G. Chen
Prof. Qian Wang**



1. Introduction

Overview: Theory of Hyperbolic PDEs

is a large subject, which has close connections with the other areas of mathematics including

Analysis, Mechanics, Mathematical Physics, Differential Geometry/Topology, ...

Besides its mathematical importance, it has a wide range of applications in

Engineering, Physics, Biology, Economics, ...

- Backbone of the year: Introduction to most facets of the theory of hyperbolic PDEs and related PDEs:
Features, methods, approaches, connections,
- Knowledge with PDE Foundation module and Analysis of PDEs (Parts I-II) is desirable

Synopsis - I:

1. Introduction

Part I: Hyperbolic Systems of First-Order Equations

2. Linear Theory:

Spaces involving time;

Hyperbolic systems of first-order equations, examples;

Weak solutions, well-posedness;

Vanishing viscosity method, energy methods,

Fourier transform method.

3. Nonlinear Theory I - Multi-D Scalar Conservation Laws:

L^1 - well-posedness theory, test function methods,
vanishing viscosity method;

*Other methods (numerical methods, kinetic method, relaxation
method, the layering method, ...);

*Further results (compactness, regularity, decay, trace, structure).

Synopsis - II:

4. Nonlinear Theory II – One-Dimensional Systems of Conservation Laws:

Riemann problem, Cauchy problem;

Elementary waves: shock waves, rarefaction waves,
contact discontinuities;

Lax entropy conditions;

Glimm scheme, front-tracking, BV solutions;

Compensated compactness, entropy analysis, L^p solutions,
vanishing viscosity methods;

*Uniqueness and continuous dependence; ...

5. *Nonlinear Theory III - Multidimensional Systems of Conservation Laws:

Basic features/phenomena (re-visit);

Local existence and stability; formation of singularities;

Discontinuities and free boundary problems;

Stability of shock waves, rarefaction waves, vortex sheets,
entropy waves.

Synopsis - III:

Part II: Second-Order Wave Equations

6. Energy estimates and local existence, Galerkin method;
Global existence of semi-linear wave equations with small data
(Quasilinear case could be similarly treated);
Lower regularity results for large data;
*Littlewood-Paley theory and Strichartz estimates.

* Optional


References:

1. **P. D. Lax: Hyperbolic Systems of Conservation Laws and the Mathematical Theory of Shock Waves.** CBMS, SIAM, 1973.
2. **C. M. Dafermos: Hyperbolic Conservation Laws in Continuum Physics,** Third edition. Springer-Verlag: Berlin, 2010.
3. **L. C. Evans: Partial Differential Equations,** Second edition. AMS: Providence, RI, 2010.
4. **L. Hormander: Lectures on Nonlinear Hyperbolic Differential Equations,** Springer-Verlag: Berlin-Heidelberg, 1997
5. **Bressan, G.-Q. Chen, M. Lewicka, D. Wang: Nonlinear Conservation Laws and Applications,** IMA Volume 153, Springer: New York, 2011.
6. **D. Serre, Systems of Conservation Laws, Vols. I, II,** Cambridge University Press: Cambridge, 1999, 2000.
7. **C. D. Sogge, Lectures on Nonlinear Wave Equations,** Second edition. International Press, Boston, MA, 2008.
8. **P. D. Lax: Hyperbolic Differential Equations,** AMS: Providence, 2000

Conservation Laws:

Rate of change of the total amount of certain quantity contained in a fixed region Ω

= Flux of this quantity across the boundary $\partial\Omega$ of the region




- The amount of such a quantity in any region can be measured by accounting for how much of it is currently present and how much of it enters or leaves the region in any fixed period of time.

Examples: Three Fundamental Laws of Nature

- ❑ **Conservation Laws of Mass and Energy:** Mass and Energy can be neither created nor destroyed.
- ❑ **Conservation Law of Momentum:** The total momentum of a closed system of objects remains constant through time.

Conservation Laws

Rate of Change of the Total Amount of
Certain Quantity in a Fixed Region Ω
= Flux of the Quantity across the Boundary $\partial\Omega$.



Conservation Law via Calculus

$$\iff \frac{d}{dt} \int_{\Omega} u dx = - \int_{\partial\Omega} \mathbf{f} \cdot \mathbf{n} dS$$

u – Density of the Quantity
 \mathbf{n} – Outward Normal to Ω

\mathbf{f} – Flux of the Quantity
 dS – Surface Element on $\partial\Omega$

Calculus Manipulations \implies

$$\partial_t u + \nabla_x \cdot \mathbf{f}(u) = 0$$

Physical Systems with $m \geq 2$ Quantities – Density Functions

\implies **Systems of Conservation Laws:**

$$\partial_t \mathbf{u} + \nabla \cdot \mathbf{f}(\mathbf{u}) = 0$$

$$\mathbf{u} = (u_1, \dots, u_m)^T$$

$$\mathbf{f}(\mathbf{u}) = (\mathbf{f}_1(\mathbf{u}), \dots, \mathbf{f}_d(\mathbf{u}))$$

Euler Equations for Compressible Fluids

$$\begin{cases} \partial_t \rho + \nabla_x \cdot (\rho \mathbf{v}) = 0 & \text{(conservation of mass)} \\ \partial_t (\rho \mathbf{v}) + \nabla_x \cdot (\rho \mathbf{v} \otimes \mathbf{v}) + \nabla_x p = 0 & \text{(conservation of momentum)} \\ \partial_t \left(\frac{1}{2} \rho |\mathbf{v}|^2 + \rho e \right) + \nabla_x \cdot \left(\left(\frac{1}{2} \rho |\mathbf{v}|^2 + \rho e + p \right) \mathbf{v} \right) = 0 & \text{(conservation of energy)} \end{cases}$$

Constitutive Relations: $p = p(\rho, e)$

- ρ – density, $\mathbf{v} = (v_1, v_2, v_3)^T$ – fluid velocity
- p – pressure, e – internal energy



Leonhard Euler



George Stokes

***Govern** the Flows when Convective Motions

Dominate Diffusion/Dispersion, ...

e.g. , shock waves in **Gases, Elastic Fluids, Shallow Water,**

Poisson, Challis, Stokes, Kelvin, Rayleigh, Airy, Earnshaw, Riemann, Rankine, Christoffel, Mach, Clausius, Kirchhoff, Gibbs, Hugoniot, Duhem, Hadamard, Jouguet, Zampfen, Weber, Taylor, Becker, Bethe, Weyl, von Neumann, Courant, Friedrichs,

Euler Equations for Potential Flow

$$\begin{cases} \partial_t \rho + \nabla_{\mathbf{x}} \cdot (\rho \nabla_{\mathbf{x}} \Phi) = 0, & \text{(Conservation of mass)} \\ \partial_t \Phi + \frac{1}{2} |\nabla_{\mathbf{x}} \Phi|^2 + \frac{\rho^{\gamma-1}}{\gamma-1} = \frac{B_\infty}{\gamma-1} := \frac{\rho_\infty^{\gamma-1} + \frac{\gamma-1}{2} u_\infty^2}{\gamma-1}, & \text{(Bernoulli's law)} \end{cases}$$

for $\gamma > 1$ or, equivalently,

$$\partial_t \rho(\partial_t \Phi, \nabla_{\mathbf{x}} \Phi, B_\infty) + \nabla_{\mathbf{x}} \cdot (\rho(\partial_t \Phi, \nabla_{\mathbf{x}} \Phi, B_\infty) \nabla_{\mathbf{x}} \Phi) = 0,$$

with

$$\rho(\partial_t \Phi, \nabla_{\mathbf{x}} \Phi, B_\infty) = \left(B_\infty - (\gamma - 1) \left(\partial_t \Phi + \frac{1}{2} |\nabla_{\mathbf{x}} \Phi|^2 \right) \right)^{\frac{1}{\gamma-1}}.$$

- Aerodynamics/Gas Dynamics: Fundamental PDE
- The potential flow equations and the full Euler equations coincide or are close each other in many important physical situations

J. Hadamard: Leçons sur la Propagation des Ondes,


Hermann: Paris 1903

Conservation Laws and Einstein Equations

$$G_{\mu\nu} = 8\pi T_{\mu\nu}$$

$T_{\mu\nu}$ – Stress-energy tensor (Energy-momentum tensor)

$G_{\mu\nu}$ – Einstein tensor (Function of the metric)



**These equations, with the geodesic equation,
form the core of the mathematical formulation
of General Relativity**

Structure of the Einstein Equations
 \implies **Conservation Laws of
Energy and Momentum:**

$$\nabla_b T^{ab} = T^{ab}_{;b} = 0$$

CALCULUS OF VARIATIONS

A Field of Mathematics that deals with extremizing functionals, as opposed to ordinary calculus which deals with functions:

$$I[w] = \int_{\Omega} L(\nabla_x w(x), w(x), x) dx$$

- Energy or Action Functionals in Physics/Engineering/industry....
- Distance/Metric Functional in Optics (light),
Geometry (geodesics, minimal surfaces, ...),
- Cost Functionals in Optimization (controls, games, image processing, design, finance, transportation, ...), ...

POINT: Seek a Minimizer or Critical Point **u** of **I[·]:**

$$I'[u]=0$$

Great Progress has been made in the recent four decades...

Conservation Laws and Calculus of Variations

□ Systems of Euler-Lagrange Equations

□ Noether's Theorem:

Any Differentiable Symmetry of
the Action of a Physical System Has
a Corresponding **Conservation Law**.

Any Invariance of the Variational
Integral $I[w]$ leads to a corresponding
Conservation Law for the critical
point of $I[\cdot]$



Hyperbolicity

$$\partial_t \mathbf{u} + \nabla \cdot \mathbf{f}(\mathbf{u}) = 0, \quad \mathbf{u} \in \mathbb{R}^m, \quad \mathbf{x} \in \mathbb{R}^d$$

Plane Wave Solutions: $\mathbf{u}(t, \mathbf{x}) = \mathbf{w}(t, \boldsymbol{\omega} \cdot \mathbf{x})$

$\mathbf{w}(t, \boldsymbol{\xi})$ is determined by: $\partial_t \mathbf{w} + (\nabla_{\mathbf{w}} \mathbf{f}(\mathbf{w}) \cdot \boldsymbol{\omega}) \partial_{\boldsymbol{\xi}} \mathbf{w} = 0$

?? Existence of stable plane wave solutions ??

Hyperbolicity in D : For any $\boldsymbol{\omega} \in S^{d-1}$, $\mathbf{u} \in D$,

$$(\nabla_{\mathbf{u}} \mathbf{f}(\mathbf{u}) \cdot \boldsymbol{\omega})_{m \times m} \mathbf{r}_j(\mathbf{u}, \boldsymbol{\omega}) = \lambda_j(\mathbf{u}, \boldsymbol{\omega}) \mathbf{r}_j(\mathbf{u}, \boldsymbol{\omega}), \quad 1 \leq j \leq m$$

$\lambda_j(\mathbf{u}, \boldsymbol{\omega})$ are real

Main Features:

Finiteness of Propagation Speeds;

Discontinuities of Solutions,

Well-Posedness: Existence, Uniqueness, Stability, ...

Hadamard's Example: $\mathbf{u} \in \mathbb{R}^2$, $x \in \mathbb{R}$

Cauchy Problem:
$$\begin{cases} \partial_t \mathbf{u} + \partial_x \mathbf{f}(\mathbf{u}) = 0 \\ \mathbf{u}|_{t=0} = \begin{pmatrix} \frac{\sin(nx)}{n} \\ 0 \end{pmatrix} \longrightarrow \begin{pmatrix} 0 \\ 0 \end{pmatrix} \text{ as } n \rightarrow \infty. \end{cases}$$

$$\mathbf{u} = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix}, \quad \mathbf{f}(\mathbf{u}) = \begin{pmatrix} u_2 \\ -u_1 \end{pmatrix}, \quad \nabla \mathbf{f}(\mathbf{u}) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \lambda_j = (-1)^j \sqrt{-1}$$

Well-Posedness: Expect that **the Limit Solution** $\mathbf{u}^0(t, x) \equiv 0$

Fixed $n > 0$, the Unique Solution:

$$\mathbf{u}^n(t, x) = \left(\frac{\sin(nx)(e^{nt} + e^{-nt})}{2n}, \frac{\cos(nx)(e^{nt} - e^{-nt})}{2n} \right)^\top$$

$$\overline{\lim}_{n \rightarrow \infty} |\mathbf{u}^n(t, x) - \mathbf{u}^0(t, x)| = \infty \quad \text{for } t > 0 \text{ for any reasonable topology}$$

***Small Changes in the Data \implies Large Changes in the Solutions**

***Mapping: "Data Space" \rightarrow "Solutions Space" Is Not Continuous**

UNSTABLE!!

Scalar Conservation Laws

$$\partial_t u + \nabla \cdot \mathbf{f}(u) = 0, \quad u \in \mathbb{R}, \mathbf{x} \in \mathbb{R}^d$$

$$\mathbf{f} : \mathbb{R} \rightarrow \mathbb{R}^d$$

Then

$$\lambda(u, \omega) = \mathbf{f}'(u) \cdot \omega, \quad r(u, \omega) \equiv 1$$

\implies **Scalar conservation laws
are always hyperbolic**

*** This is not the case for Systems,
or High-Order Equations**