

Donaldson–Thomas theory of Calabi–Yau 3-folds, and generalizations

Lecture 2 of 3: Derived Algebraic Geometry, PTVV, and
Darboux Theorems

Dominic Joyce, Oxford University

'Modern Moduli Theory' graduate school, September 2017

Based on: Pantev–Toën–Vaquié–Vezzosi arXiv:1111.3209 and
papers arXiv:1305.6302 and arXiv:1312.0090, by Ben-Bassat,
Brav, Bussi, and Joyce

Plan of talk:

- 4 Derived Algebraic Geometry and PTVV
 - 4.1 Derived Algebraic Geometry for dummies
 - 4.2 Tangent and cotangent complexes
 - 4.3 Classical symplectic geometry
 - 4.4 PTVV's shifted symplectic geometry
- 5 A 'Darboux theorem' for shifted symplectic schemes
 - 5.1 The case of -1 -shifted symplectic derived schemes
 - 5.2 The case of -2 -shifted symplectic derived schemes

4. Derived Algebraic Geometry and PTVV

4.1. Derived Algebraic Geometry for dummies

Let \mathbb{K} be an algebraically closed field of characteristic zero, e.g. $\mathbb{K} = \mathbb{C}$. Work in the context of Toën and Vezzosi's theory of *Derived Algebraic Geometry (DAG)*. This gives ∞ -categories of *derived \mathbb{K} -schemes* $\mathbf{dSch}_{\mathbb{K}}$ and *derived stacks* $\mathbf{dSt}_{\mathbb{K}}$. In this talk, for simplicity, we will mostly discuss derived schemes, though the results also extend to derived stacks.

This is a very technical subject. It is not easy to motivate DAG, or even to say properly what a derived scheme is, in an elementary talk. So I will lie a little bit.

What is a derived scheme?

\mathbb{K} -schemes in classical algebraic geometry are geometric spaces X which can be covered by Zariski open sets $Y \subseteq X$ with $Y \cong \text{Spec } A$ for A a commutative \mathbb{K} -algebra. General \mathbb{K} -schemes are very singular, but *smooth \mathbb{K} -schemes* X are very like smooth manifolds over \mathbb{K} , many differential geometric ideas like cotangent bundles TX , T^*X work nicely for them.

Think of a derived \mathbb{K} -scheme \mathbf{X} as a geometric space which can be covered by Zariski open sets $\mathbf{Y} \subseteq \mathbf{X}$ with $\mathbf{Y} \simeq \text{Spec } A^\bullet$ for $A^\bullet = (A, d)$ a commutative differential graded algebra (cdga) over \mathbb{K} , in degrees ≤ 0 .

We require \mathbf{X} to be *locally finitely presented*, that is, we can take the A^\bullet to be finitely presented, a strong condition.

Why derived algebraic geometry?

One reason derived algebraic geometry can be a powerful tool, is the combination of two facts:

- (A) Many algebro-geometric spaces one wants to study, such as moduli spaces of coherent sheaves, or complexes, or representations, etc., which in classical algebraic geometry may be very singular, also have an incarnation as (locally finitely presented) derived schemes (or derived stacks).
- (B) Within the framework of DAG, one can treat (locally finitely presented) derived schemes or stacks very much like smooth, nonsingular objects (Kontsevich's 'hidden smoothness philosophy'). Some nice things work in the derived world, which do not work in the classical world.

4.2. Tangent and cotangent complexes

In going from classical to derived geometry, we always replace vector bundles, sheaves, representations, . . . , by *complexes* of vector bundles, A classical smooth \mathbb{K} -scheme X has a tangent bundle TX and dual cotangent bundle T^*X , which are vector bundles on X , of rank the dimension $\dim X \in \mathbb{N}$.

Similarly, a derived \mathbb{K} -scheme \mathbf{X} has a *tangent complex* $\mathbb{T}_{\mathbf{X}}$ and a dual *cotangent complex* $\mathbb{L}_{\mathbf{X}}$, which are perfect complexes of coherent sheaves on \mathbf{X} , of rank the virtual dimension $\mathrm{vdim} \mathbf{X} \in \mathbb{Z}$. A complex \mathcal{E}^\bullet on \mathbf{X} is called *perfect in the interval* $[a, b]$ if locally on \mathbf{X} it is quasi-isomorphic to a complex

$\cdots 0 \rightarrow E_a \rightarrow E_{a+1} \rightarrow \cdots \rightarrow E_b \rightarrow 0 \rightarrow \cdots$, with E_i a vector bundle in position i . For \mathbf{X} a derived scheme, $\mathbb{T}_{\mathbf{X}}$ is perfect in $[0, \infty)$ and $\mathbb{L}_{\mathbf{X}}$ perfect in $(-\infty, 0]$; for \mathbf{X} a derived stack, $\mathbb{T}_{\mathbf{X}}$ is perfect in $[-1, \infty)$ and $\mathbb{L}_{\mathbf{X}}$ perfect in $(-\infty, 1]$.

Tangent complexes of moduli stacks

Suppose X is a smooth projective scheme, and \mathcal{M} is a derived moduli stack of coherent sheaves E on X . Then for each point $[E]$ in \mathcal{M} and each $i \in \mathbb{Z}$ we have natural isomorphisms

$$H^i(\mathbb{T}_{\mathcal{M}}|_{[E]}) \cong \mathrm{Ext}^{i-1}(E, E). \quad (1)$$

In effect, the derived stack \mathcal{M} remembers the entire deformation theory of sheaves on X . In contrast, if $\mathcal{M} = t_0(\mathcal{M})$ is the corresponding classical moduli scheme, (1) holds when $i \leq 1$ only. This shows that derived structure on a moduli scheme/stack can remember useful information forgotten by the classical moduli scheme/stack, e.g. the Ext groups $\mathrm{Ext}^i(E, E)$ for $i \geq 2$. If X has dimension n then (1) implies that $H^i(\mathbb{T}_{\mathcal{M}}|_{[E]}) = 0$ for $i \geq n$, so $\mathbb{T}_{\mathcal{M}}$ is perfect in $[-1, n-1]$.

Quasi-smooth derived schemes and virtual cycles

A derived scheme \mathbf{X} is called *quasi-smooth* if $\mathbb{T}_{\mathbf{X}}$ is perfect in $[0, 1]$, or equivalently $\mathbb{L}_{\mathbf{X}}$ is perfect in $[-1, 0]$.

A proper quasi-smooth derived scheme \mathbf{X} has a *virtual cycle* $[\mathbf{X}]_{\mathrm{virt}}$ in the Chow homology $A_*(X)$, where $X = t_0(\mathbf{X})$ is the classical truncation. This is because the natural morphism $\mathbb{L}_{\mathbf{X}}|_X \rightarrow \mathbb{L}_X$ induced by the inclusion $X \hookrightarrow \mathbf{X}$ is a 'perfect obstruction theory' in the sense of Behrend and Fantechi.

Most theories of invariants in algebraic geometry – e.g.

Gromov–Witten invariants, Mochizuki invariants counting sheaves on surfaces, Donaldson–Thomas invariants – can be traced back to the existence of quasi-smooth derived moduli schemes.

For an (ordinary) derived moduli scheme \mathcal{M} of coherent sheaves E on X to be quasi-smooth, we need $\mathrm{Ext}^i(E, E) = 0$ for $i \geq 3$. This is automatic if $\dim X \leq 2$. For Calabi–Yau 3-folds X , you would expect a problem with $\mathrm{Ext}^3(E, E) \neq 0$, but stable sheaves E with fixed determinant have trace-free Ext groups $\mathrm{Ext}^3(E, E)_0 = 0$.

An example of nice behaviour in the derived world

Here is an example of the 'hidden smoothness philosophy'. Suppose we have a Cartesian square of smooth \mathbb{K} -schemes (or indeed, smooth manifolds)

$$\begin{array}{ccc} W & \xrightarrow{\quad f \quad} & Y \\ \downarrow e & & \downarrow h \\ X & \xrightarrow{\quad g \quad} & Z, \end{array}$$

with g, h transverse. Then we have an exact sequence of vector bundles on W , which we can use to compute TW :

$$0 \rightarrow TW \xrightarrow{T_e \oplus T_f} e^*(TX) \oplus f^*(TY) \xrightarrow{e^*(Tg) \oplus -f^*(Th)} (g \circ e)^*(TZ) \rightarrow 0.$$

Similarly, if we have a homotopy Cartesian square of derived \mathbb{K} -schemes

$$\begin{array}{ccc} W & \xrightarrow{\quad f \quad} & Y \\ \downarrow e & & \downarrow h \\ X & \xrightarrow{\quad g \quad} & Z, \end{array}$$

with no transversality, we have a distinguished triangle on W

$$\mathbb{T}_W \xrightarrow{T_e \oplus T_f} e^*(\mathbb{T}_X) \oplus f^*(\mathbb{T}_Y) \xrightarrow{e^*(Tg) \oplus -f^*(Th)} (g \circ e)^*(\mathbb{T}_Z) \rightarrow \mathbb{T}_W[+1],$$

which we can use to compute \mathbb{T}_W . This is false for classical schemes. So, derived schemes with arbitrary morphisms, have good behaviour analogous to smooth classical schemes with transverse morphisms, and are better behaved than classical schemes.

4.3. Classical symplectic geometry

Let M be a smooth manifold. Then M has a tangent bundle and cotangent bundle T^*M . We have k -forms $\omega \in C^\infty(\Lambda^k T^*M)$, and the de Rham differential $d_{dR} : C^\infty(\Lambda^k T^*M) \rightarrow C^\infty(\Lambda^{k+1} T^*M)$. A k -form ω is *closed* if $d_{dR}\omega = 0$.

A 2-form ω on M is *nondegenerate* if $\omega \cdot : TM \rightarrow T^*M$ is an isomorphism. This is possible only if $\dim M = 2n$ for $n \geq 0$. A *symplectic structure* is a closed, nondegenerate 2-form ω on M . Symplectic geometry is the study of symplectic manifolds (M, ω) . A *Lagrangian* in (M, ω) is a submanifold $i : L \rightarrow M$ such that $\dim L = n$ and $i^*(\omega) = 0$.

4.4. PTVV's shifted symplectic geometry

Pantev, Toën, Vaquié and Vezzosi (arXiv:1111.3209) defined a version of symplectic geometry in the derived world.

Let \mathbf{X} be a derived \mathbb{K} -scheme. The cotangent complex $\mathbb{L}_{\mathbf{X}}$ has exterior powers $\Lambda^p \mathbb{L}_{\mathbf{X}}$. The *de Rham differential* $d_{dR} : \Lambda^p \mathbb{L}_{\mathbf{X}} \rightarrow \Lambda^{p+1} \mathbb{L}_{\mathbf{X}}$ is a morphism of complexes. Each $\Lambda^p \mathbb{L}_{\mathbf{X}}$ is a complex, so has an internal differential $d : (\Lambda^p \mathbb{L}_{\mathbf{X}})^k \rightarrow (\Lambda^p \mathbb{L}_{\mathbf{X}})^{k+1}$. We have $d^2 = d_{dR}^2 = d \circ d_{dR} + d_{dR} \circ d = 0$.

A *p-form of degree k* on \mathbf{X} for $k \in \mathbb{Z}$ is an element $[(\omega^0)]$ of $H^k(\Lambda^p \mathbb{L}_{\mathbf{X}}, d)$. A *closed p-form of degree k* on \mathbf{X} is an element

$$[(\omega^0, \omega^1, \dots)] \in H^k\left(\bigoplus_{i=0}^{\infty} \Lambda^{p+i} \mathbb{L}_{\mathbf{X}}[i], d + d_{dR}\right).$$

There is a projection $\pi : [(\omega^0, \omega^1, \dots)] \mapsto [\omega^0]$ from closed p -forms $[(\omega^0, \omega^1, \dots)]$ of degree k to p -forms $[\omega^0]$ of degree k .

Nondegenerate 2-forms and symplectic structures

Let $[\omega^0]$ be a 2-form of degree k on \mathbf{X} . Then $[\omega^0]$ induces a morphism $\omega^0 : \mathbb{T}_{\mathbf{X}} \rightarrow \mathbb{L}_{\mathbf{X}}[k]$, where $\mathbb{T}_{\mathbf{X}} = \mathbb{L}_{\mathbf{X}}^{\vee}$ is the tangent complex of \mathbf{X} . We call $[\omega^0]$ *nondegenerate* if $\omega^0 : \mathbb{T}_{\mathbf{X}} \rightarrow \mathbb{L}_{\mathbf{X}}[k]$ is a quasi-isomorphism.

If \mathbf{X} is a derived scheme then the complex $\mathbb{L}_{\mathbf{X}}$ lives in degrees $(-\infty, 0]$ and $\mathbb{T}_{\mathbf{X}}$ in degrees $[0, \infty)$. So $\omega^0 : \mathbb{T}_{\mathbf{X}} \rightarrow \mathbb{L}_{\mathbf{X}}[k]$ can be a quasi-isomorphism only if $k \leq 0$, and then $\mathbb{L}_{\mathbf{X}}$ lives in degrees $[k, 0]$ and $\mathbb{T}_{\mathbf{X}}$ in degrees $[0, -k]$. If $k = 0$ then \mathbf{X} is a smooth classical \mathbb{K} -scheme, and if $k = -1$ then \mathbf{X} is quasi-smooth.

A closed 2-form $\omega = [(\omega^0, \omega^1, \dots)]$ of degree k on \mathbf{X} is called a *k-shifted symplectic structure* if $[\omega^0] = \pi(\omega)$ is nondegenerate.

Although the details are complex, PTVV are following a simple recipe for translating some piece of geometry from smooth manifolds/smooth classical schemes to derived schemes:

- (i) replace manifolds/smooth schemes X by derived schemes \mathbf{X} .
- (ii) replace vector bundles $TX, T^*X, \Lambda^p T^*X, \dots$ by complexes $\mathbb{T}_{\mathbf{X}}, \mathbb{L}_{\mathbf{X}}, \Lambda^p \mathbb{L}_{\mathbf{X}}, \dots$.
- (iii) replace sections of $TX, T^*X, \Lambda^p T^*X, \dots$ by cohomology classes of the complexes $\mathbb{T}_{\mathbf{X}}, \mathbb{L}_{\mathbf{X}}, \Lambda^p \mathbb{L}_{\mathbf{X}}, \dots$, in degree $k \in \mathbb{Z}$.
- (iv) replace isomorphisms of vector bundles by quasi-isomorphisms of complexes.

Note that in (iii), we can specify the degree $k \in \mathbb{Z}$ of the cohomology class (e.g. $[\omega] \in H^k(\Lambda^p \mathbb{L}_{\mathbf{X}})$), which doesn't happen at the classical level.

Calabi–Yau moduli schemes and moduli stacks

PTVV prove that if Y is a Calabi–Yau m -fold over \mathbb{K} and \mathcal{M} is a derived moduli scheme or stack of (complexes of) coherent sheaves on Y , then \mathcal{M} has a $(2 - m)$ -shifted symplectic structure ω . This suggests applications — lots of interesting geometry concerns Calabi–Yau moduli schemes, e.g. Donaldson–Thomas theory. We can understand the associated nondegenerate 2-form $[\omega^0]$ in terms of *Serre duality*. At a point $[E] \in \mathcal{M}$, we have $h^i(\mathbb{T}_{\mathcal{M}})|_{[E]} \cong \text{Ext}^{i-1}(E, E)$ and $h^i(\mathbb{L}_{\mathcal{M}})|_{[E]} \cong \text{Ext}^{1-i}(E, E)^*$. The Calabi–Yau condition gives $\text{Ext}^i(E, E) \cong \text{Ext}^{m-i}(E, E)^*$, which corresponds to $h^{i+1}(\mathbb{T}_{\mathcal{M}})|_{[E]} \cong h^{i+1}(\mathbb{L}_{\mathcal{M}}[2 - m])|_{[E]}$. This is the cohomology at $[E]$ of the quasi-isomorphism $\omega^0 : \mathbb{T}_{\mathcal{M}} \rightarrow \mathbb{L}_{\mathcal{M}}[2 - m]$.

Lagrangians and Lagrangian intersections

Let (\mathbf{X}, ω) be a k -shifted symplectic derived scheme or stack. Then Pantev et al. define a notion of *Lagrangian* \mathbf{L} in (\mathbf{X}, ω) , which is a morphism $i : \mathbf{L} \rightarrow \mathbf{X}$ of derived schemes or stacks together with a homotopy $i^*(\omega) \sim 0$ satisfying a nondegeneracy condition, implying that $\mathbb{T}_{\mathbf{L}} \simeq \mathbb{L}_{\mathbf{L}/\mathbf{X}}[k - 1]$. If \mathbf{L}, \mathbf{M} are Lagrangians in (\mathbf{X}, ω) , then the fibre product $\mathbf{L} \times_{\mathbf{X}} \mathbf{M}$ has a natural $(k - 1)$ -shifted symplectic structure. If (S, ω) is a classical smooth symplectic scheme, then it is a 0-shifted symplectic derived scheme in the sense of PTVV, and if $L, M \subset S$ are classical smooth Lagrangian subschemes, then they are Lagrangians in the sense of PTVV. Therefore the (derived) Lagrangian intersection $L \cap M = L \times_S M$ is a -1 -shifted symplectic derived scheme.

Summary of the story so far

- Derived schemes behave better than classical schemes in some ways – they are analogous to smooth schemes, or manifolds. So, we can extend stories in smooth geometry to derived schemes. This introduces an extra degree $k \in \mathbb{Z}$.
- PTVV define a version of (' k -shifted') symplectic geometry for derived schemes. This is a new geometric structure.
- 0-shifted symplectic derived schemes are just classical smooth symplectic schemes.
- Calabi–Yau m -fold moduli schemes and stacks are $(2 - m)$ -shifted symplectic. This gives a *new geometric structure* on Calabi–Yau moduli spaces – relevant to Donaldson–Thomas theory and its generalizations.
- One can go from k -shifted symplectic to $(k - 1)$ -shifted symplectic by taking intersections of Lagrangians.

5. A 'Darboux theorem' for shifted symplectic schemes

Theorem 5.1 (Brav, Bussi and Joyce arXiv:1305.6302)

Suppose (\mathbf{X}, ω) is a k -shifted symplectic derived \mathbb{K} -scheme for $k < 0$. If $k \not\equiv 2 \pmod{4}$, then each $x \in \mathbf{X}$ admits a Zariski open neighbourhood $\mathbf{Y} \subseteq \mathbf{X}$ with $\mathbf{Y} \simeq \text{Spec}(A, d)$ for (A, d) an explicit cdga generated by graded variables x_j^{-i}, y_j^{k+i} for $0 \leq i \leq -k/2$, and $\omega|_{\mathbf{Y}} = [(\omega^0, 0, 0, \dots)]$ where x_j^l, y_j^l have degree l , and

$$\omega^0 = \sum_{i=0}^{\lfloor -k/2 \rfloor} \sum_{j=1}^{m_i} d_{dR} y_j^{k+i} d_{dR} x_j^{-i}.$$

Also the differential d in (A, d) is given by Poisson bracket with a Hamiltonian H in A of degree $k + 1$.

If $k \equiv 2 \pmod{4}$, we have two statements, one étale local with ω^0 standard, and one Zariski local with the components of ω^0 in the degree $k/2$ variables depending on some invertible functions.

Sketch of the proof of Theorem 5.1

Suppose (\mathbf{X}, ω) is a k -shifted symplectic derived \mathbb{K} -scheme for $k < 0$, and $x \in \mathbf{X}$. Then $\mathbb{L}_{\mathbf{X}}$ lives in degrees $[k, 0]$. We first show that we can build Zariski open $x \in \mathbf{Y} \subseteq \mathbf{X}$ with $\mathbf{Y} \simeq \text{Spec}(A, d)$, for $A = \bigoplus_{i \leq 0} A^i$, d a cdga over \mathbb{K} with A^0 a smooth \mathbb{K} -algebra, and such that A is freely generated over A^0 by graded variables x_j^{-i}, y_j^{k+i} in degrees $-1, -2, \dots, k$. We take $\dim A^0$ and the number of x_j^{-i}, y_j^{k+i} to be minimal at x .

Using theorems about periodic cyclic cohomology, we show that on $\mathbf{Y} \simeq \text{Spec}(A, d)$ we can write $\omega|_{\mathbf{Y}} = [(\omega^0, 0, 0, \dots)]$, for ω^0 a 2-form of degree k with $d\omega^0 = d_{dR}\omega^0 = 0$. Minimality at x implies ω^0 is strictly nondegenerate near x , so we can change variables to write $\omega^0 = \sum_{i,j} d_{dR}y_j^{k+i} d_{dR}x_j^{-i}$. Finally, we show d in (A, d) is a symplectic vector field, which integrates to a Hamiltonian H .

A 'Darboux Theorem' for atlases of derived stacks

Here is an extension of Theorem 5.1 to derived stacks.

Theorem 5.2 (Ben-Bassat, Bussi, Brav, Joyce, arXiv:1312.0090)

Let $(\mathbf{X}, \omega_{\mathbf{X}})$ be a k -shifted symplectic derived Artin stack for $k < 0$, and $p \in \mathbf{X}$. Then there exist 'standard form' affine derived schemes $\mathbf{U} = \text{Spec } A$, $\mathbf{V} = \text{Spec } B$, points $u \in \mathbf{U}$, $v \in \mathbf{V}$ with A, B minimal at u, v , morphisms $\varphi : \mathbf{U} \rightarrow \mathbf{X}$ and $\mathbf{i} : \mathbf{U} \rightarrow \mathbf{V}$ with $\varphi(u) = p$, $\mathbf{i}(u) = v$, such that φ is smooth of relative dimension $\dim H^1(\mathbb{L}_{\mathbf{X}}|_p)$, and $t_0(\mathbf{i}) : t_0(\mathbf{U}) \rightarrow t_0(\mathbf{V})$ is an isomorphism on classical schemes, and $\mathbb{L}_{\mathbf{U}/\mathbf{V}} \simeq \mathbb{T}_{\mathbf{U}/\mathbf{X}}[1 - k]$, and a 'Darboux form' k -shifted symplectic form ω_B on $\mathbf{V} = \text{Spec } B$ such that $\mathbf{i}^(\omega_B) \sim \varphi^*(\omega_{\mathbf{X}})$ in k -shifted closed 2-forms on \mathbf{U} .*

5.1. The case of -1 -shifted symplectic derived schemes

When $k = -1$ the Hamiltonian H in the theorem has degree 0.
Then Theorem 5.1 reduces to:

Corollary 5.3

Suppose (\mathbf{X}, ω) is a -1 -shifted symplectic derived \mathbb{K} -scheme. Then (\mathbf{X}, ω) is Zariski locally equivalent to a derived critical locus $\text{Crit}(H : U \rightarrow \mathbb{A}^1)$, for U a smooth classical \mathbb{K} -scheme and $H : U \rightarrow \mathbb{A}^1$ a regular function. Hence, the underlying classical \mathbb{K} -scheme $X = t_0(\mathbf{X})$ is Zariski locally isomorphic to a classical critical locus $\text{Crit}(H : U \rightarrow \mathbb{A}^1)$.

Combining this with results of Pantev et al. from §4 gives interesting consequences in classical algebraic geometry:

Corollary 5.4

Let Y be a Calabi–Yau 3-fold over \mathbb{K} and \mathcal{M} a classical moduli \mathbb{K} -scheme of coherent sheaves, or complexes of coherent sheaves, on Y . Then \mathcal{M} is Zariski locally isomorphic to the critical locus $\text{Crit}(H : U \rightarrow \mathbb{A}^1)$ of a regular function on a smooth \mathbb{K} -scheme.

Here we note that $\mathcal{M} = t_0(\mathcal{M})$ for \mathcal{M} the corresponding derived moduli scheme, which is -1 -shifted symplectic by PTVV.

A complex analytic analogue of this for moduli of coherent sheaves was proved using gauge theory by Joyce and Song arXiv:0810.5645 (Lecture 1, key idea 3), and for moduli of complexes was claimed by Behrend and Getzler.

Note that the proof of the corollary is wholly algebro-geometric.

Outlook for generalizations of Donaldson–Thomas theory

We now know that 3-Calabi–Yau moduli spaces are locally modelled on critical loci, and we have nice geometric structures encoding this (-1 -shifted symplectic structures).

There is some interesting geometry associated with critical loci:

- Perverse sheaves of vanishing cycles.
- Motivic Milnor fibres.
- Categories of matrix factorizations.

It seems natural to try and construct global structures on 3-Calabi–Yau moduli spaces, which are locally modelled on perverse vanishing cycles, motivic Milnor fibres, or matrix factorizations. This leads to questions of *categorification* of Donaldson–Thomas theory, and *motivic Donaldson–Thomas invariants*.

5.2. The case of -2 -shifted symplectic derived schemes

Let (\mathbf{X}, ω) be a -2 -shifted symplectic derived \mathbb{K} -scheme. Then the Zariski local models for (\mathbf{X}, ω) given by the Theorem 5.1 depend on the following data:

- A smooth \mathbb{K} -scheme U
- An algebraic vector bundle $E \rightarrow U$
- A section $s \in H^0(E)$
- A nondegenerate quadratic form Q on E with $Q(s, s) = 0$.

The underlying classical \mathbb{K} -scheme X of \mathbf{X} is locally $s^{-1}(0) \subset U$.

The virtual dimension of \mathbf{X} is $\mathrm{vdim}_{\mathbb{K}} \mathbf{X} = 2 \dim_{\mathbb{K}} U - \mathrm{rank}_{\mathbb{K}} E$.

The cotangent complex $\mathbb{L}_{\mathbf{X}}|_X$ of \mathbf{X} is locally given by

$$\left[\begin{array}{c} TU \\ -2 \end{array} \Big|_{s^{-1}(0)} \xrightarrow{Q \circ ds} \begin{array}{c} E^* \\ -1 \end{array} \Big|_{s^{-1}(0)} \xrightarrow{ds} \begin{array}{c} T^*U \\ 0 \end{array} \Big|_{s^{-1}(0)} \right].$$

This is the local model for (derived) moduli schemes of (simple) coherent sheaves F on a Calabi–Yau 4-fold Y . Think of U as an étale open neighbourhood of 0 in $\mathrm{Ext}^1(F, F)$, and $E \rightarrow U$ as a trivial vector bundle with fibre $\mathrm{Ext}^2(F, F)$, and Q as the nondegenerate quadratic form on $\mathrm{Ext}^2(F, F)$

$$\mathrm{Ext}^2(F, F) \times \mathrm{Ext}^2(F, F) \xrightarrow{\wedge} \mathrm{Ext}^4(F, F) \xrightarrow{\text{Serre duality}} \mathrm{Ext}^0(F, F)^* \xrightarrow{\mathrm{id}_F^*} \mathbb{K},$$

and s as a Kuranishi map $s : \mathrm{Ext}^1(E, E) \supseteq U \rightarrow \mathrm{Ext}^2(E, E)$. The special thing the theorem tells us is that we can choose U, E, s, Q with $Q(s, s) = 0$, rather than just $Q(s, s) = 0 \bmod s^3$, for instance. Borisov–Joyce arXiv:1504.00690 use these local models to construct virtual classes in homology for proper, oriented -2 -shifted symplectic derived \mathbb{C} -schemes. Surprisingly, the virtual class has half the expected dimension. This will lead to a theory of Donaldson–Thomas style invariants for Calabi–Yau 4-folds (in progress, joint work with Yalong Cao).