



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Journal of  
Combinatorial  
Theory

Series B

Journal of Combinatorial Theory, Series B 92 (2004) 163–175

<http://www.elsevier.com/locate/jctb>

# Stability theorems for cancellative hypergraphs

Peter Keevash<sup>a</sup> and Dhruv Mubayi<sup>b</sup>

<sup>a</sup>*Department of Mathematics, Princeton University, Princeton, NJ 08540, USA*

<sup>b</sup>*Department of Mathematics, Statistics, and Computer Science, University of Illinois at Chicago,  
851 S. Morgan Street, Chicago, IL 60607, USA*

Received 11 December 2002

Available online 7 July 2004

---

## Abstract

A *cancellative* hypergraph has no three edges  $A, B, C$  with  $A\Delta B \subset C$ . We give a new short proof of an old result of Bollobás, which states that the maximum size of a cancellative triple system is achieved by the balanced complete tripartite 3-graph.

One of the two forbidden subhypergraphs in a cancellative 3-graph is  $F_5 = \{abc, abd, cde\}$ . For  $n \geq 33$  we show that the maximum number of triples on  $n$  vertices containing no copy of  $F_5$  is also achieved by the balanced complete tripartite 3-graph. This strengthens a theorem of Frankl and Füredi, who proved it for  $n \geq 3000$ .

For both extremal results, we show that a 3-graph with almost as many edges as the extremal example is approximately tripartite. These stability theorems are analogous to the Simonovits stability theorem for graphs.

© 2004 Elsevier Inc. All rights reserved.

---

## 1. Introduction

Given an  $k$ -uniform hypergraph  $\mathcal{F}$ , the Turán number  $ex(n, \mathcal{F})$  of  $\mathcal{F}$  is the maximum number of edges in an  $k$ -uniform hypergraph on  $n$  vertices that does not contain a copy of  $\mathcal{F}$ . Determining these numbers is one of the central problems in extremal combinatorics. Much is known for ordinary graphs (the case  $k = 2$ ), going back to Turán [15], who solved the problem for complete graphs. The *Turán graph*  $T_r(n)$  is the complete  $r$ -partite graph on  $n$  vertices, which is ‘balanced’, i.e. no two part sizes differ by more than one. Write  $t_r(n)$  for the number of edges in  $T_r(n)$ . Then

---

*E-mail addresses:* keevash@math.princeton.edu (P. Keevash), mubayi@math.uic.edu (D. Mubayi).

Turán's theorem states that a graph  $G$  on  $n$  vertices containing no copy of  $K_{r+1}$  can have at most  $t_r(n)$  edges, and equality holds only when  $G = T_r(n)$ .

In contrast, for  $k > 2$ , the problem of finding the numbers  $ex(n, \mathcal{F})$  is notoriously difficult. Exact results on hypergraph Turán numbers are very rare (see [6,13] for surveys). It is even difficult to determine the *Turán density* of a  $k$ -graph  $\mathcal{F}$ , i.e.  $\pi(\mathcal{F}) = \lim_{n \rightarrow \infty} ex(n, \mathcal{F}) / \binom{n}{k}$ . Recently, a new technique was developed by de Caen and Füredi [4] to show that, when  $\mathcal{F}$  is the Fano plane,  $\pi(\mathcal{F}) = 3/4$ . Since then there has been considerable progress on Turán problems for 3-uniform hypergraphs. The second author and Rödl [10] developed this method to determine many new Turán densities, and the Turán number of the Fano plane was determined by the first author and Sudakov [8], and independently by Füredi and Simonovits [7], thus answering a conjecture of Sós.

In the 1960's, Katona posed the problem of determining the maximum number of triples in an  $n$  vertex 3-graph  $H$  such that no three distinct triples  $A, B, C$  have the property that  $B \Delta C \subset A$ . Here  $\Delta$  is the symmetric difference, so  $B \Delta C = (B - C) \cup (C - B)$ . Equivalently,  $H$  has the property that whenever  $A \cup B = A \cup C$ , we have  $B = C$ . Such 3-graphs are called *cancellative*.

For graphs, the condition  $B \Delta C \subset A$  can only occur if  $ABC$  form a triangle. Then the special case of Turán's theorem due to Mantel tells us that  $e(H) \leq t_2(n) = \lfloor n^2/4 \rfloor$ . This was the motivation for Katona's problem. Moving on to 3-graphs, we see that if  $B \neq C$  then  $B \Delta C \subset A$  can only occur when  $B$  and  $C$  have two points in common. This leads us to identify the two non-isomorphic configurations that are forbidden in a cancellative 3-graph:  $F_4 = \{abc, abd, bcd\}$  and  $F_5 = \{abc, abd, cde\}$ . (Note that  $F_4$  is  $K_4^{(3)}$  with an edge removed.)

**Definition 1.1.**  $S(n)$  is the complete 3-partite 3-graph on  $n$  vertices with part sizes  $\lfloor n/3 \rfloor$ ,  $\lfloor (n+1)/3 \rfloor$ ,  $\lfloor (n+2)/3 \rfloor$ . Let  $s(n)$  be the number of edges in  $S(n)$ .

Katona conjectured that the unique cancellative 3-graph on  $n$  vertices with the maximum number of triples is  $S(n)$ , and this was proved by Bollobás [1] (see also [3]).

**Theorem 1.2** (Bollobás). *A cancellative 3-graph on  $n$  vertices has at most  $s(n)$  edges, with equality only for  $S(n)$ .*

Bollobás conjectured that a similar result holds for  $k$ -graphs when  $k \geq 4$ , namely that the unique cancellative  $k$ -graph on  $n$  vertices with the most edges is the complete  $k$ -partite  $k$ -graph with almost equal part sizes. Sidorenko [12] proved Bollobás' conjecture for  $k = 4$ , but Shearer [11] gave an example showing that the conjecture is false for  $k > 10$ .

As observed above, the case  $k = 3$  of Katona's problem asks for  $ex(n, \mathcal{F})$  where  $\mathcal{F} = \{F_4, F_5\}$ . Frankl and Füredi [5] sharpened Bollobás' theorem by proving the following:

**Theorem 1.3** (Frankl–Füredi). *Let  $n \geq 3000$ . Then  $ex(n, F_5) = s(n)$ .*

In [10] a new proof of an asymptotic version of Theorem 1.3 is given. In this paper, we refine those ideas to give a new proof of Bollobás’ result, Theorem 1.2 (see Section 3). We also extend the ideas of that proof to improve Theorem 1.3. Our approach has the advantages that it significantly lowers the minimum  $n$  for which Theorem 1.3 holds, and characterizes the extremal example.

**Theorem 1.4** (Section 4). *Let  $n \geq 33$ . Then  $ex(n, F_5) = s(n)$ , with equality only for  $S(n)$ .*

Using our method, we prove stability versions of both extremal results, showing that any 3-graph that is cancellative, or even just  $F_5$ -free, and has nearly  $s(n)$  edges, must be approximately tripartite. These results are analogous to the Simonovits stability theorem for graphs, which says that a  $K_{r+1}$ -free graph with nearly  $t_r(n)$  edges is approximately  $r$ -partite (we will state this precisely later). The strongest result is the following.

**Theorem 1.5** (Section 5). *For any  $\varepsilon > 0$  there exists  $\delta > 0$  such that if  $H$  is an  $F_5$ -free 3-graph on  $n$  vertices with at least  $(1 - \delta)s(n)$  edges, then there is a partition of the vertex set of  $H$  as  $V(H) = U_1 \cup U_2 \cup U_3$  so that all but at most  $\varepsilon n^3$  edges of  $H$  have one point in each  $U_i$ .*

As far as we know, these are among the first stability theorems for hypergraphs. As well as being of interest in their own right, it seems that stability theorems may play a valuable role in the development of Turán theory for hypergraphs. The program, established in [8,9], consists of first proving a stability result for the problem in question, and then refining it to prove a Turán result. It would be interesting to see if any other problems are susceptible to this line of attack.

## 2. Preliminaries

In a 3-graph  $H$  there are two natural analogies to the notion of ‘neighborhood’ in 2-graphs. The *link* of a vertex  $x \in V(H)$  is  $L(x) = \{\{a, b\} : abx \in E(H)\}$ . We can think of the link as a graph on  $V(H)$ . The *degree* of  $x$  is  $d(x) = |L(x)|$ , the number of edges containing  $x$ . Given a pair of vertices  $x, y$ , their *neighborhood* is  $N(x, y) = \{a : axy \in E(H)\}$ . We record some simple properties of these definitions in the following useful lemma. Call an edge-colored graph *rainbow* if no two edges have the same color. Given a graph  $G$  and a subset  $X$  of its vertex set we write  $G_X$  for the restriction of  $G$  to  $X$ , i.e. the edges are those edges of  $G$  with both endpoints in  $X$ .

**Lemma 2.1.** *Let  $H$  be a 3-graph, and suppose that  $xyz \in E(H)$ . Let  $W = V(H) - \{x, y, z\}$ .*

(i) If  $H$  contains no  $F_4$ , then  $L(x)$  contains no triangle.

(ii) If  $H$  contains no  $F_5$ , then  $L(x)_W, L(y)_W$  and  $L(z)_W$  are edge-disjoint graphs. If  $H$  is cancellative then  $L(x), L(y)$  and  $L(z)$  are edge-disjoint graphs.

When  $H$  contains no  $F_5$ , we think of  $G = (L(x) \cup L(y) \cup L(z))_W$  as a 3-edge-colored graph.

(iii) If  $H$  contains no  $F_5$ , then any triangle in  $G$  is either monochromatic or rainbow. If  $H$  is cancellative then all triangles in  $G$  are rainbow.

(iv) If  $H$  is cancellative and  $G$  contains a  $K_4$  then this  $K_4$  must be properly 3-edge-colored.

**Proof.** (i) If  $abc$  is a triangle in  $L(x)$ , then  $xab, xac, xbc$  forms a copy of  $F_4$ .

(ii) If  $a, b$  are in  $W$  with  $ab \in L(x) \cap L(y)$ , then  $xab, yab, xyz$  forms a copy of  $F_5$ . If  $ae \in L(x) \cap L(y)$ , then  $xaz, yaz, xyz$  forms a copy of  $F_4$ .

(iii) If  $abc$  is a triangle with  $ab$  and  $bc$  in  $L(x)$  and  $ac$  in  $L(y)$ , then  $xab, xbc, yac$  forms a copy of  $F_5$ . If  $H$  is also  $F_4$ -free there are no monochromatic triangles by (i).

(iv) Every triangle is rainbow, so no two incident edges have the same color. In a copy of  $K_4$ , this implies that every color appears at most twice. There are three colors and six edges, so equality holds.  $\square$

We conclude this section by deriving some recurrences for the number of edges in  $T_3(n)$  and  $S(n)$ . First we give the approximate formulae,

$$t_3(n) = \left(\frac{2}{3} + o(1)\right) \binom{n}{2}, \quad s(n) = \left(\frac{2}{9} + o(1)\right) \binom{n}{3}.$$

Now consider both as being defined on the same vertex set, with the same partition  $X, Y, Z$ . Let  $xyz$  be an edge of  $S(n)$ . By inclusion/exclusion the number of edges meeting the set  $\{x, y, z\}$  is

$$|L(x)| + |L(y)| + |L(z)| - |N(x, y)| - |N(y, z)| - |N(z, x)| + 1. \tag{1}$$

Note that the links  $L(x), L(y), L(z)$  partition the edges of  $T_3(n)$  so  $|L(x)| + |L(y)| + |L(z)| = t_3(n)$ . The sets  $N(x, y), N(y, z), N(x, z)$  partition the vertex set. Consequently, (1) is  $t_3(n) - n + 1$  and we get the identity

$$s(n) = s(n - 3) + t_3(n) - n + 1. \tag{2}$$

Finally, every vertex is adjacent to exactly two of  $x, y, z$  in  $T_3(n)$ , so

$$t_3(n) = t_3(n - 3) + 2n - 3. \tag{3}$$

### 3. A new proof of Bollobás’ theorem

In this section we will illustrate the application of the ideas of de Caen and Füredi to cancellative hypergraphs by giving a new proof of Theorem 1.2.

**Proof of Theorem 1.2.** Suppose that  $H$  is a cancellative 3-graph with  $s(n)$  edges. We will prove by induction on  $n$  that  $H$  is isomorphic to  $S(n)$ . Since  $S(n)$  is a maximal cancellative 3-graph, this will prove the theorem. The result is obvious for  $n \leq 4$ , so assume that  $n \geq 5$ . If there is any triple of vertices from  $H$  (not necessarily an edge) that meets at most  $t_3(n) - n$  edges, then delete its vertices to form  $H'$ . By (2),  $H'$  has more than  $s(n - 3)$  edges. Since  $H'$  is cancellative, the induction hypothesis applies to give a contradiction. We may therefore assume that every triple of  $H$  meets at least  $t_3(n) - n + 1$  edges of  $H$ .

Fix some edge  $e = x_1x_2x_3$ . We claim that at most  $t_3(n - 3)$  edges can meet  $e$  in exactly one point. For by Lemma 2.1 (ii) the links  $L(x_i)$  form a 3-colored graph  $G$ . Suppose for the sake of contradiction that  $G$  contains a  $K_4$  disjoint from  $e$ , which we denote  $K$ . Then we must have  $n \geq 7$ . By Lemma 2.1 (iv) each color meets  $K$  in a matching, so each vertex of  $K$  is incident to edges of all three colors. Then every pair from  $K \cup e$  belongs to some edge of  $H$ , so by Lemma 2.1 (ii), the seven link graphs of the vertices in  $K \cup e$  are pairwise edge-disjoint. There are  $\binom{7}{3} = 35$  triples in  $K \cup e$ , and for each the total size of the links of its vertices is at least the number of edges it meets, which is at least  $t_3(n) - n + 1$ . Each vertex belongs to  $\binom{6}{2} = 15$  triples, so by averaging the seven link graphs contain at least  $\lceil \frac{35}{15}(t_3(n) - n + 1) \rceil$  edges. This is greater than  $\binom{n}{2}$  for  $n \geq 7$ , a contradiction. We deduce that  $G$  contains no copy of  $K_4$ . Applying Turán’s theorem, we see that at most  $t_3(n - 3)$  edges can meet  $e$  in exactly one point.

Next note that the three sets  $N(x_1, x_2)$ ,  $N(x_2, x_3)$ ,  $N(x_3, x_1)$  are pairwise disjoint (or we find  $F_4$ ) so at most  $n - 3$  edges meet  $e$  in exactly two points. Therefore the number of edges meeting  $e$  (including  $e$  itself) is at most

$$t_3(n - 3) + (n - 3) + 1 = t_3(n) - n + 1$$

by (3). As argued before, we must have equality in all these computations. It follows that  $H' = H - \{x_1, x_2, x_3\}$  has  $s(n - 3)$  edges, so is  $S(n - 3)$  by induction hypothesis. Also,  $G$  restricted to  $V(H')$  has  $t_3(n - 3)$  edges and is  $K_4$ -free, so it must be  $T_3(n - 3)$ . Let  $(X_1, X_2, X_3)$  be its tripartition. By Lemma 2.1 (iii), all of its triangles are rainbow, and this is only possible when for each  $i, j$  all edges between  $X_i$  and  $X_j$  have the same color, which by relabelling we can take to be  $k$ , where  $\{1, 2, 3\} = \{i, j, k\}$ . With this notation  $N(x_i, x_j) \subset X_k \cup x_k$  as  $H$  is cancellative, and so  $N(x_i, x_j) = X_k \cup x_k$  by the above computations. Therefore  $\{X_i \cup \{x_i\}\}_{i=1}^3$  defines a tripartition of  $H$ , i.e.  $H$  is  $S_n$ .  $\square$

#### 4. An improvement of the Frankl–Füredi theorem

In this section we prove Theorem 1.4. First we prove an easy lemma on edge colorings of  $K_8$ .

**Lemma 4.1.** *Suppose that  $K_8$  is 4-edge-colored with no monochromatic  $K_4$  and no 2-colored triangle. Then each color class consists of two disjoint triangles and a single isolated edge.*

**Proof.** The hypothesis implies that in the graph formed by all edges of some fixed color, adjacency is an equivalence relation on the vertex set. Thus each color class is a union of complete subgraphs of size at most three, and therefore has at most seven edges. Since  $K_8$  has 28 edges, each color class has exactly seven edges, and the result follows.  $\square$

It will be useful to consider the following 5 vertex hypergraph, which has the property that each pair of its vertices belongs to an edge. Let  $F_{1,4}$  denote the 3-graph with vertex set  $v \cup U$ ,  $|U| = 4$ , and edge set  $\{vuu' : u, u' \in U\}$ .

**Proof of Theorem 1.4.** Suppose that  $H = (V, E)$  is an  $n$  vertex 3-graph that contains no  $F_5$ . We will prove by induction on  $n$  that  $e(H) \leq s(n) + h(n)$ , where  $h(n)$  is a function that is equal to zero for  $n \geq 32$  (this technique is sometimes called progressive induction). Consequently for  $n \geq 32$ , we have  $e(H) \leq s(n)$ . For  $n > 32$ , we will show that either  $H$  is cancellative, or  $e(H) < s(n)$ . By Theorem 1.2, this implies that if  $e(H) = s(n)$  for  $n > 32$ , then  $H$  is  $S(n)$ .

To define  $h(n)$  we first define a function  $g(n)$  by  $g(0) = g(1) = g(2) = g(3) = 0$ ,  $g(4) = 2$ , and

$$g(n) = \max\{g(n - 4) + t_7(n - 4) + 2(n - 4) + 3 - [s(n) - s(n - 4)],$$

$$g(n - 5) + \binom{n - 5}{2} + 4(n - 5) + 6 - [s(n) - s(n - 5)],$$

$$g(n - 12) + \binom{n - 12}{2} + 14(n - 12) + 39 - [s(n) - s(n - 12)]\}.$$

Asymptotically, we have

$$g(n) = \max\left\{g(n - 4) - \frac{3n^2}{189} + O(n), g(n - 5) - \frac{n^2}{18} + O(n), g(n - 12) - \frac{5n^2}{3} + O(n)\right\}.$$

Consequently,  $g(n)$  is both decreasing and negative for large  $n$ . Direct computations yield  $g(32) = 0$ ,  $g(n) < 0$  for  $n > 32$  and  $g(n) - g(m) < 0$  when  $n > 32$  and  $n \geq m + 4$ . Let  $h(n) = \max\{g(n), 0\}$ . Then

- (a)  $h(n) = 0$  for  $n \geq 32$ , and
- (b)  $h(n) - h(m) \geq g(n) - g(m)$  for  $n \geq m + 4$  with strict inequality for  $n > 32$ .

This implies that

$$h(n) \geq \max\{h(n-4) + t_7(n-4) + 2(n-4) + 3 - [s(n) - s(n-4)], \tag{4}$$

$$h(n-5) + \binom{n-5}{2} + 4(n-5) + 6 - [s(n) - s(n-5)], \tag{5}$$

$$h(n-12) + \binom{n-12}{2} + 14(n-12) + 39 - [s(n) - s(n-12)] \tag{6}$$

and the inequality is strict for  $n > 32$ .

For  $n \leq 4$  any 3-graph has at most  $s(n) + h(n)$  edges, so the basis for the induction holds trivially, and we can assume that  $n \geq 5$ . If  $H$  is cancellative, then by Theorem 1.2,  $e(H) \leq s(n) \leq s(n) + h(n)$ , with equality only for  $S(n)$ . We may therefore assume that  $H$  is not cancellative.

First consider the case that  $H$  contains a copy of  $F_{1,4}$ , denoted  $v \cup U$  as before. For  $0 \leq i \leq 3$  let  $f_i$  be the number of edges in  $H$  with  $i$  points in  $v \cup U$ . By the induction hypothesis,  $f_0 \leq s(n-5) + h(n-5)$ . Note that by definition of  $F_{1,4}$  we have  $f_3 = 6$ . Also, there cannot be an edge of the form  $xu'u'$  with  $u, u' \in U$ , or for any other  $u'' \in U$  we have a copy of  $F_5$  with edges  $xu'u', vu'u'', vu'u''$ . So any edge with 2 points in  $v \cup U$  must contain  $v$ , giving  $f_2 \leq 4(n-5)$ . Now every pair of vertices in  $F_{1,4}$  belongs to an edge, so Lemma 2.1 (ii) implies that the links of its vertices are edge-disjoint, i.e.  $f_1 \leq \binom{n-5}{2}$ . Therefore by (5) we have

$$e(H) = \sum_0^3 f_i \leq s(n-5) + h(n-5) + \binom{n-5}{2} + 4(n-5) + 6 \leq s(n) + h(n),$$

with strict inequality for  $n > 32$ , and we are done in this case.

Therefore we can assume that  $H$  contains no copy of  $F_{1,4}$ . Since  $H$  is not cancellative, it contains a copy of  $F_4$ . Let  $S = \{s_1, s_2, s_3, s_4\}$  be its vertex set and  $\{s_1s_2s_3, s_1s_2s_4, s_1s_3s_4\}$  its edges. Note that every pair of vertices in  $S$  belong to an edge, so by Lemma 2.1 (ii)  $G = \cup L(s_i)$  is a simple 4-colored graph on  $V - S$ .

*Case 1:* Suppose that  $G$  contains no copy of  $K_8$ .

Let  $f_i$  denote the number of edges of  $H$  with  $i$  points in  $S$ . By induction, we have  $f_0 \leq s(n-4) + h(n-4)$ . Also  $f_1 \leq t_7(n-4)$  by Turán’s theorem, and  $f_3 = 3$  by definition. To bound  $f_2$ , note that for any  $t \in V - S$  at most 2 pairs  $s_i s_j$  can belong to  $L(t)$ . For suppose there are 3 such pairs. They cannot all contain  $s_1$ , as then  $S \cup t$  would be a copy of  $F_{1,4}$ , which gives a case we have already excluded. Any other choice of 3 pairs must contain 2 of the form  $s_i s_j, s_i s_k$  with  $i \neq 1$ . Writing  $\{1, 2, 3, 4\} = \{i, j, k, \ell\}$  we see that  $s_j s_k s_\ell$  is an edge, and so  $ts_i s_j, ts_i s_k, s_j s_k s_\ell$  is a copy of  $F_5$ . This contradiction shows that  $L(t)$  contains at most 2 pairs from  $S$ , and so  $f_2 \leq 2(n-4)$ . Consequently, (4) yields

$$e(H) = \sum_0^3 f_i \leq s(n-4) + h(n-4) + t_7(n-4) + 2(n-4) + 3 \leq s(n) + h(n),$$

with strict inequality for  $n > 32$ , and we are done in this case.

Case 2: Suppose that  $G$  contains a copy of  $K_8$ .

Let  $T$  denote the vertex set of this copy of  $K_8$ , and let  $f_i$  denote the number of edges of  $H$  with exactly  $i$  elements in  $S \cup T$ . By induction, we have  $f_0 \leq s(n - 12) + h(n - 12)$ . Consider the 4-edge coloring on  $T$  given by the links  $L(s_i)$ . There can be no monochromatic  $K_4$  (as this gives a copy of  $F_{1,4}$ ) and no 2-colored triangle (this gives a copy of  $F_5$ ) so the coloring satisfies the hypothesis of Lemma 4.1. It follows that every vertex of  $T$  is incident to an edge of each of the four colors, so  $S \cup T$  induces a subhypergraph in which every pair of vertices belongs to an edge. Then Lemma 2.1 (ii) implies that the links  $L(v)$  are edge-disjoint for  $v \in S \cup T$ , and so  $f_1 \leq \binom{n-12}{2}$ .

To estimate  $f_2$ , we fix  $z \notin S \cup T$  and count pairs  $xy$  in  $S \cup T$  such that  $xyz$  is an edge. By the same argument as in Case 1, there are at most 2 such pairs with both  $x$  and  $y$  in  $S$ . For each  $y$  in  $T$  there is at most one  $x$  in  $S$  such that  $xyz$  is an edge, for if  $x'$  is another then  $yz$  is in  $L(x)$  and  $L(x')$ , contrary to Lemma 2.1 (ii). Thus there are at most 8 pairs with  $x$  in  $S$  and  $y$  in  $T$ . Next note that  $L(z)_T$  forms a matching. Otherwise there are edges  $xyz, x'yz$  and  $s_i xx'$  (where  $xx'$  has color  $i$ ) and this is a copy of  $F_5$ . This gives at most 4 pairs with  $x, y$  in  $T$ . In total there are at most 14 pairs  $xy$  for each  $z$ , so  $f_2 \leq 14(n - 12)$ .

Now we bound  $f_3$ , the number of edges  $xyz$  contained in  $S \cup T$ . There are 3 edges with  $x, y, z$  all in  $S$ . There are no edges with  $x, y$  in  $S$  and  $z$  in  $T$ . Otherwise Lemma 4.1 shows that there is  $w \in T$  with  $wz \in L(x)$  and then  $w' \in T$  with  $ww' \in L(y)$ . This gives  $xyz, xwz, ww'y$  forming a copy of  $F_5$ . Consequently, there are no edges of this form. The edges with two elements of  $T$  and one of  $S$  we bound simply by  $\binom{8}{2} = 28$ . Finally we claim that if  $x, y, z$  are all in  $T$  then  $xyz$  is a monochromatic triangle. For suppose it is rainbow, and say  $xy$  has color 1. There is some  $w \in T - xy$  such that  $wz$  has color 1, and then  $xyz, xyw, wz$  forms  $F_5$ . Thus  $xyz$  is monochromatic, and by Lemma 4.1, there are at most 8 such edges. In total we have  $f_3 \leq 39$ .

Putting everything together and recalling Eq. (6) we have,

$$e(H) = \sum_0^3 f_i \leq s(n - 12) + h(n - 12) + \binom{n - 12}{2} + 14(n - 12) + 39 \leq s(n) + h(n),$$

with strict inequality for  $n > 32$ , and the theorem is proved.  $\square$

### 5. Stability theorems

In this section, we will show that a cancellative 3-graph with almost as many edges as  $S(n)$  looks approximately like  $S(n)$ . Moreover, we will also show that an  $F_5$ -free 3-graph with almost as many edges as  $S(n)$  looks approximately like  $S(n)$ . Such theorems can be thought of as *stability* theorems, after the classical result of Simonovits [14] for graphs. To the best of our knowledge, these are among the first such theorems for hypergraphs.

We will use Theorems 1.2 and 1.3 in their proofs, and we also need the following slight variation on a case of the Simonovits stability theorem. It differs from the usual formulation, in that the conclusion refers to deleting vertices rather than edges, but this version follows easily from the proof given in [2, pp. 340–342]. See also [9] for a version with explicit constants.

**Proposition 5.1.** *For any  $\varepsilon' > 0$  there exists  $\delta' > 0$  such that if  $G$  is a  $K_4$ -free graph on  $n$  vertices with at least  $(1 - \delta')t_3(n)$  edges, then one can delete  $\varepsilon'n$  vertices from  $G$  so that the remaining graph is tripartite.*

The following theorem is a stability version of Bollobás' theorem.

**Theorem 5.2.** *For any  $\varepsilon > 0$  there exists  $\delta > 0$  such that if  $H$  is a cancellative 3-graph on  $n$  vertices with at least  $(1 - \delta)s(n)$  edges, then there is a partition of the vertex set of  $H$  as  $V(H) = U_1 \cup U_2 \cup U_3$  so that all but at most  $\varepsilon n^3$  edges of  $H$  have one point in each  $U_i$ .*

**Proof.** Note that we may and will assume that  $\varepsilon$  is sufficiently small, as this only strengthens the conclusion we reach. We can also assume that  $n$  is sufficiently large, as if we prove the theorem for  $n > n_0$  then it automatically holds for all  $n$  with  $\delta$  replaced by  $\min\{\delta, n_0^{-3}\}$ . Our constants satisfy the hierarchy  $\delta \ll \delta' \ll \varepsilon' \ll \varepsilon$ . More precisely, given  $\varepsilon > 0$ , choose  $\varepsilon' < (1/2)10^{-7}\varepsilon^2$ . Fix  $\delta' < (1/2)10^{-7}\varepsilon^2$  so that Proposition 5.1 applies with  $\varepsilon'$ , and let  $\delta < 27 \times 10^{-10}(\varepsilon\delta')^2$ .

In the course of the proof we will form a set  $V_0$  of bad vertices, which will always have size at most  $\frac{1}{2}\varepsilon n$ . Our arguments will always apply to  $H$  restricted to  $V - V_0$ , and we will show that we can delete at most  $\frac{1}{2}\varepsilon n^3$  edges from  $V - V_0$  to make it tripartite. Since  $V_0$  is incident to less than  $\frac{1}{2}\varepsilon n^3$  edges we can delete them and then extend to a tripartition of  $V$  by distributing the vertices of  $V_0$  arbitrarily.

Suppose there are  $10^{-5}\varepsilon\delta'n$  vertices of degree at most  $(1 - 10^{-4}\varepsilon\delta')n^2/9$ . Deleting them we arrive at a hypergraph  $H'$  on  $(1 - 10^{-5}\varepsilon\delta')n$  vertices with at least  $(1 - \delta - 3 \times 10^{-5}\varepsilon\delta'(1 - 10^{-4}\varepsilon\delta'))n^3/27$  edges. The choice of  $\delta$  implies that

$$1 - \delta - 3 \times 10^{-5}\varepsilon\delta'(1 - 10^{-4}\varepsilon\delta') > 1 - 3 \times 10^{-5}\varepsilon\delta' + 3(10^{-5}\varepsilon\delta')^2 > (1 - 10^{-5}\varepsilon\delta')^3.$$

Consequently,  $e(H') > \frac{1}{27}[(1 - 10^{-5}\varepsilon\delta')n]^3$ , which contradicts Bollobás' theorem. We deduce that there are at most  $10^{-5}\varepsilon\delta'n$  vertices of degree at most  $(1 - 10^{-4}\varepsilon\delta')n^2/9$ , and we will put them in the bad set  $V_0$ .

Now fix some edge  $x_1x_2x_3$  in  $V - V_0$ . By Lemma 2.1 (ii), the 3 links  $L(x_i)$  are edge-disjoint graphs, and by the previous remark each has at least  $(1 - 10^{-4}\varepsilon\delta')n^2/9$  edges. We think of their union as a graph  $J$  with at least  $(1 - 10^{-4}\varepsilon\delta')n^2/3$  edges, where  $L(x_i)$  is colored with color  $i$ . As in our proof of Theorem 1.2, we see that  $J$  cannot contain a  $K_4$ . For together with  $x_1x_2x_3$  this would give a 7 vertex subhypergraph in which every pair of vertices is contained in an edge. This gives 7

pairwise disjoint link graphs each with at least  $(1 - 10^{-4}\epsilon\delta')n^2/9$  edges, so their total number of edges is more than  $\binom{n}{2}$ , which is impossible.

Suppose that  $J$  has  $10^{-1}\delta'n$  vertices of degree at most  $(1 - 10^{-3}\epsilon)2n/3$ . Deleting them we arrive at a graph  $J'$  on  $(1 - 10^{-1}\delta')n$  vertices with at least  $(1 - 10^{-4}\epsilon\delta' - 2 \times 10^{-1}\delta'(1 - 10^{-3}\epsilon))n^2/3$  edges. Since  $\delta' < 10^{-2}\epsilon$  we have  $e(J') > \frac{1}{3}[(1 - 10^{-1}\delta')n]^2$ . But  $J' \subset J$  is  $K_4$ -free, so this contradicts Turán’s theorem. We deduce that there are at most  $10^{-1}\delta'n$  vertices of degree at most  $(1 - 10^{-3}\epsilon)2n/3$  in  $J$ , and we will put them in the bad set  $V_0$ .

Since  $J$  has at least  $(1 - 10^{-4}\epsilon\delta' - 2 \times 10^{-1}\delta'(1 - 10^{-3}\epsilon))n^2/3$  remaining edges, and this is trivially at least  $(1 - \delta')n^2/3$ , we can apply Proposition 5.1. This shows that we can add at most  $\epsilon'n$  vertices to  $V_0$ , and then partition  $V - V_0$  into 3 sets  $V_1, V_2, V_3$  each containing no edges of  $J$ . Then  $J$  restricted to  $V_1 \cup V_2 \cup V_3$  has at least  $(1 - \delta' - \epsilon')n^2/3 > (1 - 10^{-7}\epsilon^2)n^2/3$  edges (by the choice of  $\delta', \epsilon'$ ).

Note that  $|V_i - n/3| < 10^{-3}\epsilon n$  for each  $i$ , or  $J$  would have at most

$$\begin{aligned} |V_i|(n - |V_i|) + (n - |V_i|)^2/4 &= \frac{1}{3}n^2 - \frac{1}{12}(3|V_i| - n)^2 \\ &< \frac{1}{3}n^2 - \frac{3}{4}10^{-6}\epsilon^2n^2 < (1 - 10^{-7}\epsilon^2)n^2/3 \end{aligned}$$

edges which is impossible. Since each vertex in  $V - V_0$  has degree at least  $(1 - 10^{-3}\epsilon)2n/3$ , we see that each vertex  $v_i$  in  $V_i$  has degree at least  $(1 - 10^{-3}\epsilon)2n/3 - (1/3 + 10^{-3}\epsilon)n > n/3 - 10^{-2}\epsilon n$  in both  $V_j, j \neq i$ .

Since  $J$  restricted to  $V_1 \cup V_2 \cup V_3$  has more than  $|V_1 \cup V_2 \cup V_3|^2/4$  edges (for small  $\epsilon$ ) it contains a triangle by Turán’s theorem. Each  $V_i$  is independent, so we can take the vertices of the triangle to be  $v_1v_2v_3$  with  $v_i$  in  $V_i$ . For each  $V_i$  add any vertex that is not adjacent to both  $v_j, j \neq i$  to  $V_0$ . There are at most  $6 \times 10^{-2}\epsilon n$  such vertices. By Lemma 2.1 (iii) all triangles of  $J$  are multicolored, so we can suppose  $v_iv_j$  has color  $k$ , where  $\{i, j, k\} = \{1, 2, 3\}$ . Then each vertex of  $V_k$  is joined to the vertices  $v_i, v_j$  by one edge of color  $i$  and one of color  $j$ . Let  $V_k^1$  consist of those vertices  $v$  in  $V_k$  for which  $vv_i$  has color  $i$  and  $vv_j$  has color  $j$  (so the color matches the subscript), and  $V_k^2 = V_k - V_k^1$ .

All edges from  $v_1$  to  $V_2^1 \cup V_3^1$  have color 1. Therefore there are no edges between  $V_2^1$  and  $V_3^1$ , and the same holds between  $V_i^1$  and  $V_j^1$  for any two distinct  $i, j \in \{1, 2, 3\}$ . If both  $V_i^1$  and  $V_j^1$  have size at least  $10^{-2}\epsilon n$ , then  $J$  has at most  $n^2/3 - (10^{-2}\epsilon n)^2 < (1 - 10^{-7}\epsilon^2)n^2/3$  edges, which is impossible. It follows that there is at most one  $l$  for which  $|V_l^1| \geq 10^{-2}\epsilon n$ . Without loss of generality we assume that  $l = 1$ . Thus both  $V_2^1$  and  $V_3^1$  have size at most  $10^{-2}\epsilon n$ , and we add their vertices to  $V_0$ .

Now take any edge  $pqr$  of  $H$  in  $V - V_0$ . Note that it cannot have 2 of its vertices in one of the sets  $V_1^1, V_1^2, V_2^2$  or  $V_3^2$ . For example, if  $p$  and  $q$  both belonged to  $V_1^1$  then  $x_2v_2$  belongs to both the links  $L(p)$  and  $L(q)$ , which contradicts Lemma 2.1 (ii). The other cases are similar. Next suppose that we have  $p \in V_1^1, q \in V_1^2$  and  $r \in V_2^2$ . This situation is illustrated in Fig. 1 above (the numbers on the edges are their colors).




Fig. 1.

Note that  $qv_3$  has color 2 and  $rv_3$  has color 1, so if  $qr$  is an edge of  $J$  it must have color 3. But  $pv_3$  has color 3, i.e.  $x_3pv_3$  is an edge, and  $qr$  belongs to  $L(p)$  and  $L(x_3)$ , which contradicts Lemma 2.1 (ii). We deduce that  $qr$  is not an edge of  $J$ . Since  $J$  has at least  $(1 - 10^{-7}\epsilon^2)n^2/3$  edges respecting the partition  $(V_1, V_2, V_3)$  out of at most  $n^2/3$  possible edges, there are at most  $10^{-7}\epsilon^2n^2/3$  choices for  $qr$ , so at most  $10^{-7}\epsilon^2n^3/3$  such edges  $pqr$ .

Similarly there are at most  $10^{-7}\epsilon^2n^3/3$  edges  $pqr$  with  $p \in V_1^1$ ,  $q \in V_1^2$  and  $r \in V_3^2$ . These edges are exceptional; all others have one point in each of  $V_1^1 \cup V_1^2$ ,  $V_2^2$  and  $V_3^2$ . Define a tripartition  $V = U_1 \cup U_2 \cup U_3$  so that  $V_1^1 \cup V_1^2 \subset U_1$ ,  $V_2^2 \subset U_2$ ,  $V_3^2 \subset U_3$  and the bad vertices  $V_0$  are distributed arbitrarily. Since  $|V_0| < \frac{1}{2}\epsilon n$  and there are less than  $\frac{1}{2}\epsilon n^3$  exceptional edges we see that all but at most  $\epsilon n^3$  edges of  $H$  have one point in each  $U_i$ , so the theorem is proved.  $\square$

**Remark.** It is possible that Theorem 5.2 can also be proved by extending the ideas of Bollobás’ original proof of Theorem 1.2. His method was to construct three large disjoint sets with the property that every edge had at most one point from each and then use a counting argument to show that the number of edges is maximized when there are no other points outside these sets. (See [3] for an exposition.) Our framework seems more general however, since the same approach gives a proof of Theorem 1.5 below. As far as we can tell, there is no straightforward modification of Bollobás’ original proof that yields this stronger result.

Now we use the preceding theorem to prove a stability version of the Frankl–Füredi theorem.

**Proof of Theorem 1.5.** Again we can assume  $\epsilon$  sufficiently small and  $n$  sufficiently large. By Theorem 5.2 we can choose  $\delta < \epsilon/100$  so that any cancellative 3-graph on  $n'$  vertices with at least  $(1 - \delta)s(n')$  edges has a tripartition with at most  $\frac{1}{2}\epsilon n'^3$  wrong edges.

Suppose that there are  $\frac{1}{2}\varepsilon n$  vertices of degree at most  $(1 - 10^{-2})n^2/9$ . Deleting them we arrive at a hypergraph  $H'$  on  $(1 - \varepsilon/2)n$  vertices with at least  $(1 - \delta - 3 \cdot \varepsilon/2 \cdot (1 - 10^{-2}))n^3/27 > \frac{1}{27}[(1 - \varepsilon/2)n]^3$  edges, which contradicts Theorem 1.4. Consequently, there are at most  $\frac{1}{2}\varepsilon n$  vertices of degree at most  $(1 - 10^{-2})n^2/9$ , and we denote them  $V_0$ .

Set  $n' = |V - V_0|$ . Then  $H$  restricted to  $V - V_0$  has at least  $(1 - \delta)s(n')$  edges. If  $H$  contains no copy of  $F_4$ , then it is cancellative, and by choice of  $\delta$  there is a partition  $V(H) = U_1 \cup U_2 \cup U_3$  so that all but at most  $\frac{1}{2}\varepsilon n^3$  edges of  $H$  have one point in each  $U_i$ . Since  $|V_0| < \frac{1}{2}\varepsilon n$ , we can distribute the vertices of  $V_0$  arbitrarily to obtain the required partition. Therefore we can assume that  $H$  contains a copy of  $F_4$ .

Now we argue as in our proof of Theorem 1.4. Let  $S = s_1s_2s_3s_4$  be the vertex set of this  $F_4$ . Then  $\cup L(s_i)$  is a simple 4-colored graph on  $V - V_0 - S$  with at least  $(1 - 10^{-2})4n^2/9$  edges, so it contains a copy of  $K_8$ . Again, by Lemma 4.1 this  $K_8$  either contains a monochromatic  $K_4$  or each of its vertices is incident to all 4 colors. Thus we find a set  $X$  with  $|X| = 5$  or  $|X| = 12$  that induces a subhypergraph in which every pair of vertices belongs to an edge. Then the links of its vertices form edge-disjoint graphs on  $V - V_0 - S$  each with at least  $(1 - 10^{-2})n^2/9$  edges. This gives a total of more than  $\binom{n}{2}$ , which is a contradiction, so the theorem is proved.  $\square$

## Acknowledgments

Research of the second author supported by grants DMS-9970325 and DMS-0400812.

## References

- [1] B. Bollobás, Three-graphs without two triples whose symmetric difference is contained in a third, *Discrete Math.* 8 (1974) 21–24.
- [2] B. Bollobás, *Extremal graph theory*, London Mathematical Society Monographs, Vol. 11, Academic Press, Inc., London, New York, 1978.
- [3] B. Bollobás, *Combinatorics, Set Systems, Hypergraphs, Families of Vectors and Combinatorial Probability*, Cambridge University Press, Cambridge, 1986.
- [4] D. de Caen, Z. Füredi, The maximum size of 3-uniform hypergraphs not containing a Fano plane, *J. Combin. Theory Ser. B* 78 (2) (2000) 274–276.
- [5] P. Frankl, Z. Füredi, A new generalization of the Erdős-Ko-Rado theorem, *Combinatorica* 3 (3–4) (1983) 341–349.
- [6] Z. Füredi, Turán type problems, in: *Surveys in Combinatorics*, London Mathematical Society Lecture Note Series 166, Cambridge University Press, Cambridge, 1991, pp. 253–300.
- [7] Z. Füredi, M. Simonovits, Triple systems not containing a Fano Configuration, *Combinatorics, Probab. Comput.*, to appear.
- [8] P. Keevash, B. Sudakov, The exact Turán number of the Fano plane, *Combinatorica*, to appear.
- [9] P. Keevash, B. Sudakov, On a hypergraph Turán problem of Frankl, *Combinatorica*, to appear.

- [10] D. Mubayi, V. Rödl, On the Turán number of triple systems, *J. Combin. Theory Ser. A* 100 (2002) 136–152.
- [11] J.B. Shearer, A new construction for cancellative families of sets, *Electron. J. Combin.* 3 (1) (1996) Research Paper 15, approx. 3pp (electronic).
- [12] A.F. Sidorenko, On the maximal number of edges in a homogeneous hypergraph that does not contain prohibited subgraphs, *Mat. Zametki* 41 (3) (1987) 433–455.
- [13] A. Sidorenko, What we know and what we do not know about Turán numbers, *Graphs and Combinatorics* 11 (1995) 179–199.
- [14] M. Simonovits, A method for solving extremal problems in graph theory, stability problems, 1968 *Theory of Graphs (Proc. Colloq., Tihany, 1966)*, Academic Press, New York, 1968, pp. 279–319.
- [15] P. Turán, Eine Extremalaufgabe aus der Graphentheorie, *Mat. Fiz. Lapok* 48 (1941) 436–452.