


TOPOLOGY & GROUPS
 MICHAELMAS 2008
 QUESTION SHEET 4

1. Triangulate the torus as shown below



Let x and y be the loops $(1, 2, 3, 1)$ and $(1, 4, 7, 1)$, and let K be the union of these two loops (ie. K comes from the boundary of the square).

- (i) Prove that any edge loop based at 1 is equivalent to an edge loop lying entirely in K .
 - (ii) Deduce that any edge loop based at 1 is equivalent to a word in the alphabet $\{x, y\}$.
 - (iii) Show that the edge loops xy and yx are equivalent.
 - (iv) Deduce that any edge loop based at 1 is equivalent to $x^m y^n$, for $m, n \in \mathbb{Z}$.
 - (v) Prove that if $x^m y^n \sim x^M y^N$, then $m = M$ and $n = N$. [Hint: define 'winding numbers' as in the proof of Theorem III.32.]
 - (vi) Deduce that the fundamental group of the torus is isomorphic to $\mathbb{Z} \times \mathbb{Z}$.
2. Prove that every non-trivial element of a free group has infinite order.
3. The centre $Z(G)$ of a group G is $\{g \in G : gh = hg \forall h \in G\}$. Let S be a set with more than one element. Prove that the centre of $F(S)$ is the identity element.

4. (i) Let F be the free group on the three generators x , y and z . For non-zero integers r , s and t , show that the subgroup of F generated by x^r , y^s and z^t is freely generated by these elements.
- (ii) Let H be the subgroup of $F(\{x, y\})$ generated by x^2 , y^2 , xy and yx . Show that H is not freely generated by these elements.
5. Compute an explicit free generating set for the fundamental group of the following graph:

