

# COMPUTING ZETA FUNCTIONS OF ARTIN-SCHREIER CURVES OVER FINITE FIELDS

ALAN G.B. LAUDER AND DAQING WAN

ABSTRACT. We present a practical polynomial-time algorithm for computing the zeta function of certain Artin-Schreier curves over finite fields. This yields a method for computing the order of the Jacobian of an elliptic curve in characteristic 2, and more generally, any hyperelliptic curve in characteristic 2 whose affine equation is of a particular form.

## 1. INTRODUCTION

We present a low-degree polynomial-time algorithm for computing the zeta function of certain Artin-Schreier curves defined over finite fields. One consequence is a practical method for computing the order of the Jacobian of an elliptic curve in characteristic 2, and more generally, any hyperelliptic curve whose affine equation is of a particular form. Hyperelliptic curves have been proposed for use in public key cryptosystems by Koblitz [2, 14]. Our algorithm provides the first method of finding “random” hyperelliptic curves of arbitrary genus defined over large finite fields of characteristic 2 whose Jacobians have orders suitable for cryptographic use. Our method can be extended to more general curves, and we plan to present one such generalisation in a sequel paper.

We now introduce some notation which will allow us to explain our results. Let  $p$  denote a prime number, and  $a$  a positive integer. Define  $q = p^a$  and denote by  $\mathbb{F}_q$  the finite field with  $q$  elements. Fix an algebraic closure  $\bar{\mathbb{F}}_q$  of  $\mathbb{F}_q$  and let  $\mathbb{F}_{q^k}$  be the unique subfield of order  $q^k$ . We write  $\bar{\mathbb{F}}_q^*$  for the set of non-zero elements in  $\bar{\mathbb{F}}_q$ . The Artin-Schreier curves over  $\mathbb{F}_q$  we consider in this paper are defined by an equation of the form

$$(1) \quad Z^p - Z = f(X)$$

---

*Date:* November 21, 2001.

Alan Lauder gratefully acknowledges the support of the EPSRC (Grant GR/N35366/01) and St John’s College, Oxford. Daqing Wan is partially support by the NSF. The research was partly undertaken when the authors were visiting the Lorentz Center, University of Leiden, and the second author the Institute for Mathematical Sciences, National University of Singapore. We wish to thank Hendrik Lenstra and Harald Niederreiter, respectively, for their support during these visits, and also Pierrick Gaudry and Frederik Vercauteren for helpful comments on an earlier version of the paper.

*Mathematics Subject Classification 2000:* 11T99, 11Y16, 14Q05 *Key words and phrases:* Artin-Schreier, hyperelliptic, finite field, zeta function, algorithm, cryptography.

where  $f \in \mathbb{F}_q[X, X^{-1}]$  is a Laurent polynomial. Specifically, denote by  $C_f$  the curve embedded in  $\mathbb{F}_q^* \times \mathbb{F}_q$  with equation (1) and let  $\tilde{C}_f$  be the unique smooth projective curve birational to  $C_f$ . Let  $d$  denote the largest absolute value of any exponent which occurs in a non-zero term of  $f$ . For example, if  $f \in \mathbb{F}_q[X]$  this is just the degree. Our main theorem is

**Theorem 1.** *The zeta function of the smooth projective curve  $\tilde{C}_f$  may be computed deterministically in  $\tilde{O}(p^4 a^3 d^{5+\delta})$  bit operations. Here  $\delta = 0$  for  $p > 2$  and  $\delta = 1$  for  $p = 2$ .*

Here we use Soft-Oh notation  $\tilde{O}$  which ignores logarithmic factors, as in [15, Section 6.3]. More details of the complexity using different methods of arithmetic, and also the space complexity, can be found in Section 8.2.

We now explain how our algorithm may be applied to certain hyperelliptic curves in characteristic 2. Let  $C$  denote the affine curve with equation

$$Y^2 + X^m Y = h(X)$$

where  $h(X) \in \mathbb{F}_{2^a}[X]$  is of degree  $2g+1$  and  $m$  is a non-negative integer not greater than  $g$ . Let  $\tilde{C}$  be the unique smooth projective curve birational to  $C$ . Then  $\tilde{C}$  is birational to an Artin-Schreier curve, as explained in Note 5, and thus one may compute the zeta function of  $\tilde{C}$  in the complexity bounds of Theorem 1. From this it follows

**Corollary 2.** *The order of the Jacobian of the curve  $\tilde{C}$  may be computed deterministically in  $\tilde{O}(a^3 g^6)$  bit operations.*

This algorithm for hyperelliptic curves in characteristic 2 has been implemented by Vercauteren [23]. With regard to the dependence on  $a$ , we note that our method when restricted to elliptic curves has comparable time complexity to [18]. Moreover, it is the first practical algorithm for hyperelliptic curves in characteristic 2 which has polynomial-time growth in both the field size and genus. (The problem of polynomial-time computability for arbitrary varieties in small characteristic was already solved in [15], but the general algorithm there is not very practical. Also, a practical algorithm for hyperelliptic curves in odd characteristic is presented in [12] using different, though related, methods.) We refer to the references in [2] for the large literature on point counting, including [7, 19], and the more recent work [8, 9, 10, 11, 12, 17, 18, 22, 24, 26].

Sections 2, 3, 4 and 5 lay the mathematical foundation of our algorithm: it is based mainly upon an extension of the work of Dwork [6] due to Adolphson and Sperber [1]. Section 6 contains a statement of the algorithm for what we call Type 1 Artin-Schreier curves, and Section 7 describes exactly how to perform the main steps. The complexity analysis is tied up in Section 8, and Section 9 discusses the remaining type of Artin-Schreier curve in a more condensed fashion. As in [15] we aim to give a largely self-contained presentation.

## 2. L-FUNCTIONS AND ARTIN-SCHREIER CURVES

**2.1. General theory.** Let  $\bar{\mathbb{Q}}$  denote an algebraic closure of the rationals  $\mathbb{Q}$ . Let  $\Psi : \mathbb{F}_p \rightarrow \bar{\mathbb{Q}}$  be a non-trivial additive character, and  $\text{Tr}_k : \mathbb{F}_{q^k} \rightarrow \mathbb{F}_p$  the absolute trace map. A specific  $\Psi$  will be constructed in Section 4.2 but for now it may be arbitrary. Define  $\Psi_k : \mathbb{F}_{q^k} \rightarrow \bar{\mathbb{Q}}$  to be the non-trivial additive character  $\Psi \circ \text{Tr}_k$ .

For  $f \in \mathbb{F}_q[X, X^{-1}]$  define

$$(2) \quad S_k^*(f, \Psi) := \sum_{x \in \mathbb{F}_{q^k}^*} \Psi_k(f(x))$$

$$(3) \quad L^*(f, \Psi, T) := \exp \left( \sum_{k=1}^{\infty} \frac{S_k^*(f, \Psi)}{k} T^k \right).$$

For simplicity we shall omit the  $\Psi$  in this notation. Let  $C_f$  be the curve embedded in  $\bar{\mathbb{F}}_q^* \times \bar{\mathbb{F}}_q$  with equation

$$Z^p - Z = f(X).$$

Let  $\tilde{C}_f$  denote the unique smooth projective curve birational to  $C_f$ .

**Lemma 3.** *For each  $x \in \mathbb{F}_{q^k}^*$  there are exactly  $n_x$  points of the form  $(x, z) \in \mathbb{F}_{q^k}^* \times \mathbb{F}_{q^k}$  on  $C_f$  where*

$$n_x = \begin{cases} p & \text{if } \text{Tr}_k(f(x)) = 0 \\ 0 & \text{if } \text{Tr}_k(f(x)) \neq 0. \end{cases}$$

*Proof.* Follows from [16, Theorem 2.25].  $\square$

Denote by  $C_f(\mathbb{F}_{q^k})$  the set of  $\mathbb{F}_{q^k}$ -rational points on  $C_f$ . From Lemma 3 one deduces

$$\#(C_f(\mathbb{F}_{q^k})) = \sum_{a=0}^{p-1} \sum_{x \in \mathbb{F}_{q^k}^*} \Psi_k(af(x)) = \sum_{\theta \in G} \theta(S_k^*(f)) + (q^k - 1),$$

where  $G$  is the Galois group of  $\mathbb{Q}(\zeta)$  over  $\mathbb{Q}$ , with  $\zeta := \Psi(1)$  a primitive  $p$ th root of unity. Writing  $Z(C_f, T)$  for the zeta function [15, Section 1] of  $C_f$  it follows that

$$(4) \quad Z(C_f, T) = \frac{\{\prod_{\theta \in G} \theta(L^*(f, T))\}(1 - T)}{(1 - qT)}.$$

Here  $G$  acts on power series  $\mathbb{Q}(\zeta)[[T]]$  coefficient-wise, fixing monomials  $T^k$ .

To proceed further it is necessary to split the possible Laurent polynomials  $f$  into three types.

**2.2. Three types of Artin-Schreier curves.** *Type 1:* Assume that  $f \in \mathbb{F}_q[X]$  has degree  $d$  not divisible by  $p$ . Here we can also define

$$(5) \quad S_k(f) := \sum_{x \in \mathbb{F}_{q^k}} \Psi_k(f(x)) = S_k^*(f) + \Psi(k \text{Tr}_1(f(0)))$$

$$(6) \quad L(f, T) := \exp \left( \sum_{k=1}^{\infty} \frac{S_k(f)}{k} T^k \right) = L^*(f, T)(1 - \Psi(\text{Tr}_1(f(0)))T)^{-1}.$$

Hence from (4)

$$Z(C_f, T) = \frac{\{\prod_{\theta \in G} \theta(L(f, T))\}(1 - T)}{(1 - qT)\{\prod_{\theta \in G} \theta(1 - \Psi(\text{Tr}_1(f(0)))T)\}}.$$

(The latter term on the denominator is  $(1 - T)^{p-1}$  or  $1 + T + \dots + T^{p-1}$  depending upon whether  $\text{Tr}_1(f(0))$  is zero or not, although this does not concern us.) By Weil, the L-function  $L(f, T)$  is a polynomial of degree  $d - 1$ , and all of its reciprocal roots have complex absolute value  $\sqrt{q}$  under all complex embeddings of  $\bar{\mathbb{Q}}$ . Thus  $\prod_{\theta \in G} \theta(L(f, T))$  is a polynomial of degree  $(p - 1)(d - 1)$ , pure of weight 1 (see [25, Section 3] for definitions).

Let  $Z(\tilde{C}_f, T)$  denote the zeta function of the smooth projective curve  $\tilde{C}_f$ . Let  $g$  be the genus of  $\tilde{C}_f$ . Again, by Weil, we know that

$$Z(\tilde{C}_f, T) = \frac{P(\tilde{C}_f, T)}{(1 - T)(1 - qT)},$$

where the numerator is a polynomial of degree  $2g$ , pure of weight 1. Since  $C_f$  and  $\tilde{C}_f$  are birational they differ by a finite number of points, and hence their zeta functions differ by a factor of weight 0. Comparing the pure weight 1 parts in  $Z(C_f, T)$  and  $Z(\tilde{C}_f, T)$ , we deduce

$$(7) \quad P(\tilde{C}_f, T) = \prod_{\theta \in G} \theta(L(f, T)).$$

In particular, the genus  $g$  of the curve  $\tilde{C}_f$  is given by the formula

$$(8) \quad g = (p - 1)(d - 1)/2.$$

*Type 2:* Assume that  $f \in \mathbb{F}_q[X^{-1}]$  has negative degree  $d^-$  not divisible by  $p$ . That is,  $d^-$  is the lowest exponent occurring in  $f$ . Then  $C_f$  is birational to  $C_{f^*}$  where  $f^* := f(X^{-1})$ , and we have reduced to type 1.

*Type 3:* Assume that  $f \in \mathbb{F}_q[X, X^{-1}]$  but not in  $\mathbb{F}_q[X] \cup \mathbb{F}_q[X^{-1}]$ . Let the negative degree be  $d^-$  and the positive degree be  $d^+$ . Assume  $p$  does not divide  $d^- d^+$ . In this case, by Weil,  $L^*(f, T)$  is a polynomial of degree  $d^+ - d^-$ , pure of weight 1. By comparing the pure weight 1 parts in  $Z(C_f, T)$  and  $Z(\tilde{C}_f, T)$  we find that

$$(9) \quad P(\tilde{C}_f, T) = \prod_{\theta \in G} \theta(L^*(f, T)),$$



is the numerator of  $Z(\tilde{C}_f, T)$ . In particular, the genus  $g$  of the curve  $\tilde{C}_f$  is given by the formula

$$g = (p-1)(d^+ - d^-)/2.$$

Thus in all cases the computation of the zeta function of the smooth projective curve  $\tilde{C}_f$  reduces to the evaluation of the L-function of certain one variable exponential sums.

**Note 4** The degree conditions on  $f$  are essential for cohomological arguments; however, given a Laurent polynomial  $f$  which does not satisfy them one may replace it by a new polynomial  $\tilde{f}$  which does such that  $C_f$  and  $C_{\tilde{f}}$  are isomorphic. For each term  $a_{jp}X^{jp}$ , the isomorphism  $Z \rightarrow Z + bX^j$  ( $X \rightarrow X$ ) shows that we can replace the term  $a_{jp}X^{jp}$  with  $b_jX^j$  where  $b := a_{jp}^{1/p} \in \mathbb{F}_q$ . Repeat this procedure until no term has a non-zero exponent divisible by  $p$ . The resulting polynomial  $\tilde{f}$  has no terms with non-zero exponents divisible by  $p$ , and thus it certainly satisfies any necessary degree restrictions.

**Note 5** Let  $C$  be the curve in Corollary 2. Then  $C$  is birational to the curve  $C_f$  with equation  $Z^2 + Z = f(X)$  where  $f := X^{-2m}h$ . This can be seen by making the change of variable  $Y = ZX^m$ . Thus  $\tilde{C}$ , as in Corollary 2, is birational to  $\tilde{C}_f$  and the zeta function of  $\tilde{C}_f$  can then be computed using Note 4, to get  $f$  in the correct form, and the algorithm we will present. By the special value formula for the zeta function at  $T = 1$ , the order of the Jacobian is the numerator of the zeta function evaluated at  $T = 1$  [2, page 175]. Similar comments apply to curves over  $\mathbb{F}_q$  of the form  $Y^p - X^mY = h(X)$  where  $q = p^a$  and  $p-1$  divides  $m$ . Also, general hyperelliptic curves in characteristic 2 are birational to Artin-Schreier curves of the form  $Z^2 + Z = f(X)$  for  $f$  a rational function. As such they may be tackled using a generalisation of our approach.

**2.3. Type 1 Artin-Schreier curves.** In Sections 3, 4, 5, 6, 7 and 8 we shall denote by  $f$  a polynomial in  $\mathbb{F}_q[X]$  of degree  $d$  not divisible by  $p$ . We write  $f = \sum_{j \in J} a_j X^j$  where  $a_j \neq 0$ . In these sections we will explain how to compute the zeta function of the smooth projective curve  $\tilde{C}_f$ . That is, we shall cover classical “type 1” Artin-Schreier curves. In Section 9 we shall discuss the modifications required when  $f \in \mathbb{F}_q[X, X^{-1}]$  is a Laurent polynomial with both positive and negative exponents. These are “type 3” Artin-Schreier curves. As mentioned before, “type 2” may be reduced to “type 1” and we shall not discuss them again.

### 3. $p$ -ADIC THEORY

**3.1.  $p$ -adic rings.** Let  $\mathbb{Q}_p$  denote the  $p$ -adic numbers with ring of integers  $\mathbb{Z}_p$ . Fix  $\Omega$  the completion of an algebraic closure of  $\mathbb{Q}_p$ . Denote by  $\epsilon$  a

primitive  $(q-1)$ th root of unity in  $\Omega$ , and  $\pi \in \Omega$  an element which satisfies  $\pi^{p-1} = -p$ . Define

$$(10) \quad A := \mathbb{Z}_p[\epsilon, \pi].$$

In particular,  $A$  has residue field  $\mathbb{F}_q$ . Elements in  $A$  may be represented via  $\pi$ -adic expansions whose coefficients are from some distinct set of representatives for the quotient  $A/(\pi)$  of size  $q$ . For any primitive  $p$ th root of unity  $\zeta$  we have that both  $\mathbb{Q}_p(\zeta)$  and  $\mathbb{Q}_p(\pi)$  are the unique totally ramified extensions of  $\mathbb{Q}$  of degree  $p-1$  (since  $\text{ord}(1-\zeta) = \text{ord}(\pi) = 1/(p-1)$ ). Thus  $\mathbb{Z}_p[\zeta] = \mathbb{Z}_p[\pi]$  and so  $A$  contains the exponential sums defined in the previous section. Let  $G$  denote the group of automorphisms  $\theta_j : \mathbb{Q}_p(\pi) \rightarrow \mathbb{Q}_p(\pi)$  for  $1 \leq j \leq p-1$  where each  $\theta_j$  fixes  $\mathbb{Q}_p$  and

$$(11) \quad \theta_j(\pi) := \eta^j \pi.$$

Here  $\eta \in \mathbb{Z}_p$  is a primitive  $(p-1)$ th root of unity. Then  $G$  is the Galois group of the extension  $\mathbb{Q}_p(\pi)/\mathbb{Q}_p$  (its action on  $\zeta$  is  $\theta_j : \zeta \mapsto \zeta^{\eta^j \bmod p}$ , although we shall not need this explicitly).

It will be convenient to work in complete rings which contain arbitrary roots  $\pi^r$  of  $\pi$ , for  $r$  a rational number. To this end, let  $\Pi = \bigcup_{z \in \mathbb{N}} \{\pi^{1/z}\}$  and  $\tilde{A}$  denote the completion of the ring  $\mathbb{Z}_p[\epsilon, \Pi]$ . Here  $\mathbb{N}$  denotes the positive integers.

Denote by  $\tau$  the map on  $\tilde{A}$  defined as

$$(12) \quad \tau(\epsilon) := \epsilon^p, \tau \text{ fixes } \mathbb{Z}_p[\Pi] \text{ and is continuous.}$$

Let  $\text{ord}$  and  $|\cdot|_p$  denote the  $p$ -adic valuation and norm on  $\tilde{A}$  normalised so that  $\text{ord}(p) = 1$  and  $|p|_p = 1/p$ .

Note that the ring  $A$  can be constructed and computed in easily; see [15, Section 3] for more details on  $p$ -adic fields. (The larger ring  $\tilde{A}$  is only introduced for mathematical convenience; all of our computations are performed in  $A$ .)

**3.2. A weight function.** As in [15, Section 4] we define a weight function: For each non-negative integer  $u \in \mathbb{Z}_{\geq 0}$  let

$$\text{wt}(u) := \lceil u/d \rceil.$$

Here  $\lceil * \rceil$  is the least integer not less than  $*$  and  $d$  the degree of the polynomial  $f \in \mathbb{F}_q[X]$ . Define  $\tilde{\text{wt}}(u) := u/d$  and so  $\lceil \tilde{\text{wt}}(u) \rceil = \text{wt}(u)$ . Notice that  $\pi^{\text{wt}(r)} \in A$  and  $\pi^{\tilde{\text{wt}}(r)} \in \tilde{A}$  for every  $r \in \mathbb{Z}_{\geq 0}$ .

**3.3. Banach modules.** Let  $B$  denote a complete subring of  $\Omega$ . A Banach module over  $B$  is an ultrametrically normed complete module  $E$  over  $B$ , such that  $\|re\| = |r|_p \|e\|$  for  $r \in B$  and  $e \in E$ , where  $\|\cdot\|$  is the module norm. An orthonormal basis for  $E$  is a set  $\{e_i \mid i \in \mathbb{N}\}$  such that every element in  $E$  can be written uniquely in the form  $\sum_i b_i e_i$  where  $b_i \in B$  with  $|b_i|_p \rightarrow 0$  as  $i \rightarrow \infty$ . (See [20] and [4, Section A].) In the case that  $B$  is a field we call it a Banach space.

**Definition 6.** For each rational number  $\delta > 0$ , let  $\tilde{L}(\delta)$  be the Banach module over  $\tilde{A}$  whose orthonormal basis consists of all terms  $\pi^{\tilde{\text{wt}}(r)\delta} X^r$  for  $r \in \mathbb{Z}_{\geq 0}$ . Let  $A\{X\}$  ( $\tilde{A}\{X\}$  respectively) denote the Banach module over  $A$  ( $\tilde{A}$  resp.) whose orthonormal basis consists of all terms  $X^r$  for  $r \in \mathbb{Z}_{\geq 0}$ .

Note that  $\tilde{L}(\delta') \subset \tilde{L}(\delta)$  for  $\delta' > \delta$  and all the above spaces lie in  $\tilde{A}\{X\}$ . One may check that all the above spaces are closed under multiplication, and are in fact rings. Extend  $\tau$  to act on each power series in the ring  $\tilde{A}\{X\}$  by taking

$$(13) \quad \tau(X) := X, \tau \text{ is } \tilde{A}\text{-linear and continuous.}$$

#### 4. ANALYTIC REPRESENTATION OF ADDITIVE CHARACTERS

**4.1. Dwork's splitting functions.** Let  $\theta(t)$  denote the splitting function [15, Section 4.1]

$$(14) \quad \theta(t) := \theta_1(t) = \exp(\pi(t - t^p)).$$

Write

$$(15) \quad \hat{f} := \sum_{j \in J} \hat{a}_j X^j$$

for the polynomial over  $A$  obtained by taking the Teichmüller lifting of each coefficient of  $f$ .

**Lemma 7.** For each term  $\hat{a}_j X^j$  in  $\hat{f}$ , we have  $\theta(\hat{a}_j X^j) \in \tilde{L}(\delta)$  for any

$$\delta < \left( \frac{p-1}{p} \right)^2$$

*Proof.* Writing  $\theta(t) =: \sum_{r=0}^{\infty} \lambda_r t^r$  we see from [15, Equation (3)] that  $\text{ord}(\lambda_r) \geq (p-1)r/p^2$ . Terms in  $\theta(\hat{a}_j X^j)$  are of the form  $\lambda_r \hat{a}_j^r X^{jr}$  for  $r$  a non-negative integer. Now  $\text{ord}(\lambda_r \hat{a}_j^r) \geq (p-1)r/p^2$  since  $\text{ord}(\hat{a}_j) = 0$ . Also  $0 \leq j \leq d$  and so  $\tilde{\text{wt}}(jr) = jr/d \leq r$ . Thus

$$\text{ord}(\lambda_r \hat{a}_j^r) \geq \frac{p-1}{p^2} r \geq \frac{p-1}{p^2} \tilde{\text{wt}}(jr).$$

Also for  $\delta < ((p-1)/p)^2$  we have

$$\text{ord}(\pi^{\delta \tilde{\text{wt}}(jr)}) < \frac{p-1}{p^2} \tilde{\text{wt}}(jr)$$

and the result follows.  $\square$

**Definition 8.** Let  $F$  and  $F^{(a)}$  be defined as

$$(16) \quad F := \prod_{j \in J} \theta(\hat{a}_j X^j)$$

$$(17) \quad F^{(a)} := \prod_{i=0}^{a-1} \tau^i(F(X^{p^i})).$$

Recall here that  $q = p^a$ . Both  $F$  and  $F^{(a)}$  are one-way infinite power series in  $A\{X\}$ . By Lemma 7 and the fact that each  $\tilde{L}(\delta)$  is a ring we have

**Lemma 9.** *The power series  $F \in \tilde{L}(\delta)$  for any*

$$\delta < \left(\frac{p-1}{p}\right)^2$$

**4.2. Dwork's additive character.** For  $k \geq 1$  define

$$\Phi_k(t) := \prod_{i=0}^{ak-1} \theta(t^{p^i}) \in \mathbb{Z}_p[\pi][[t]]$$

and let

$$\Phi(t) := \prod_{i=0}^{a-1} \theta(t^{p^i}) \in \mathbb{Z}_p[\pi][[t]].$$

Denote by Teich the Teichmüller lifting map from  $\bar{\mathbb{F}}_q$  to  $\mathbb{Z}_p^{\text{unram}}$ , where  $\mathbb{Z}_p^{\text{unram}}$  is the unramified integral closure of  $\mathbb{Z}_p$  in  $\Omega$ . Then

$$\Psi_k := \Phi_k \circ \text{Teich}, \Psi := \Phi \circ \text{Teich}$$

are non-trivial characters from  $\mathbb{F}_{q^k}$  and  $\mathbb{F}_p$ , respectively, to  $\mathbb{Z}_p[\pi]$ . We have that  $\Psi_k(x) = \Psi(\text{Tr}_k(x))$  where  $\text{Tr}_k$  is the trace map from  $\mathbb{F}_{q^k}$  to  $\mathbb{F}_p$ . (See [15, Lemma 6].)

The following lemma is proved exactly as [15, Proposition 9]. It gives an analytic expression for the exponential sum  $S_k^*(f) := S_k^*(f, \Psi)$

**Lemma 10.** *Let  $S_k^*(f)$  be the exponential sum defined in (2) using Dwork's additive character  $\Psi$ , and  $F^{(a)}(X)$  the power series from Definition 8. Then*

$$S_k^*(f) = \sum_{x^{q^k-1}=1} F^{(a)}(x) F^{(a)}(x^q) \dots F^{(a)}(x^{q^{k-1}}).$$

Here the sum is over the Teichmüller liftings in  $\mathbb{Z}_p^{\text{unram}}$  of the points on the torus  $\mathbb{F}_{q^k}^*$ .

**4.3. Completely continuous maps.** The next step is to introduce operators on the module  $L$  so that the righthand side of the above expression can be interpreted as the “trace” of a map on a Banach module.

**Definition 11.** *Let  $\psi_p$  be the map on  $\tilde{A}\{X\}$  which acts on monomials as*

$$\psi_p(X^r) := \begin{cases} X^{r/p} & \text{if } p \text{ divides } r \\ 0 & \text{otherwise} \end{cases}$$

and extends to all of  $\tilde{A}\{X\}$  by  $\tau^{-1}$ -linearity and continuity. Specifically,  $\psi_p(\sum_r A_r X^r) = \sum_{r, p|r} \tau^{-1}(A_r) X^{r/p}$ . Write  $\psi_q := \psi_p^a$ , a linear map since  $\tau^{-a}$  is the identity on  $\tilde{A}$ . Let  $\alpha := \psi_p \circ F$  and  $\alpha_a := \psi_q \circ F^{(a)}$ . Precisely,  $\alpha$  is multiplication by  $F$  followed by the map  $\psi_p$ , and likewise for  $\alpha_a$  (see [15, Definitions 20, 21]).

**Lemma 12.** *The maps satisfy  $\alpha_a = \alpha^a$ .*

*Proof.* Proved exactly as in [15, Lemma 22].  $\square$

**Lemma 13.** *The map  $\alpha$  is stable on  $\tilde{L}(\delta)$  for*

$$\delta < \frac{(p-1)^2}{p}$$

*Proof.* We first claim that  $\psi_p(\tilde{L}(\delta)) \subseteq \tilde{L}(p\delta)$  for any rational  $\delta > 0$ : For this it is enough to observe that  $\psi_p(\pi^{\tilde{\text{wt}}(r)\delta} X^r) = \pi^{\tilde{\text{wt}}(r/p)p\delta} X^{r/p}$  for any  $r$  divisible by  $p$ . Now let  $G \in \tilde{L}(\delta)$  where  $\delta$  satisfies the necessary inequality, depending on  $p$ . Then by Lemma 9  $F \in \tilde{L}(\delta/p)$  and so since also  $G \in \tilde{L}(\delta/p)$  by closure under multiplication we find that  $FG \in \tilde{L}(\delta/p)$ . Hence  $\psi_p(FG) \in \tilde{L}(\delta)$ , that is  $\alpha(G) \in \tilde{L}(\delta)$  as required.  $\square$

**Definition 14.** *Let  $L$  be defined as*

$$L := \begin{cases} \tilde{L}(1) \cap A\{X\} & \text{if } p > 2 \\ (\tilde{L}(\gamma) \cap A\{X\}) \otimes \mathbb{Q} & \text{if } p = 2. \end{cases}$$

Here  $\gamma$  may be taken to be any rational number in the range  $1/(2 + (1/2d)) < \gamma < 1/2$ . The key point is that  $\alpha$  is stable on  $\tilde{L}(\gamma)$ , since  $\gamma < 1/2$ , and the space  $\tilde{L}(\gamma)$  is small enough such that an ad hoc argument we present later (Lemma 28) works. Thus for  $p > 2$  we have that  $L$  is just the Banach module over  $A$  with orthonormal basis the terms  $\pi^{\text{wt}(u)} X^u$  for non-negative integers  $u$ . For  $p = 2$  it is the Banach space over  $A \otimes \mathbb{Q}$  with orthonormal basis the terms  $\pi^{\lceil \gamma \tilde{\text{wt}}(u) \rceil} X^u$  for non-negative  $u$ .

**Lemma 15.** *The maps  $\alpha$  and  $\alpha_a$  are stable on  $L$ .*

*Proof.* First suppose  $p > 2$ . Then  $\alpha$  is stable on  $\tilde{L}(1)$  by Lemma 13. Certainly  $\alpha$  is stable on the ring of convergent power series  $A\{X\}$ , since  $F \in A\{X\}$ . Thus  $\alpha$  is stable on  $\tilde{L}(1) \cap A\{X\} = L$ . That  $\alpha_a$  is stable on  $L$  now follows from Lemma 12. Second, consider the case  $p = 2$ . Putting  $p = 2$  in Lemma 13 we find  $\alpha$  is stable on  $\tilde{L}(\gamma) \cap A\{X\}$  since  $\gamma < 1/2$ . Given  $G \in L$  we have that  $mG \in \tilde{L}(\gamma) \cap A\{X\}$  for some  $p$ -adic integer  $m$ . The result now follows easily.  $\square$

**Note 16** The rings  $\tilde{A}$  and  $\tilde{L}(\delta)$  and the function  $\tilde{\text{wt}}$  were introduced to prove the above result in as simple a manner as possible; we shall have little further need for them, working from now on mainly with  $A$  and  $L$ .

For certain classes of linear maps on Banach modules the trace and determinant are defined. This is done in the usual way, via matrices for the maps with respect to an orthonormal basis. We refer to [20] and [4, Section A2] for definitions. The key result is the chain level Dwork Trace Formula.

**Theorem 17.** *With  $S_k^*(f)$  and  $L^*(f, T)$  the exponential sum and  $L$ -function defined as in (2) and (3) using Dwork's additive character  $\Psi$  (Section 4.2), and  $\alpha_a$  the map on  $L$  given in Definition 11 we have*

$$S_k^*(f) = (q^k - 1) \text{Tr}(\alpha_a^k | L).$$

*Thus we have*

$$L^*(f, T) = \frac{\det(1 - T\alpha_a | L)}{\det(1 - Tq\alpha_a | L)}.$$

*Here the trace and determinant are defined via matrices for the maps with respect to the orthonormal basis of  $L$ .*

*Proof.* This is in essence a case of [15, Theorem 25] (with “ $n = 0$ ”). Note that the matrix “ $M_a$ ” in [15, Theorem 25] is that for the map  $\alpha_a$  with respect to a “formal basis” [15, Section 5.2] of the form  $\{X^i \mid i \in \mathbb{Z}_{\geq 0}\}$ . For the above formulae we require a matrix with respect to the orthonormal basis  $\{\pi^{\text{wt}(i)} X^i \mid i \in \mathbb{Z}_{\geq 0}\}$  (when  $p > 2$ , with a slightly different basis for  $p = 2$ ). One may verify, using for example [15, Equation (12)], that the trace of powers of these two matrices are the same.  $\square$

These formulae may be used to compute the zeta function in a similar fashion to [15]. Namely, a finite matrix may be computed which represents the map  $\alpha$  acting on some appropriate modular reduction of  $L$ . This matrix is then used to compute the characteristic polynomial of  $\alpha_a$  itself up to a necessary  $p$ -adic accuracy. (This algorithm has been implemented by Vercauteren [23].) However, this “chain level” method results, for example, in a time complexity of  $\tilde{O}(a^{4.38})$  with space  $\tilde{O}(a^4)$  bits using the fastest methods for matrix multiplication and ring arithmetic. Using some homological algebra one can derive a better “cohomological” formula leading to an improved algorithm. This is what we do in this paper.

At this stage since  $f$  is an ordinary polynomial rather than a Laurent polynomial with negative and positive terms, we can do a little more work to derive a better chain level formula: Let  $L_{>0}$  denote the Banach module comprising of those power series in  $L$  with zero constant term. For  $p > 2$  the module  $L_{>0}$  is defined over  $A$ , and for  $p = 2$  over  $A \otimes \mathbb{Q}$ . It follows easily from Lemma 15 that

**Lemma 18.** *The maps  $\alpha$  and  $\alpha_a$  are stable on  $L_{>0}$ .*

Now one may check via a matrix for the map  $\alpha_a$  with respect to the orthonormal basis  $\{\pi^{\text{wt}(r)} X^r\}_{r \geq 0}$  for  $p > 2$ , and  $\{\pi^{\lceil \gamma \text{wt}(r) \rceil} X^r\}_{r \geq 0}$  for  $p = 2$ , that

$$\det(1 - T\alpha_a | L) = (1 - F^{(a)}(0)T) \det(1 - T\alpha_a | L_{>0}).$$

Here

$$F^{(a)}(0) = \Psi(\text{Tr}_1(a_0))$$

is a root of unity, where  $\Psi$  is Dwork's additive character from  $\mathbb{F}_p$  to  $\mathbb{Z}_p[\pi]$ . Let  $S_k(f)$  and  $L(f, T)$  be as in (5) and (6). Then one has  $S_k(f) = S_k^*(f) + \Psi(k\text{Tr}_1(a_0))$  and so  $L(f, T) = L^*(f, T)(1 - F^{(a)}(0)T)^{-1}$ . Hence we have

**Theorem 19.** *Let  $L(f, T)$  be the  $L$ -function for the exponential sum over the affine line from (6). Then*

$$L(f, T) = \frac{\det(1 - T\alpha_a|_{L_{>0}})}{\det(1 - qT\alpha_a|_L)}.$$

## 5. DWORK COHOMOLOGY

Let  $H$  be the polynomial in  $L$  defined as

$$(18) \quad H = \pi \hat{f}$$

(In fact  $H \in \tilde{L}(1) \cap A\{X\}$  in all cases, and this latter ring equals  $L$  for  $p > 2$ , and lies strictly within  $L$  for  $p = 2$ .)

Let  $D$  be the operator on  $L$  defined as

$$D = X \frac{d}{dX} + H_X$$

where

$$H_X := X \frac{dH}{dX}.$$

Here  $d/dX$  is the usual differential operator on polynomials extended to power series by continuity, with  $X$  and  $H_X$  just acting by multiplication. Now  $X(d/dX)$  is stable on  $L$ , and  $L$  is a ring. From this it follows that  $D$  is stable on  $L$ , and in fact maps  $L$  to  $L_{>0}$ .

**Note 20** We pause to explain the motivation behind the above definitions: Define  $\hat{\theta}(t) := \prod_{i=0}^{\infty} \theta(t^{p^i})$ . One may check that  $\hat{\theta}(t) = \exp(\pi t)$ . Now  $F = \prod_{j \in J} \theta(\hat{a}_j X^j)$ . Defining  $\hat{F}(X) := \prod_{j \in J} \hat{\theta}(\hat{a}_j X^j)$  we find that this equals  $\exp(\pi \hat{f})$  which is just  $\exp(H)$ , with  $H$  as in (18). Since  $\hat{a}_j^q = \hat{a}_j$  and  $\tau^i(\hat{a}_j) = \hat{a}_j^{p^i}$  for each  $j \in J$  it follows from Definition 8 that  $F^{(a)}(X) = \hat{F}(X)/\hat{F}(X^q)$ . From this we see  $\alpha_a = \psi_q \circ F^{(a)} = \exp(-H) \circ \psi_q \circ \exp(H)$ . Define the operator  $E := X(d/dX)$ , and so  $E \circ (q\psi_q) = \psi_q \circ E$ . Then with  $D$  as above one may check  $D = E + E(H) = \exp(-H) \circ E \circ \exp(H)$ . Thus  $D$  and  $\alpha_a$  are obtained from  $E$  and  $\psi_q$  by some kind of twisting; also, it now follows that  $D \circ (q\alpha_a) = \alpha_a \circ D$ , which is the crucial relation. (See [6, pages 55-60] and [21, pages 267-270] for more details.)

Let  $\mathcal{L}$  be the complex

$$0 \longrightarrow L \xrightarrow{D} L_{>0} \longrightarrow 0.$$

This is a complex of  $A$ -modules when  $p > 2$  and  $A \otimes \mathbb{Q}$ -spaces for  $p = 2$ . Denote by  $H_1$  and  $H_0$  the kernel and co-kernel of the map  $D$ . In particular,  $H_0 := L_{>0}/D(L)$ .

**Proposition 21.** *The map  $D$  is injective and so  $H_1 = 0$ . Moreover, for  $p > 2$  ( $p = 2$  respectively)  $H_0$  is a finite free  $A$ -module ( $A \otimes \mathbb{Q}$ -space respectively) of rank  $d - 1$ . A basis for  $H_0$  may be taken as the set of terms*

$$\{\pi^{\text{wt}(i)} X^i \mid 0 < i < d\} = \{\pi X, \pi X^2, \dots, \pi X^{d-1}\}.$$

*Proof.* Over the formal power series ring  $\Omega[[X]]$ , the formal solutions of the first order linear differential equation  $D = 0$  is the one-dimensional subspace generated by  $\exp(-H)$ . But the power series  $\exp(-H) \notin L$  (the decay rate of coefficients is too slow). This shows that the restriction of the operator  $D$  to  $L$  is injective. The second part of the proposition follows from the normal form computations in section 7.  $\square$

By Note 20 we have

$$D \circ q\alpha_a = \alpha_a \circ D.$$

Thus the map  $\alpha_a$  defines a chain map on  $\mathcal{L}$ :

$$\begin{array}{ccccccc} 0 & \longrightarrow & L & \xrightarrow{D} & L_{>0} & \longrightarrow & 0 \\ & & \downarrow q\alpha_a & & \downarrow \alpha_a & & \\ 0 & \longrightarrow & L & \xrightarrow{D} & L_{>0} & \longrightarrow & 0. \end{array}$$

Denote by  $H_0(\alpha_a)$  and  $H_1(q\alpha)$  the maps induced on the homology  $H_0$  and  $H_1$  by this chain map, and  $\det(1 - H_0(\alpha_a)T)$ ,  $\det(1 - H_1(q\alpha_a)T)$  the corresponding determinants.

**Theorem 22.** *The  $L$ -function from Theorem 19 satisfies*

$$L(f, T) = \det(1 - H_0(\alpha_a)T).$$

*Proof.* We have

$$\frac{\det(1 - T\alpha_a|_{L_{>0}})}{\det(1 - Tq\alpha_a|_L)} = \frac{\det(1 - H_0(\alpha_a)T)}{\det(1 - H_1(q\alpha_a)T)}.$$

This identity is proved in the same way that [20, Proposition 9] is derived from [20, Lemma 2]. Now  $H_1 = 0$  and so the denominator on the righthand side is 1. The expression for the  $L$ -function now follows from Theorem 19.  $\square$

**Corollary 23.** *The zeta function  $Z(\tilde{C}_f, T)$  of the smooth projective curve  $\tilde{C}_f$  birational to the affine curve with equation  $Z^p - Z = f(X)$  satisfies*

$$Z(\tilde{C}_f, T) = \frac{\prod_{j=1}^{p-1} \theta_j(\det(1 - H_0(\alpha_a)T))}{(1 - T)(1 - qT)}.$$

Here  $\theta_j$  are the automorphisms (11) of  $\mathbb{Z}_p[\pi]$  extended to act on polynomials by fixing  $T$ . The numerator is a polynomial of degree  $(p - 1)(d - 1)$ .

*Proof.* Follows from Theorem 22 and equation (7).  $\square$



Thus the strategy of the algorithm is to compute the determinant of the map  $H_0(\alpha_a)$  on the zeroth homology  $H_0$ , up to a suitable modular precision. This may be done efficiently via the following lemma, which is an immediate consequence of Lemma 12.

**Lemma 24.** *Let  $H_0(\alpha)$  denote the map induced on  $H_0$  by  $\alpha$ . Then  $H_0(\alpha_a) = H_0(\alpha)^a$ .*

It will be enough to compute the coefficients of the characteristic polynomial of Frobenius modulo  $p^N$  for

$$N = \lfloor (p-1)(d-1)(1+a/2) + 1 \rfloor.$$

This follows since the L-function of the exponential sum  $L(f, T)$  has reciprocal roots whose complex absolute values are  $\sqrt{q}$ . Thus the coefficient of  $T^k$  in the polynomial  $\prod_{\theta_j \in G} \theta_j(L(f, T))$  are integers of absolute value at most  $\binom{2g}{k} p^{ak/2} \leq 2^{2g} p^{ak/2}$ . Since the polynomial  $\prod_{\theta_j \in G} \theta_j(L(f, T))$  has degree  $2g = (p-1)(d-1)$ , it follows that determining the coefficients modulo  $p^N$  for  $N > (p-1)(d-1)(1+a/2)$  is sufficient. (Due to a certain loss of accuracy when one performs the homological reduction, it is necessary to initially compute the coefficients of  $F$  modulo  $p^{\varepsilon(N+1)}$ , for a small positive integer  $\varepsilon$ .)

## 6. THE ALGORITHM

We now present our point counting algorithm for Type 1 Artin-Schreier curves (see Section 2 for our classification of Artin-Schreier curves).

**Algorithm 25** [Artin-Schreier - Type 1]

Input: An equation  $Z^p - Z = f(X)$  over  $\mathbb{F}_q$ , where  $f \in \mathbb{F}_q[X]$  and  $q = p^a$ .

Output: The zeta function  $Z(\tilde{C}_f, T)$  of the unique smooth projective curve birational to the affine curve defined by this equation.

Step 0: Replace  $f$  by a polynomial all of whose terms have exponents not divisible by  $p$ , in the manner explained in Note 4. Denote this new polynomial also by  $f$ . This will not change the zeta function. Set  $N := \lfloor (p-1)(d-1)(1+a/2) + 1 \rfloor$ , where  $d$  is the degree of  $f$ . Let  $\varepsilon := 4$  when  $p > 2$  and  $\varepsilon := (4d+1)$  when  $p = 2$ . We shall compute the coefficients of the numerator of the zeta function modulo  $p^N$ .

Step 1: Compute the power series  $F$  given in Definition 8 with coefficients determined modulo  $p^{\varepsilon(N+1)}$ . Let  $\alpha$  be the map on the ring  $L$  (Definition 6) defined as  $\alpha = \psi_p \circ F$  (Definition 11). Let  $H_0(\alpha)$  be the map induced on the zeroth homology  $H_0$  of the complex  $\mathcal{L}$  by  $\alpha$ .

Step 2: Let  $\pi X, \pi X^2, \dots, \pi X^{d-1}$  be the basis for the zeroth homology  $H_0$ . For each basis element  $e$ , compute the image  $H_0(\alpha)(e) \in H_0$  with coefficients

determined modulo  $p^N$ . Construct  $M$ , defined as the matrix representing the map  $H_0(\alpha)$  with respect to the basis, with coefficients determined modulo  $p^N$ . Specifically,  $M = (m_{ij})$  where  $i$  is the row index, and  $j$  the column index, and  $H_0(\alpha)(\pi X^j) = \sum_{i=1}^{d-1} m_{ij}(\pi X^i) \bmod p^N$  for  $1 \leq j \leq d-1$ .

Step 3: Compute

$$M_a := M\tau^{-1}(M)\tau^{-2}(M)\dots\tau^{-(a-1)}(M)$$

modulo  $p^N$ , where the map  $\tau$  is the lifting of Frobenius to  $A$  as given in (12). Thus  $M_a$  is a matrix for the map  $H_0(\alpha_a)$ .

Step 4: Output the rational function

$$Z(\tilde{C}_f, T) := \frac{\prod_{j=1}^{p-1} \theta_j(\det(I - M_a T))}{(1-T)(1-qT)}$$

where  $\theta_j$  is the automorphism from (11) extended to act on  $\mathbb{Z}_p[\pi][T]$  by fixing monomials.

The correctness of the algorithm follows from Corollary 23 and Lemma 24, along with the discussion of the choice of  $N$  at the end of Section 5, and the choice of  $\varepsilon$  from Lemmas 27 and 31. The matrix  $M$  is called the absolute Frobenius matrix, and  $\det(I - M_a T)$  the characteristic polynomial of Frobenius. In Section 7 we will describe exactly how this first matrix is computed, allowing us to give a complexity analysis of the algorithm in Section 8. This will complete the proof of Theorem 1 for Type 1 and 2 Artin-Schreier curves. We present the algorithm for Type 3 curves in Section 9.

**Note 26** The above algorithm can be practically improved by using the functional equation

$$T^{2g} q^g P(\tilde{C}_f, \frac{1}{qT}) = P(\tilde{C}_f, T).$$

This functional equation shows that it is enough to determine the coefficients modulo  $p^{N'}$  where  $N' := \lfloor (p-1)(d-1)(1+a/4) + 1 \rfloor$ , computing only the first half of the coefficients directly in  $P(\tilde{C}_f, T)$ , and then recovering the second half by the functional equation. Moreover, our choice of  $\varepsilon$  is perhaps rather large, especially in the case  $p = 2$ . It may be enough in practice, as observed in [23], to work initially to  $p$ -adic accuracy  $N' + \epsilon$  where  $\epsilon$  is some small variable which can be determined “experimentally” (see also Note 32).

## 7. PERFORMING THE MAIN STEPS

We shall work with elements in  $L$  with coefficients determined modulo  $p^{\varepsilon(N+1)}$ . (In the case  $p = 2$  by this we mean that given  $G \in L$  we have

that  $G = G' + p^{\varepsilon(N+1)}G''$  where  $G'$  is a known polynomial over  $A \otimes \mathbb{Q}$  and  $G''$  is a power series with coefficients in  $A$ .) If an element is given to this accuracy we say it lies in  $L \bmod p^{\varepsilon(N+1)}$ . Similarly,  $H_0$  is the free module over  $A$  when  $p > 2$ , and  $A \otimes \mathbb{Q}$  when  $p = 2$ , spanned by the basis monomials

$$(19) \quad \{\pi X, \pi X^2, \dots, \pi X^{d-1}\}.$$

We write  $s \in H_0 \bmod p^N$  if an element  $s$  is given in  $H_0$  with coefficients modulo  $p^N$ . In the next two sections we shall explain how given an element  $G \in L_{>0} \bmod p^{\varepsilon(N+1)}$  we can compute  $s \in H_0 \bmod p^N$  such that

$$G = D(r) + s \bmod p^N$$

for some  $r \in L$ . We call this process finding a “normal form” in  $L_{>0} \bmod p^{\varepsilon(N+1)}$  and say that  $s$  is “cohomologous” to  $G$ . (For an arbitrary  $G \in L_{>0}$  the choice of  $\varepsilon$  would in fact depend upon the decay rate of the coefficients of  $G$ . As such, in what follows  $\varepsilon$  should be thought of as a variable; in Lemmas 27 and 31 we determine which values for  $\varepsilon$  suffice in the cases of interest.)

**7.1. Normal forms in  $L_{>0}/D(L)$  - Case  $p > 2$ .** In the case  $p > 2$  because of the decay rate of power series in  $L$  working in  $L_{>0}/D(L)$  is particularly simple. From (15) and (18)

$$H = \pi \sum_{j=0}^d \hat{a}_j X^j$$

and thus

$$H_X = \pi \sum_{j=1}^d \hat{a}_j j X^j$$

where  $\text{ord}(\hat{a}_d d) = 0$  since  $d \neq 0$  in  $\mathbb{F}_q$ . Consider the basis monomial  $\pi^{\text{wt}(u)} X^u$  for the Banach module  $L$  where  $u \geq d$ . We have the trivial identity

$$(20) \quad \begin{aligned} \pi^{\text{wt}(u)} X^u &= \left( \pi \left( \sum_{j=1}^d \hat{a}_j j X^j \right) + X \frac{d}{dX} \right) ((\hat{a}_d d)^{-1} \pi^{\text{wt}(u)-1} X^{u-d}) \\ &\quad - \left( \pi \left( \sum_{j=1}^{d-1} \hat{a}_j j X^j \right) + X \frac{d}{dX} \right) ((\hat{a}_d d)^{-1} \pi^{\text{wt}(u)-1} X^{u-d}) =: D(r) + r' \end{aligned}$$

where  $r \in L$  and  $r' \in L_{>0}$ . Moreover,  $r'$  is a sum of monomials of degree less than  $u$  and greater or equal to  $u - d + 1$ . Now let  $G \in L_{>0} \bmod p^{\varepsilon(N+1)}$  and assume that  $b\pi^{\text{wt}(u)} X^u$  is the highest term which occurs in  $G$  for some  $b \in A$ . (Note that by the decay rate on the coefficients of elements in  $L$  we see that  $u = \mathcal{O}(\varepsilon N p d)$ .) We may suppose for our purposes that  $u \geq d$ . We can write this term as  $D(br) + br'$  where  $br'$  is a sum of monomials of degree less than  $u$  but not less than  $u - d + 1$ . Write  $G = G' + b\pi^{\text{wt}(u)} X^u$ . Then in  $L_{>0}/D(L)$  we find that  $G$  is “cohomologous” to  $G_1 := G' + br'$ . To compute  $G_1$  requires  $d$  multiplications in  $A \bmod p^{\varepsilon(N+1)}$ , and the same

number of additions (plus a little precomputation which can be ignored). Now continue in this way until the highest term in some  $G_m$  has degree less than  $d$ . Precisely, we need  $m$  at most  $u - d + 1 = \mathcal{O}(\varepsilon N p d)$ .

In this way we may find a “normal form” for any element in  $L_{>0}$ . That is, given  $G \in L_{>0} \bmod p^{\varepsilon(N+1)}$  we can write it as

$$G = D(r) + s$$

where  $r \in L$ , and  $s \in H_0 \bmod p^N$  is a linear combination with coefficients in  $A \bmod p^N$  of the basis monomials

$$\{\pi X, \pi X^2, \dots, \pi X^{d-1}\}.$$

The process above has time complexity

$$(21) \quad \tilde{\mathcal{O}}((\varepsilon N p d) d (N p a)^c)$$

bit operations where  $c$  is the exponent for multiplication as defined in Section 8.1. (Strictly speaking using the method that we describe the final factor in the bracket should be  $(\varepsilon N p a)^c$ . However, a simple analysis based upon the proof of the next lemma shows that for each coefficient  $c_u X^u$  in  $G$  where  $c_u$  is given modulo  $p^{\varepsilon(N+1)}$ , one must only keep track of the first  $N$  terms in the  $p$ -adic expansion of  $c_u$ , not including the leading zero terms.)

The next lemma justifies the choice for  $\varepsilon$  in the case  $p > 2$ .

**Lemma 27.** *Let  $e \in \{\pi X, \dots, \pi X^{d-1}\}$  with  $\alpha(e)$  cohomologous to  $s$ , a linear combination of the basis elements. To determine  $s \bmod p^N$  it is enough to compute the coefficients of  $\alpha(e)$  modulo  $p^{4(N+1)}$ .*

*Proof.* Let  $e = \pi X^j$ . By Lemma 9 terms in  $F(e)$  are of the form  $c_v X^{v+j}$  where  $\text{ord}(c_v) \geq ((p-1)/p)^2(v/d)$ . By the action of  $\psi_p$  terms in  $\alpha(e)$  are of the form  $c'_u X^u$  where  $\text{ord}(c'_u) \geq ((p-1)/p)^2((pu-j)/d)$ . Equality (20) shows that for  $u$  such that

$$\left(\frac{p-1}{p}\right)^2 \left(\frac{up-j}{d}\right) - \frac{u}{d} \geq N$$

the normal form of the term  $c'_u X^u$  vanishes modulo  $p^N$ . Thus for

$$\frac{u}{d} \geq 3(N+1) \geq \frac{N+1}{((p-1)^2/p) - 1}$$

the term  $c'_u X^u$  does not contribute to  $s \bmod p^N$ . It is easy to now check that computing the coefficients  $c'_u X^u$  for  $u/d$  less than this bound with coefficients determined modulo  $p^b$  where

$$b = 3(N+1) + (N+1) = 4(N+1)$$

is enough to determine the normal form of  $\alpha(e)$  modulo  $p^N$ .  $\square$

**7.2. Normal forms in  $L_{>0}/D(L)$  - Case  $p = 2$ .** The reduction process for  $p = 2$  is in essence the same; namely, one uses a trivial identity to reduce the degree at each step. However, the justification that it works is somewhat more involved, since the power series in  $L$  decay rather more slowly.

We shall assume that  $d\hat{a}_d = 1$ ; the more general situation just involves some notational complications, the essential point being that we always have  $\text{ord}(d\hat{a}_d) = 0$ . Let  $J_1 := J - \{d\}$  be the support set  $J$  of  $f$  excluding the element  $d$ , and  $b_j := j\hat{a}_j$ . We may assume that all non-constant terms in  $f$  have odd degree (Note 4), so each integer in  $J_1$  is odd. (The argument below in fact works provided only that  $d$  is odd, the key point being that in any case  $\text{ord}(b_j) = \text{ord}(j)$  for all  $j \in J_1$ , and this is at least 1 for  $j$  even.) In a similar manner to before we have the identity

$$(22) \quad X^u = D\left(\frac{1}{\pi}X^{u-d}\right) - \left(\frac{(u-d)}{\pi}X^{u-d} + \sum_{j \in J_1} b_j X^{u-d+j}\right).$$

This is used to reduce a power series given in finite precision in  $L_{>0}$  to its normal form, that is, a polynomial over  $A \otimes \mathbb{Q}$  of degree less than  $d$  with no constant term. As before the complexity is (21).

We must also address one theoretical problem. Let  $G := \sum_u c_u X^u \in L$ , and suppose that  $c_u X^u = D(r_u) + s_u$  with  $s_u$  a polynomial of degree less than  $d$  over  $A \otimes \mathbb{Q}$ . Then  $G = D(\sum_u r_u) + \sum_u s_u$  provided that  $|r_u|, |s_u| \rightarrow 0$  as  $u \rightarrow \infty$ . To show that these sequences indeed converge, and to get a bound on their  $p$ -adic orders, requires a more careful analysis which we now perform in a rather ad hoc fashion.

**Lemma 28.** *We may write any monomial  $X^u$  in the form  $X^u = D(r) + s$  where  $r$  is a polynomial of degree at most  $u-d$ , and  $s$  is a linear combination of the monomials  $X, X^2, \dots, X^{d-1}$ , with coefficients in  $A \otimes \mathbb{Q}$ . Moreover, the coefficients of  $s$  have  $p$ -adic order at least  $-mu - 1$  and for any  $v$  the coefficient of  $X^{u-v}$  in  $r$  has order at least  $-mv - 2$ . Here*

$$m := \frac{1}{2d + (1/2)}.$$

*Proof.* Our approach will be to show that  $X^u = D(r') + s'$  where  $r'$  satisfies the conditions in the statement of the lemma, and  $s'$  has the following property: it is a sum of terms of the form  $c_v X^{u-v}$  where  $v \geq 1$  and either  $\text{ord}(c_v) \geq -mv$ , or  $\text{ord}(c_v) \geq -mv - 1$  with  $u - v < d$ . (In particular if  $v \geq 2d + 1$  with  $\text{ord}(c_v) \geq -1$ , or  $v \geq 4d + 1$  with  $\text{ord}(c_v) \geq -2$ , then the term  $c_v X^{u-v}$  is of the required form.) The result then follows by induction. For simplicity we shall only consider the remainder term “ $s'$ ”; one may verify that the other term “ $r'$ ”, which we abbreviate as “ $*$ ”, has the required properties in all cases.

If  $u < d$  there is nothing to prove, so we assume  $u \geq d$ . By (22) if  $u - d$  is even we are done, since then  $(u - d)/\pi$  is integral.

Assume then that  $u - d$  is odd. Applying (22) we are reduced to considering the term  $((u - d)/\pi)X^{u-d}$ . If  $u < 2d$  then  $u - d < d$  and once again we are done. Thus assume  $u \geq 2d$ . Applying (22) to this new term we get that  $((u - d)/\pi)X^{u-d}$  equals  $D(*)$  plus

$$(23) \quad -\frac{(u-d)(u-2d)}{\pi^2}X^{u-2d} - \frac{u-d}{\pi} \sum_{j \in J_1} b_j X^{u-d-(d-j)}.$$

We first examine the terms in the final summation of (23). If  $u-d-(d-j) < d$  we are done. Otherwise, applying (22) we find that  $((u-d)/\pi)X^{u-d-(d-j)}$  equals  $D(*)$  plus

$$(24) \quad -\frac{(u-d)(u-3d+j)}{\pi^2}X^{u-3d+j} - \frac{u-d}{\pi} \sum_{i \in J_1} b_i X^{u-d-(d-j)-(d-i)}.$$

For the first term,  $(u-3d+j)$  is divisible by 2 and also  $u-3d+j \leq u-(2d+1)$ . Thus this term is of the required form " $c_v X^{u-v}$ " where  $\text{ord}(c_v) \geq -1$  and  $v \geq 2d+1$ . For the terms in the summation in (24) one repeatedly applies (22). The first time one uses the middle term on the righthand side of (22) one gets the term  $c_v X^{u-v}$ , say. Here the order of the coefficient  $c_v$  is  $-1$ , since  $(u-2d-(d-j)-(d-i)-\dots)$  is even. Moreover, we have  $v = 2d + (d-j) + (d-i) + \dots \geq 2d+1$ , and once again this term is of the required form. If one never uses the middle term then the resulting term  $c_v X^{u-v}$ , say, has coefficient  $c_v$  of order at least  $-1$  and  $u-v < d$ , and once again we are done.

It remains to consider the first term of (23). We may assume that  $u \geq 3d$  (for otherwise it is already of the correct form). Applying (22) to this term we get  $D(*)$  plus

$$(25) \quad \frac{(u-d)(u-2d)(u-3d)}{\pi^3}X^{u-3d} + \frac{(u-d)(u-2d)(u-3d)}{\pi^2} \sum_{j \in J_1} b_j X^{u-2d-(d-j)}.$$

The terms in the last summation are of the required form since  $(u-2d)$  is even and  $2d+(d-j) \geq 2d+1$ . For the first term of (25) applying (22) we get  $D(*)$  minus

$$(26) \quad \frac{(u-d) \dots (u-4d)}{\pi^4}X^{u-4d} + \frac{(u-d)(u-2d)(u-3d)}{\pi^3} \sum_{j \in J_1} b_j X^{u-3d-(d-j)}.$$

The first term is of the required form since  $(u-2d)(u-4d)$  is divisible by  $2^3$  and also  $4d \geq 2d+1$ . For the terms in the summation of (26) one repeatedly applies (22). The first time one uses the middle term on the righthand side of (22) one gets  $c_v X^{u-v}$ , say. Here the order of the coefficient  $c_v$  is  $-2$ , since both  $(u-2d)$  and  $(u-4d-(d-j)-(d-i)-\dots)$  are divisible by 2. Also  $v = 4d + (d-j) + (d-i) + \dots \geq 4d+1$  which shows that this term is of the required form. If one never uses the middle term then the resulting term

$c_v X^{u-v}$ , say, has coefficient  $c_v$  of order at least  $-1$ , since  $(u - 2d)$  is even, and  $u - v < d$ , and once again we are done.  $\square$

**Corollary 29.** *For  $p = 2$  the set  $\{\pi X, \dots, \pi X^{d-1}\}$  is a basis for  $L_{>0}/D(L)$ .*

*Proof.* Since  $\gamma > 1/(2 + (1/2d))$ , by Lemma 28 and a straightforward continuity argument we know that this set spans  $L_{>0}/D(L)$ . Any basis cannot have fewer than  $d - 1$  elements by consideration of the degree of the L-function  $L(f, T)$  (using (8)), and so it must be a basis.  $\square$

The above result is primarily of theoretical interest. The next lemma shows that in the case in which we are interested, denominators do not in fact occur.

**Lemma 30.** *Let  $p = 2$  and  $e \in \{\pi X, \pi X^2, \dots, \pi X^{d-1}\}$ . Then  $\alpha(e)$  is cohomologous to an element  $\sum_{j=1}^{d-1} m_j(\pi X^j)$  where  $m_j \in A$ .*

*Proof.* Dividing through by  $\pi$ , we show that  $\alpha(\pi^{-1}e)$  is cohomologous to an element  $\sum_{j=1}^{d-1} m_j X^j$  with  $m_j \in A$ . Let  $e = X^j$  where  $1 \leq j \leq d - 1$ . Then the terms in  $F(e)$  are of the form  $c_v X^{v+j}$  where  $\text{ord}(c_v) \geq v/4d$  (from Lemma 9). By the action of  $\psi_2$  the terms in  $\alpha(X^j)$  are of the form  $c'_u X^u$  where  $\text{ord}(c'_u) \geq \lceil (u/2d) - (j/4d) \rceil$ . We claim that such terms are cohomologous to polynomials of degree at most  $d - 1$  over  $A$ : If  $u \leq 2d$  the result is true by one application of (22) and induction. For  $u > 2d$  one proves the result using at most two applications of (22) and induction.  $\square$

The next lemma justifies the choice of  $\varepsilon$  in the case  $p = 2$ .

**Lemma 31.** *To compute the normal form of  $\alpha(e) \in L$  modulo  $p^N$  it is sufficient to determine the coefficients of  $\alpha(e)$  modulo  $p^{(4d+1)(N+1)}$ .*

*Proof.* Lemma 28 shows that for  $u$  such that

$$\text{ord}(c_u) - \frac{u}{2d + (1/2)} - 1 \geq N$$

the normal form of the term  $c_u X^u$  vanishes modulo  $p^N$ . Since  $\alpha(e) \in L$  terms in this power series are of the form  $c_u X^u$ , say, where  $\text{ord}(c_u) \geq u/2d$ . For  $u/2d := (4d + 1)(N + 1)$  we have  $u/(2d + (1/2)) = 4d(N + 1)$  and  $(u/2d) - (u/(2d + (1/2))) - 1 = N$ , and for  $u' > u$  we get a strict inequality in the latter. Thus, it is enough to work modulo  $p^b$  where

$$b = 4d(N + 1) + (N + 1) = (4d + 1)(N + 1).$$

$\square$

**7.3. Computing the absolute Frobenius matrix.** We now describe how to perform the main step of the algorithm, that is, constructing the matrix for the absolute Frobenius map with respect to the basis  $\{\pi X, \dots, \pi X^{d-1}\}$ . First, one may compute  $F$  with the coefficients determined modulo  $p^{\varepsilon(N+1)}$  directly from the formula in Definition 8 and the expression for  $\theta(t)$  in (14).

Working with coefficients modulo  $p^{\varepsilon(N+1)}$ , for each basis element  $e$  the polynomial  $\psi_p \circ F(e) \bmod p^{\varepsilon(N+1)}$  may be constructed. The reduction method of Sections 7.1 and 7.2 is then used to write this as a linear combination of the basis elements  $\pi X, \dots, \pi X^{d-1}$ . In this way the matrix  $M$  is found with coefficients determined modulo  $p^N$ . (Note that the entries in  $M$  are  $p$ -adic integers for all  $p$ , by Lemma 30.)

**7.4. Finding the characteristic polynomial of Frobenius.** One may compute the matrix  $M_a$  via the formula

$$(27) \quad M_a = \prod_{i=0}^{a-1} \tau^{-i}(M)$$

This is proved from Lemma 24 in the same way as [15, Lemma 26] (see also the sentence following this lemma for an alternative approach). The characteristic polynomial may then be found deterministically by computing  $\text{Tr}(M_a^k)$  for  $1 \leq k \leq d$  and using the Newton identity

$$\det(I - M_a T) = \exp\left(-\sum_{k=1}^{\infty} \frac{\text{Tr}(M_a^k)}{k} T^k\right).$$

(Alternatively, one could use an interpolation method.) The numerator of the zeta function may now be found by computing the conjugates (11) and taking a product to get a polynomial in  $\mathbb{Z}_p[T]$ .

## 8. COMPLEXITY ANALYSIS

**8.1. Exponents for ring multiplication.** Let  $B$  be some modular reduction of  $A$  and  $P$  the ring of polynomials in one variable over  $B$ . Denote by  $c$  the (deterministic) exponent for multiplication in both rings: Precisely, polynomials of degree  $\delta$  in  $P$  can be multiplied in  $\tilde{O}(\delta^c)$  operations in  $B$ , and elements in  $B$  multiplied in  $\tilde{O}((\log |B|)^c)$  bit operations. Using classical methods we take  $c = 2$ , Karatsuba's algorithm gives  $c = \log_2(3) < 1.59$ , and  $c = 1$  using Fast Fourier Transform (FFT) methods. (Note that we ignore logarithmic factors, and the space complexity is  $\tilde{O}(\delta)$ ,  $\tilde{O}(\log |B|)$  in all cases. For polynomial multiplication using FFT methods we refer to [3]; we assume as in [12, Section 5] and [10, Section 4.3] that FFT methods may be applied to  $B$ , although we do not know a convenient reference for this.) Similarly, let  $\omega$  denote the exponent for deterministic multiplication of matrices over  $B$ ; thus two  $\delta \times \delta$  matrices can be multiplied in  $\tilde{O}(\delta^\omega)$  operations in  $B$ . It will transpire that the choice of  $\omega(\leq 3)$  does not affect the overall complexity.

**8.2. Complexity of point counting algorithm.** First we must compute  $F$  with coefficients determined modulo  $p^{\varepsilon(N+1)}$ . This may be done by multiplying together the  $\mathcal{O}(d)$  polynomials  $\theta(\hat{a}_j X^j)$  in the ring " $\tilde{L}(\delta) \bmod p^{\varepsilon(N+1)}$ " for  $\delta$  as in Lemma 9. (Note that each  $\theta(t)$  can itself be constructed via multiplication of truncated power series of the form  $\exp(*t^{p^j})$ , as in [15,



Lemma 29]. This part is dominated by the computation of  $F$  itself.) From the decay rate of coefficients of power series in  $\tilde{L}(\delta)$ , we see that polynomials  $\theta(\hat{a}_j X^j) \bmod p^{\varepsilon(N+1)}$  are linear combinations of the monomials  $\pi^{\lceil \tilde{\text{wt}}(i)\delta \rceil} X^i$  for  $\tilde{\text{wt}}(i)$  bounded so that

$$\lceil \tilde{\text{wt}}(i)\delta \rceil \leq \varepsilon(N+1)(p-1).$$

There are  $\mathcal{O}(\varepsilon N p d)$  such terms. Since the coefficients of the power series  $\theta(\hat{a}_j X^j)$  lie in  $A$  it follows that the construction of  $F$  may be done in

$$(28) \quad \tilde{\mathcal{O}}((\varepsilon N p d)^c (\varepsilon N p a \log p)^c) = \tilde{\mathcal{O}}((\varepsilon^2 N^2 p^2 a d)^c)$$

bit operations. (Here  $\mathcal{O}(\varepsilon N p a \log p)$  is the bit-size of elements in the ring  $A \bmod p^{\varepsilon(N+1)}$ .) Second, finding  $\psi_p \circ F(e)$  for all  $d-1$  basis monomials requires

$$(29) \quad \tilde{\mathcal{O}}(d(\varepsilon N p d)(\varepsilon N p a)) = \tilde{\mathcal{O}}(\varepsilon^2 N^2 p^2 a d^2)$$

bit operations. Here we use the quasi-linear time method to compute the map  $\tau^{-1}$  on  $A \bmod p^{\varepsilon(N+1)}$  suggested in [12, Section 5] and [10, Step 2] (namely, precomputation of the map on each basis element  $\epsilon$  by Newton iteration). Third, we must compute a normal form for each such expression to find the coefficients in the matrix  $M$ . By the time estimate (21), we see that each column of  $M$  can be computed in

$$(30) \quad \tilde{\mathcal{O}}(\varepsilon N^{c+1} p^{c+1} d^2 a^c)$$

bit operations. We require  $d-1 = \mathcal{O}(d)$  such computations. Fourth, computing the matrix for  $M_a$  may be done using Equation (27) and the fast exponentiation method of [15, Lemma 31] in

$$(31) \quad \tilde{\mathcal{O}}(d^\omega (N p a)^c)$$

bit operations. Finally, the computation of the characteristic polynomial takes

$$(32) \quad \tilde{\mathcal{O}}(d^{\omega+1} (N p a)^c)$$

bit operations. (Computation of the products of the conjugates using (11) is absorbed in the other estimates.) Adding (28), (29),  $d \times (30)$ , (31) and (32) together and putting  $N = \mathcal{O}(p a d)$  with  $\varepsilon = \mathcal{O}(1), \mathcal{O}(d)$  we get

$$\begin{aligned} & \tilde{\mathcal{O}}(p^{4c} a^{3c} d^{4+c}) && \text{for } p > 2 \\ & \tilde{\mathcal{O}}(p^{4c} a^{3c} d^{\max(5c, c+5)}) && \text{for } p = 2 \end{aligned}$$

where  $c$  is the exponent for ring multiplication as discussed in Section 8.1. (Here we have assumed  $\omega \leq 3$ .) Using FFT methods we may take  $c = 1$ , giving the time complexity claimed in Theorem 1.

The space complexity is in all cases determined by the size of the ring  $L \bmod p^{\varepsilon(N+1)}$  and also the polynomial  $F \bmod p^{\varepsilon(N+1)}$ . These are both

$$\tilde{\mathcal{O}}((\varepsilon N p d)(\varepsilon N p a)) = \begin{cases} \tilde{\mathcal{O}}(p^4 a^3 d^3) & \text{for } p > 2 \\ \tilde{\mathcal{O}}(a^3 d^5) & \text{for } p = 2. \end{cases}$$

This completes the proof of Theorem 1 in the case of Type 1 Artin-Schreier curves, and also Type 2 since they are easily reduced to Type 1.

**Note 32** It is possible to reduce the factors  $d^{5c}$  and  $d^5$  in the time and space complexities, respectively, for  $p = 2$  to  $d^{4c}$  and  $d^4$  using a more careful analysis: we have not taken into account that only part ( $N$  terms) of the  $p$ -adic expansion of each coefficient in  $F \bmod p^{\varepsilon(N+1)}$  needs to be computed. A much more detailed analysis of the action of  $D$  could perhaps reduce the value required for  $\varepsilon$  when  $p = 2$ , giving a uniform time estimate of  $\tilde{O}(p^4 a^3 d^5)$  with space  $\tilde{O}(p^4 a^3 d^3)$  for all  $p$ .

**Note 33** We briefly describe one alternative approach for  $p = 2$  which has also been implemented by Vercauteren [23]: The idea is to use a more complicated splitting function to improve the decay rate of the coefficients in  $F$ . The result is that one may use  $L := \tilde{L}(1) \cap A\{X\}$  for the case  $p = 2$ , and the proof that  $L_{>0}/D(L)$  becomes easier. Specifically, setting  $p = 2$  take  $\theta := \theta_3$  where  $\theta_3$  is the splitting function defined in Lemma 6 of [15]. One may compute the required element  $\gamma_3 \in \mathbb{Z}_p$  in the following manner: let  $\gamma'_3$  satisfy the relation

$$\gamma'_3 = 1 + 4\gamma'^3_3 - 8\gamma'^4_3 + 8\gamma'^5_3.$$

Then  $\gamma_3 := 2\gamma'_3$ . Let  $\alpha$  be defined as before, but with the new  $\theta$ . This time it is stable on  $L := \tilde{L}(1) \cap A\{X\}$ . Define  $H$  to be

$$H := \sum_{j=0}^2 \gamma_{3,j} \tau^j(\hat{f}(X^{p^j}))$$

where

$$\gamma_{3,j} := \sum_{i=0}^j \frac{\gamma_3^{p^i}}{p^i}.$$

Then one must compute the action of  $\alpha$  on the homology  $L_{>0}/D(L)$  where  $D := X(d/dX) + H_X$ . To do this, for  $j \geq 0$  define  $\pi^j L_{>0}$  to be the Banach module over  $A$  with orthonormal basis  $\pi^{\text{wt}(u)+j} X^u$  ( $u > 0$ ). Then

$$H_X \equiv \pi X \frac{d\hat{f}}{dX} \bmod \pi L_{>0}.$$

Now one performs the reduction process from Section 7.1 coupled with a hensel lifting argument to reveal “ $r$ ” and “ $s$ ” with “ $G = D(r) + s$ ” working modulo  $\pi^j L_{>0}$  for increasing powers of  $j$ . Note that one may take any  $N \geq (d-1)(1 + (a/2)) + 1$  and  $\varepsilon = 11/3$ , and all computations are done with  $p$ -adic integers. The algorithm has complexity  $\tilde{O}(a^{\max(3c,4)} d^6)$  where  $c$  is the exponent for multiplication and space complexity  $\tilde{O}(a^3 d^3)$ . This is slightly slower in terms of  $a$  due to a step in the hensel lifting for which we were unable to find a fast method.

## 9. ARTIN-SCHREIER COVERS OF THE TORUS

In this section we present in a much more condensed fashion the algorithm for computing the zeta function of the Type 3 Artin-Schreier curves.

Let  $f \in \mathbb{F}_q[X, X^{-1}]$  have negative degree  $d^- < 0$  and positive degree  $d^+ > 0$ . We shall explain how to compute the L-function  $L^*(f, T)$ . From this one may easily compute the zeta functions of Type 3 Artin-Schreier curves using (9).

Define a weight function  $\text{wt}$  on  $\mathbb{Z}$  by

$$\text{wt}(u) := \begin{cases} \lceil u/d^- \rceil & \text{if } u < 0 \\ \lceil u/d^+ \rceil & \text{if } u \geq 0 \end{cases}$$

where  $|\cdot|$  is the usual (not  $p$ -adic) absolute value. For  $p > 2$  let  $L$  be the Banach module over  $A$  with orthonormal basis  $\pi^{\text{wt}(r)} X^r$  for  $r \in \mathbb{Z}$ . For  $p = 2$  let  $L$  denote the Banach space over  $A \otimes \mathbb{Q}$  with orthonormal basis  $\pi^{\lceil \gamma \text{wt}(r) \rceil} X^r$  for  $r \in \mathbb{Z}$ . (Here  $\gamma$  is any rational number with  $1/2 > \gamma > 1/(2 + 1/(2d'))$  where  $d' := \max(|d^-|, d^+)$ .) Note that  $L$  contains two-way infinite power series, and in fact still forms a ring because of the decay conditions on coefficients. Now define the power series  $F$  and  $F^{(a)}$  in exactly the same manner as before. In this case they are two-way infinite power series. With  $\alpha$  and  $\alpha_a$  defined exactly as before, we find that both maps are stable on  $L$ . We have the formula

$$L^*(f, T) = \frac{\det(1 - T\alpha_a|L)}{\det(1 - Tq\alpha|L)}.$$

In this case one cannot remove any unit root factors. Let  $\mathcal{L}$  be the (slightly “larger”) complex of modules

$$0 \longrightarrow L \xrightarrow{D} L \longrightarrow 0.$$

These are  $A$ -modules for  $p > 2$  and  $A \otimes \mathbb{Q}$ -spaces for  $p = 2$ . Here  $D$  is defined in exactly the same manner as before (in Section 5). In this case  $H_0 = L/D(L)$  and  $H_1 = 0$ , and one recovers the cohomological trace formula

$$L^*(f, T) = \det(1 - H_0(\alpha_a)T).$$

Once again we have the crucial relation  $H_0(\alpha_a) = H_0(\alpha)^a$ . A basis for  $H_0$  can be taken as

$$\{\pi X^{d^-}, \pi X^{d^-+1}, \dots, \pi X^{-1}, 1, \pi X, \dots, \pi X^{d^+-1}\}$$

with  $d^+ - d^-$  the dimension of this space. Computation of normal forms in  $L/D(L)$  is done in a similar manner to before. Precisely, one first uses the method in Section 7.1 and 7.2 to find an element cohomologous to a given element whose leading term has degree less than  $d^+$ . One then performs a similar process to increase the degree of the lowest term so that it is not less than  $d^-$ . Note that we need a final  $p$ -adic accuracy of any  $N$  greater than  $(p-1)(d^+ - d^-)(a/2 + 1)$ . For  $p > 2$  the factor  $\varepsilon$  can be taken to be 4, and for  $p = 2$  it is  $4d' + 1$ . The complexity of the above algorithm may be checked to be identical to that in the case when  $f$  is just an ordinary polynomial.

Precisely, if we write “ $d := d^+ - d^-$ ” then all the bounds in Section 8 are still true. This completes the proof of Theorem 1, and Corollary 2 follows from Note 5.

## REFERENCES

- [1] A. Adolphson and S. Sperber, Exponential sums and Newton polyhedra: Cohomology and estimates, *Ann. Math.* 130, (1989) 367-406.
- [2] I. Blake, G. Seroussi, N. Smart, *Elliptic Curves in Cryptography*, LMS Lecture Note Series 265, Cambridge University Press, 1999.
- [3] D. Cantor and E. Kaltofen, On fast multiplication of polynomials over arbitrary algebras, *Acta Inform.* 28 (7), (1991), 693-701.
- [4] R.F. Coleman,  $p$ -adic Banach spaces and families of modular forms, *Invent. Math.* 127, (1997), 417-479.
- [5] B. Dwork, On the rationality of the zeta function of an algebraic variety, *Amer. J. Math.*, 82, (1960), 631-648.
- [6] B. Dwork, On the zeta function of a hypersurface, *Pub. Math. IHES* No. 12, 1962.
- [7] N. Elkies, Elliptic and modular curves over finite fields and related computational issues, in “*Computational perspectives in number theory: Proceedings of a conference in honour of A.O.L. Atkin*”, (D.A. Buell and J.T. Teitelbaum), American Mathematical Society International Press 7, 1998, 21-76.
- [8] M. Fouquet, P. Gaudry and R. Harley, An extension of Satoh’s algorithm and its implementation, *J. Ramanujan Math. Soc.* 15, (2000), 281-318.
- [9] M. Fouquet, P. Gaudry and R. Harley, Finding secure curves with the Satoh-FGH algorithm and an early abort strategy, in B. Pfitzmann (ed), *Advances in Cryptology - EUROCRYPT 2001*, Lecture Notes in Computer Science 2045, Springer-Verlag, (2001), 14-29.
- [10] P. Gaudry and N. Gürel, An extension of Kedlaya’s algorithm for counting points on superelliptic curves, preprint 2001.
- [11] P. Gaudry and R. Harley, Counting points on hyperelliptic curves over finite fields, in B. Preneel (ed), *Advances in Cryptology - EUROCRYPT 2000*, Lecture notes in Computer Science 1807, Springer-Verlag, (2000), 19-34.
- [12] K.S. Kedlaya, Counting points on hyperelliptic curves using Monsky-Washnitzer cohomology, preprint 2001.
- [13] N. Koblitz,  $p$ -Adic Numbers,  $p$ -Adic Analysis and Zeta Functions, 2nd Edition, Springer, 1984.
- [14] N. Koblitz, Hyperelliptic cryptosystems, *J. of Cryptology* 1, (1989), 139-150.
- [15] A.G.B. Lauder and D. Wan, Counting points on varieties over finite fields of small characteristic, preprint 2001.
- [16] R. Lidl and H. Niederreiter, *Introduction to Finite Fields and Their Applications*, Cambridge University Press, 1986.
- [17] B. Poonen, Computational aspects of curves of genus at least 2, in “*Algorithmic Number Theory II*” (H. Cohen), Lecture Notes in Computer Science 1122, Springer, 1996, 283-306.
- [18] T. Satoh, The canonical lift of an ordinary elliptic curve over a finite fields and its points counting, *J. Ramanujan Math. Soc.* 15, (2000), 247-270.
- [19] R. Schoof, Elliptic curves over finite fields and the computation of square roots mod  $p$ , *Math. Comp.* 44, no. 170, (1985), 483-494.
- [20] J.-P. Serre, Endomorphismes complètement continus des espaces de Banach  $p$ -adique, *Pub. Math. IHES* 12, (1962), 69-85.
- [21] S. Sperber, On the  $p$ -adic theory of exponential sums, *Amer. J. Math.* 108, (1983), 255-296.

- [22] F. Vercauteren, B. Preneel and J. Vandewalle, A memory efficient version of Satoh's algorithm, in B. Pfitzmann (ed) *Advances in Cryptology - EUROCRYPT 2001*, Lecture Notes in Computer Science 2045, Springer-Verlag, (2001), 1-13.
- [23] F. Vercauteren, personal communication, 2001.
- [24] D. Wan, Computing zeta functions over finite fields, *Contemporary Mathematics*, 225 (1999), 131-141.
- [25] D. Wan, Pure L-functions from algebraic geometry over finite fields, in "Finite Fields and Applications" D. Jungnickel and H. Niederreiter (eds), Springer-Verlag, (2000), 437-461.
- [26] D. Wan, Algorithmic theory of zeta functions over finite fields, preprint 2001.

COMPUTING LABORATORY, OXFORD UNIVERSITY, OXFORD OX1 3QD, UK     *E-mail address:* ALAN.LAUDER@COMLAB.OX.AC.UK

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, IRVINE, CA 92697, USA     *E-mail address:* DWAN@MATH.UCI.EDU