HUMBOLDT-UNIVERSITÄT ZU BERLIN

Regularization of divergent Feynman integrals and their computation with hyperlogarithms

Erik Panzer[1]

Institutes of Physics and Mathematics
Humboldt-Universität zu Berlin
Unter den Linden 6
10099 Berlin, Germany

ETH Zürich (Hönggerberg), March 18th, 2014
Particle Physics Seminar

[1] panzer@mathematik.hu-berlin.de

Plan of the talk

1. The hyperlogarithm method of integration

Plan of the talk

1. The hyperlogarithm method of integration
2. Examples of linearly reducible Feynman integrals

Plan of the talk

1. The hyperlogarithm method of integration
2. Examples of linearly reducible Feynman integrals
3. Divergences and analytic regularization

Motivation: Feynman integrals in Schwinger parameters

For scalar propagators $P_e = (p_e^2 + m_e^2)^{-a_e}$, Feynman integrals become

$$\Phi(G) = \frac{\Gamma(\text{sdd})}{\prod_e \Gamma(a_e)} \int_{\mathbb{R}_+^{|E|}} F\ \Omega \quad \text{with} \quad F = \psi^{\text{sdd}-D/2} \cdot \varphi^{-\text{sdd}} \cdot \prod_{e \in E} \alpha_e^{a_e - 1}.$$

Motivation: Feynman integrals in Schwinger parameters

For scalar propagators $P_e = (p_e^2 + m_e^2)^{-a_e}$, Feynman integrals become

$$\Phi(G) = \frac{\Gamma(\mathrm{sdd})}{\prod_e \Gamma(a_e)} \int_{\mathbb{R}_+^{|E|}} F\ \Omega \quad \text{with} \quad F = \psi^{\mathrm{sdd}-D/2} \cdot \varphi^{-\mathrm{sdd}} \cdot \prod_{e \in E} \alpha_e^{a_e - 1}.$$

The graph polynomials ψ, φ are sums over all spanning trees T and all spanning two-forests F with external momentum $q(T_1)$ entering T_1:

$$\psi = \sum_T \prod_{e \notin T} \alpha_e \quad \text{and} \quad \varphi = \sum_{F = T_1 \dot\cup T_2} q^2(T_1) \prod_{e \notin F} \alpha_e + \psi \sum_e m_e^2 \alpha_e.$$

Their homogeneity is exploited in that $\Omega = \delta(H) \cdot \bigwedge_e \mathrm{d}\alpha_e$ restricts to a hyperplane H, e.g. $H = 1 - \alpha_{|E|}$. When G has $|G|$ loops,

$$\mathrm{sdd} = \sum_e a_e - \frac{D}{2}|G| \quad \text{denotes its superficial degree of divergence.}$$

Motivation: Feynman integrals in Schwinger parameters

For scalar propagators $P_e = (p_e^2 + m_e^2)^{-a_e}$, Feynman integrals become

$$\Phi(G) = \frac{\Gamma(\mathrm{sdd})}{\prod_e \Gamma(a_e)} \int_{\mathbb{R}_+^{|E|}} F\ \Omega \quad \text{with} \quad F = \psi^{\mathrm{sdd}-D/2} \cdot \varphi^{-\mathrm{sdd}} \cdot \prod_{e \in E} \alpha_e^{a_e - 1}.$$

The graph polynomials ψ, φ are sums over all spanning trees T and all spanning two-forests F with external momentum $q(T_1)$ entering T_1:

$$\psi = \sum_T \prod_{e \notin T} \alpha_e \quad \text{and} \quad \varphi = \sum_{F = T_1 \dot\cup T_2} q^2(T_1) \prod_{e \notin F} \alpha_e + \psi \sum_e m_e^2 \alpha_e.$$

Their homogeneity is exploited in that $\Omega = \delta(H) \cdot \bigwedge_e \mathrm{d}\alpha_e$ restricts to a hyperplane H, e.g. $H = 1 - \alpha_{|E|}$. When G has $|G|$ loops,

$$\mathrm{sdd} = \sum_e a_e - \frac{D}{2}|G| \quad \text{denotes its superficial degree of divergence.}$$

Idea: Expand $F = \sum_n \varepsilon^n F_n$ with $F_n \in \mathbb{Q}[\alpha_e, \psi^{-1}, \varphi^{-1}, \ln\psi, \ln\varphi, \ln\alpha_e]$ and compute the integrals $\int F_n\ \Omega$ along a sequence $\alpha_1, \alpha_2, \ldots, \alpha_{|E|-1}$ of edges.

Hyperlogarithms for Feynman integrals, F. Brown [9, 8, 7]

Definition

For any word $w \in \Sigma^{\times}$ in letters $\{\omega_\sigma : \ \sigma \in \Sigma\}$ over a finite set $0 \in \Sigma \subset \mathbb{C}$, the associated *hyperlogarithm* $L_w(z)$ is defined by

$$L_{\omega_0^n}(z) := \frac{\log^n z}{n!} \quad \text{for any } n \in \mathbb{N}_0 \text{ and } \quad L_{\omega_\sigma w}(z) := \int_0^z \frac{\mathrm{d}z'}{z' - \sigma} L_w(z').$$

Hyperlogarithms for Feynman integrals, F. Brown [9, 8, 7]

Definition

For any word $w \in \Sigma^{\times}$ in letters $\{\omega_\sigma : \ \sigma \in \Sigma\}$ over a finite set $0 \in \Sigma \subset \mathbb{C}$, the associated *hyperlogarithm* $L_w(z)$ is defined by

$$L_{\omega_0^n}(z) := \frac{\log^n z}{n!} \quad \text{for any } n \in \mathbb{N}_0 \text{ and } \quad L_{\omega_\sigma w}(z) := \int_0^z \frac{\mathrm{d}z'}{z' - \sigma} L_w(z').$$

Alternative names

Goncharov polylogarithms, *generalized harmonic polylogarithms*:

$$L_{\omega_{\sigma_1} \ldots \omega_{\sigma_n}}(z) = G\left(\sigma_1, \ldots, \sigma_n; z\right).$$

Hyperlogarithms for Feynman integrals, F. Brown [9, 8, 7]

Definition

For any word $w \in \Sigma^{\times}$ in letters $\{\omega_\sigma : \ \sigma \in \Sigma\}$ over a finite set $0 \in \Sigma \subset \mathbb{C}$, the associated *hyperlogarithm* $L_w(z)$ is defined by

$$L_{\omega_0^n}(z) := \frac{\log^n z}{n!} \quad \text{for any } n \in \mathbb{N}_0 \text{ and } \quad L_{\omega_\sigma w}(z) := \int_0^z \frac{\mathrm{d}z'}{z' - \sigma} L_w(z').$$

Example (Relation to classical and multiple polylogarithms)

For any $0 \neq \sigma \in \Sigma$ and $n \in \mathbb{N}$ note

$$L_{\omega_\sigma}(z) = \log(z - \sigma) - \log(-\sigma) = -\operatorname{Li}_1\left(\frac{z}{\sigma}\right), \quad L_{\omega_0^{n-1}\omega_\sigma}(z) = -\operatorname{Li}_n\left(\frac{z}{\sigma}\right).$$

More generally, for $\sigma_1, \ldots, \sigma_r \in \Sigma \setminus \{0\}$ and $n_1, \ldots, n_r \in \mathbb{N}$ we have

$$L_{\omega_0^{n_r-1}\omega_{\sigma_r} \cdots \omega_0^{n_2-1}\omega_{\sigma_2}\omega_0^{n_1-1}\omega_{\sigma_1}}(z) = (-1)^r \operatorname{Li}_{n_1,\ldots,n_r}\left(\frac{\sigma_2}{\sigma_1}, \ldots, \frac{\sigma_r}{\sigma_{r-1}}, \frac{z}{\sigma_r}\right).$$

Hyperlogarithms for Feynman integrals, F. Brown [9, 8, 7]

Properties

1. Algebra: $L_. : \mathbb{Q}\langle\Sigma\rangle \longrightarrow \mathcal{O}(\mathbb{C} \setminus \Sigma)$ (extended linearly) is a morphism of algebras for the *shuffle product*: $L_u \cdot L_v = L_{u \sqcup\!\sqcup v}$. For example,

$$L_{\omega_\sigma} \cdot L_{\omega_{\tau_1}\omega_{\tau_2}} = L_{\omega_\sigma\omega_{\tau_1}\omega_{\tau_2}} + L_{\omega_{\tau_1}\omega_\sigma\omega_{\tau_2}} + L_{\omega_{\tau_1}\omega_{\tau_2}\omega_\sigma}.$$

Hyperlogarithms for Feynman integrals, F. Brown [9, 8, 7]

Properties

1. Algebra: $L_\cdot : \mathbb{Q}\langle\Sigma\rangle \longrightarrow \mathcal{O}(\mathbb{C} \setminus \Sigma)$ (extended linearly) is a morphism of algebras for the *shuffle product*: $L_u \cdot L_v = L_{u \sqcup\!\sqcup v}$. For example,

$$L_{\omega_\sigma} \cdot L_{\omega_{\tau_1}\omega_{\tau_2}} = L_{\omega_\sigma\omega_{\tau_1}\omega_{\tau_2}} + L_{\omega_{\tau_1}\omega_\sigma\omega_{\tau_2}} + L_{\omega_{\tau_1}\omega_{\tau_2}\omega_\sigma}.$$

2. Grading: by *weight* $|w| := n$ for all $w = \omega_{\sigma_1} \ldots \omega_{\sigma_n} \in \Sigma^n$

Hyperlogarithms for Feynman integrals, F. Brown [9, 8, 7]

Properties

1. Algebra: $L_. : \mathbb{Q}\langle\Sigma\rangle \longrightarrow \mathcal{O}(\mathbb{C} \setminus \Sigma)$ (extended linearly) is a morphism of algebras for the *shuffle product*: $L_u \cdot L_v = L_{u \sqcup\!\sqcup v}$. For example,

$$L_{\omega_\sigma} \cdot L_{\omega_{\tau_1}\omega_{\tau_2}} = L_{\omega_\sigma\omega_{\tau_1}\omega_{\tau_2}} + L_{\omega_{\tau_1}\omega_\sigma\omega_{\tau_2}} + L_{\omega_{\tau_1}\omega_{\tau_2}\omega_\sigma}.$$

2. Grading: by *weight* $|w| := n$ for all $w = \omega_{\sigma_1} \dots \omega_{\sigma_n} \in \Sigma^n$
3. Independence: $\{L_w : \ w \in \Sigma^\times\}$ are linearly independent over algebraic functions $\overline{\mathbb{C}(z)}$.

Hyperlogarithms for Feynman integrals, F. Brown [9, 8, 7]

Properties

1. Algebra: $L_. : \mathbb{Q}\langle\Sigma\rangle \longrightarrow \mathcal{O}(\mathbb{C} \setminus \Sigma)$ (extended linearly) is a morphism of algebras for the *shuffle product*: $L_u \cdot L_v = L_{u \sqcup\!\sqcup v}$. For example,

$$L_{\omega_\sigma} \cdot L_{\omega_{\tau_1}\omega_{\tau_2}} = L_{\omega_\sigma\omega_{\tau_1}\omega_{\tau_2}} + L_{\omega_{\tau_1}\omega_\sigma\omega_{\tau_2}} + L_{\omega_{\tau_1}\omega_{\tau_2}\omega_\sigma}.$$

2. Grading: by *weight* $|w| := n$ for all $w = \omega_{\sigma_1} \ldots \omega_{\sigma_n} \in \Sigma^n$
3. Independence: $\{L_w : \ w \in \Sigma^\times\}$ are linearly independent over algebraic functions $\overline{\mathbb{C}(z)}$.
4. Monodromies: For paths $\gamma, \eta : [0,1] \longrightarrow \mathbb{C} \setminus \Sigma$ with $\gamma(1) = \eta(0)$,

$$\int_{\gamma\star\eta} \omega_{\sigma_1} \ldots \omega_{\sigma_n} = \sum_{k=0}^{n} \int_\eta \omega_{\sigma_1} \ldots \omega_{\sigma_k} \int_\gamma \omega_{\sigma_{k+1}} \ldots \omega_{\sigma_n} \quad \text{(Chen's lemma)}$$

Hyperlogarithms for Feynman integrals, F. Brown [9, 8, 7]

Properties

1. Algebra: $L_. : \mathbb{Q}\langle\Sigma\rangle \longrightarrow \mathcal{O}(\mathbb{C} \setminus \Sigma)$ (extended linearly) is a morphism of algebras for the *shuffle product*: $L_u \cdot L_v = L_{u \sqcup\!\sqcup v}$. For example,
$$L_{\omega_\sigma} \cdot L_{\omega_{\tau_1}\omega_{\tau_2}} = L_{\omega_\sigma\omega_{\tau_1}\omega_{\tau_2}} + L_{\omega_{\tau_1}\omega_\sigma\omega_{\tau_2}} + L_{\omega_{\tau_1}\omega_{\tau_2}\omega_\sigma}.$$
2. Grading: by *weight* $|w| := n$ for all $w = \omega_{\sigma_1} \ldots \omega_{\sigma_n} \in \Sigma^n$
3. Independence: $\{L_w : \ w \in \Sigma^\times\}$ are linearly independent over algebraic functions $\overline{\mathbb{C}(z)}$.
4. Monodromies: For paths $\gamma, \eta : [0,1] \longrightarrow \mathbb{C} \setminus \Sigma$ with $\gamma(1) = \eta(0)$,
$$\int_{\gamma\star\eta} \omega_{\sigma_1} \ldots \omega_{\sigma_n} = \sum_{k=0}^{n} \int_\eta \omega_{\sigma_1} \ldots \omega_{\sigma_k} \int_\gamma \omega_{\sigma_{k+1}} \ldots \omega_{\sigma_n} \quad \text{(Chen's lemma)}$$
5. Divergences: For any $w \in \Sigma^\times$ and $\sigma \in \Sigma \cup \{\infty\}$ exist unique $f_w^{\sigma,k}(z)$, $0 \leq k \leq |w|$, holomorphic at $z \to \sigma$ such that
$$L_w(z) = \sum_k \log^k(z - \sigma) \cdot f_w^{\sigma,k}(z).$$

Hyperlogarithms for Feynman integrals, F. Brown [9, 8, 7]

Integration

By construction and use of partial fractioning, every integrand

$$f(z) \in L(\Sigma) := \mathbb{Q}\left[z, \left\{\frac{1}{z-\sigma} : \ \sigma \in \Sigma\right\}, \{L_w : \ w \in \Sigma^\times\}\right]$$

has a primitive $F(z)$ in the enlarged algebra

$$R(\Sigma) := \mathbb{Q}\left[\Sigma \cup \left\{\frac{1}{\sigma_i - \sigma_j} : \ \sigma_i \neq \sigma_j \text{ from } \Sigma\right\}\right] \otimes L(\Sigma).$$

Hyperlogarithms for Feynman integrals, F. Brown [9, 8, 7]

Integration

By construction and use of partial fractioning, every integrand

$$f(z) \in L(\Sigma) := \mathbb{Q}\left[z, \left\{\frac{1}{z-\sigma} : \ \sigma \in \Sigma\right\}, \{L_w : \ w \in \Sigma^\times\}\right]$$

has a primitive $F(z)$ in the enlarged algebra

$$R(\Sigma) := \mathbb{Q}\left[\Sigma \cup \left\{\frac{1}{\sigma_i - \sigma_j} : \ \sigma_i \neq \sigma_j \text{ from } \Sigma\right\}\right] \otimes L(\Sigma).$$

Example

$$\int \frac{L_{\omega_\sigma}(z)\mathrm{d}z}{(z-\tau)^2} = -\frac{L_{\omega_\sigma}(z)}{z-\tau} + \int \frac{\mathrm{d}z}{(z-\tau)(z-\sigma)} = -\frac{L_{\omega_\sigma}(z)}{z-\tau} + \frac{L_{\omega_\tau}(z) - L_{\omega_\sigma}(z)}{\tau-\sigma}$$

Hyperlogarithms for Feynman integrals, F. Brown [9, 8, 7]

Integration

By construction and use of partial fractioning, every integrand

$$f(z) \in L(\Sigma) := \mathbb{Q}\left[z, \left\{\frac{1}{z-\sigma} : \ \sigma \in \Sigma\right\}, \{L_w : \ w \in \Sigma^\times\}\right]$$

has a primitive $F(z)$ in the enlarged algebra

$$R(\Sigma) := \mathbb{Q}\left[\Sigma \cup \left\{\frac{1}{\sigma_i - \sigma_j} : \ \sigma_i \neq \sigma_j \text{ from } \Sigma\right\}\right] \otimes L(\Sigma).$$

Example

$$\int \frac{L_{\omega_\sigma}(z)\mathrm{d}z}{(z-\tau)^2} = -\frac{L_{\omega_\sigma}(z)}{z-\tau} + \int \frac{\mathrm{d}z}{(z-\tau)(z-\sigma)} = -\frac{L_{\omega_\sigma}(z)}{z-\tau} + \frac{L_{\omega_\tau}(z) - L_{\omega_\sigma}(z)}{\tau-\sigma}$$

For the following integration, we must extract the dependence of

$$\int_0^\infty f(z)\mathrm{d}z = \lim_{z\to\infty} F(z) - \lim_{z\to 0} F(z)$$

on the next Schwinger variable.

Hyperlogarithms for Feynman integrals, F. Brown [9, 8, 7]

Rewriting polylogarithms as hyperlogarithms

Consider $w = \omega_{\sigma_1} \dots \omega_{\sigma_n}$ with $\sigma_1, \dots, \sigma_n \in \mathbb{Q}(t)$ depending rationally on a variable t (the next Schwinger parameter α_e to integrate):

$$\frac{\partial}{\partial t} L_w(z) = \frac{-\sigma_1'}{z - \sigma_1} L_{\omega_{\sigma_2} \dots}(z) + \sum_{i=1}^{n-1} \frac{(\sigma_i - \sigma_{i+1})'}{\sigma_i - \sigma_{i+1}} L_{\dots \not{\omega}_{\sigma_{i+1}} \dots - \dots \not{\omega}_{\sigma_i} \dots}(z) - \frac{\sigma_n'}{\sigma_n} L_{\dots \not{\omega}_{\sigma_n}}(z)$$

Hyperlogarithms for Feynman integrals, F. Brown [9, 8, 7]

Rewriting polylogarithms as hyperlogarithms

Consider $w = \omega_{\sigma_1} \dots \omega_{\sigma_n}$ with $\sigma_1, \dots, \sigma_n \in \mathbb{Q}(t)$ depending rationally on a variable t (the next Schwinger parameter α_e to integrate):

$$\frac{\partial}{\partial t} L_w(z) = \frac{-\sigma_1'}{z - \sigma_1} L_{\omega_{\sigma_2} \dots}(z) + \sum_{i=1}^{n-1} \frac{(\sigma_i - \sigma_{i+1})'}{\sigma_i - \sigma_{i+1}} L_{\dots \not{\omega}_{\sigma_{i+1}} \dots - \dots \not{\omega}_{\sigma_i} \dots}(z) - \frac{\sigma_n'}{\sigma_n} L_{\dots \not{\omega}_{\sigma_n}}(z)$$

Example ($\lim_{z\to\infty} L_{(\omega_{-t}-\omega_{-1})\omega_{-1}}(z)$)

$$\frac{\partial}{\partial t} \lim_{z\to\infty} \left[L_{\omega_{-t}\omega_{-1}}(z) - L_{\omega_{-1}\omega_{-1}}(z) \right] = \left[\frac{\partial}{\partial t} \log(t-1) \right] \cdot \lim_{z\to\infty} L_{\omega_{-t}-\omega_{-1}}(z)$$

$$= \frac{1}{t-1} \lim_{z\to\infty} \log \frac{(z+t)}{t(z+1)} = -\frac{1}{t-1} L_{\omega_0}(t)$$

Hyperlogarithms for Feynman integrals, F. Brown [9, 8, 7]

Rewriting polylogarithms as hyperlogarithms

Consider $w = \omega_{\sigma_1} \dots \omega_{\sigma_n}$ with $\sigma_1, \dots, \sigma_n \in \mathbb{Q}(t)$ depending rationally on a variable t (the next Schwinger parameter α_e to integrate):

$$\frac{\partial}{\partial t} L_w(z) = \frac{-\sigma_1'}{z - \sigma_1} L_{\omega_{\sigma_2} \dots}(z) + \sum_{i=1}^{n-1} \frac{(\sigma_i - \sigma_{i+1})'}{\sigma_i - \sigma_{i+1}} L_{\dots \not{\omega}_{\sigma_{i+1}} \dots - \dots \not{\omega}_{\sigma_i} \dots}(z) - \frac{\sigma_n'}{\sigma_n} L_{\dots \not{\omega}_{\sigma_n}}(z)$$

Example ($\lim_{z\to\infty} L_{(\omega_{-t}-\omega_{-1})\omega_{-1}}(z) = C - L_{\omega_1\omega_0}(t)$)

$$\frac{\partial}{\partial t} \lim_{z\to\infty} \left[L_{\omega_{-t}\omega_{-1}}(z) - L_{\omega_{-1}\omega_{-1}}(z) \right] = \left[\frac{\partial}{\partial t} \log(t-1) \right] \cdot \lim_{z\to\infty} L_{\omega_{-t}-\omega_{-1}}(z)$$

$$= \frac{1}{t-1} \lim_{z\to\infty} \log \frac{(z+t)}{t(z+1)} = -\frac{1}{t-1} L_{\omega_0}(t)$$

Hyperlogarithms for Feynman integrals, F. Brown [9, 8, 7]

Rewriting polylogarithms as hyperlogarithms

Consider $w = \omega_{\sigma_1} \dots \omega_{\sigma_n}$ with $\sigma_1, \dots, \sigma_n \in \mathbb{Q}(t)$ depending rationally on a variable t (the next Schwinger parameter α_e to integrate):

$$\frac{\partial}{\partial t} L_w(z) = \frac{-\sigma_1'}{z - \sigma_1} L_{\omega_{\sigma_2} \dots}(z) + \sum_{i=1}^{n-1} \frac{(\sigma_i - \sigma_{i+1})'}{\sigma_i - \sigma_{i+1}} L_{\dots \not{\omega}_{\sigma_{i+1}} \dots - \dots \not{\omega}_{\sigma_i} \dots}(z) - \frac{\sigma_n'}{\sigma_n} L_{\dots \not{\omega}_{\sigma_n}}(z)$$

Example ($\lim_{z\to\infty} L_{(\omega_{-t}-\omega_{-1})\omega_{-1}}(z) = C - L_{\omega_1\omega_0}(t)$)

$$\frac{\partial}{\partial t} \lim_{z\to\infty} \left[L_{\omega_{-t}\omega_{-1}}(z) - L_{\omega_{-1}\omega_{-1}}(z) \right] = \left[\frac{\partial}{\partial t} \log(t-1) \right] \cdot \lim_{z\to\infty} L_{\omega_{-t}-\omega_{-1}}(z)$$

$$= \frac{1}{t-1} \lim_{z\to\infty} \log \frac{(z+t)}{t(z+1)} = -\frac{1}{t-1} L_{\omega_0}(t)$$

The constants C of integration can be computed symbolically as well in `HyperInt` [15].

Linear reducibility

This method applies only if for all $n < N$, the partial integrals

$$f_n := \left[\prod_{e=1}^{n-1} \int_0^\infty \mathrm{d}\alpha_e\right] F \quad \text{lie in} \quad L(\Sigma_n)(\alpha_n) \tag{1.1}$$

for some sets $\Sigma_n \subset \mathbb{Q}(\alpha_{n+1}, \ldots, \alpha_N)$. In particular, at the next stage the differences $\sigma_i - \sigma_j$ have to factor linearly in α_{n+1} for all $\sigma_i, \sigma_j \in \Sigma_n$.

Linear reducibility

This method applies only if for all $n < N$, the partial integrals

$$f_n := \left[\prod_{e=1}^{n-1} \int_0^\infty \mathrm{d}\alpha_e\right] F \quad \text{lie in} \quad L(\Sigma_n)(\alpha_n) \tag{1.1}$$

for some sets $\Sigma_n \subset \mathbb{Q}(\alpha_{n+1}, \ldots, \alpha_N)$. In particular, at the next stage the differences $\sigma_i - \sigma_j$ have to factor linearly in α_{n+1} for all $\sigma_i, \sigma_j \in \Sigma_n$.

Definition

G is called *linearly reducible* if for some ordering $e_1, \ldots, e_N$ of its edges there exist sets Σ_n for all $0 \leq n < N$ such that (1.1) holds.

Linear reducibility

This method applies only if for all $n < N$, the partial integrals

$$f_n := \left[\prod_{e=1}^{n-1} \int_0^\infty \mathrm{d}\alpha_e\right] F \quad \text{lie in} \quad L(\Sigma_n)(\alpha_n) \tag{1.1}$$

for some sets $\Sigma_n \subset \mathbb{Q}(\alpha_{n+1}, \ldots, \alpha_N)$. In particular, at the next stage the differences $\sigma_i - \sigma_j$ have to factor linearly in α_{n+1} for all $\sigma_i, \sigma_j \in \Sigma_n$.

Definition

G is called *linearly reducible* if for some ordering $e_1, \ldots, e_N$ of its edges there exist sets Σ_n for all $0 \leq n < N$ such that (1.1) holds.

Polynomial reduction

A simple algorithm providing sufficient conditions for linear reducibility is given by Stembridge's procedure as explained in [9]. Much more powerful is the method of compatibility graphs developed in [8].

Are there linearly reducible Feynman graphs?

Single-scale integrals

Massless propagators: Coefficients of ε-expansions

Theorem (positive matrix graphs [8])

All graphs G of vertex-width $\mathrm{vw}(G) \leq 3$ are linearly reducible and their periods lie in $\mathcal{Z} := \mathrm{lin}_{\mathbb{Q}} \{\zeta(n_1, \ldots, n_r) : \ n_1, \ldots, n_r \in \mathbb{N} \text{ and } n_r > 1\}$.



Single-scale integrals

Massless propagators: Coefficients of ε-expansions

Theorem (positive matrix graphs [8])

All graphs G of vertex-width $\mathrm{vw}(G) \leq 3$ are linearly reducible and their periods lie in $\mathcal{Z} := \mathrm{lin}_{\mathbb{Q}} \{\zeta(n_1, \ldots, n_r) : \ n_1, \ldots, n_r \in \mathbb{N} \text{ and } n_r > 1\}$.



Theorem (vacuum graphs with four or five loops [14])

All massless propagators up to four loops are linearly reducible. Their periods are constrained to alternating Euler sums $\mathcal{Z}_2$:

$$\mathcal{Z}_2 := \mathrm{lin}_{\mathbb{Q}} \left\{ \mathrm{Li}_{n_1,\ldots,n_r}(\sigma_1, \ldots, \sigma_r) : \ n_i \in \mathbb{N}, \sigma_i^2 = 1 \forall i \text{ and } (n_r, \sigma_r) \neq (1,1) \right\}.$$



Massless on-shell four-point graphs

Massless on-shell four-point graphs

Theorem ([6])

All massless four-point on-shell graphs ($p_1^2 = p_2^2 = p_3^2 = p_4^2 = 0$) *with at most two loops are linearly reducible. In particular these include*



Massless on-shell four-point graphs

Theorem ([6])

All massless four-point on-shell graphs ($p_1^2 = p_2^2 = p_3^2 = p_4^2 = 0$) with at most two loops are linearly reducible. In particular these include



$K_4 =$ 

At three loops, there are non-reducible graphs characterized by forbidden minors like K_4.

Massless on-shell four-point graphs

Theorem ([6])

All massless four-point on-shell graphs ($p_1^2 = p_2^2 = p_3^2 = p_4^2 = 0$) with at most two loops are linearly reducible. In particular these include



$K_4 =$ 

At three loops, there are non-reducible graphs characterized by forbidden minors like K_4.
Still, K_4 evaluates to polylogarithms [12]. Linear reducibility can be restored through a change of variables, hence parametric integration of K_4 is possible.

Massless on-shell four-point graphs

Linearly reducible four-loop example

Φ



$$= \frac{\Gamma(1+4\varepsilon)}{s^{1+4\varepsilon}} \sum_{n=-1}^{\infty} f_n\left(\frac{s}{u}\right) \cdot \varepsilon^n$$

is linearly reducible along the sequence $1, 2, 8, 6, 9, 7, 5, 4$ of edges ($\alpha_3 = 1$) with final set $\{s + u\}$, so all f_n are harmonic polylogarithms of $\frac{s}{u}$:

Massless on-shell four-point graphs

Linearly reducible four-loop example

$\Phi\Big($ 

$\Big) = \dfrac{\Gamma(1+4\varepsilon)}{s^{1+4\varepsilon}} \sum_{n=-1}^{\infty} f_n\left(\dfrac{s}{u}\right)\cdot\varepsilon^n$

is linearly reducible along the sequence $1,2,8,6,9,7,5,4$ of edges ($\alpha_3 = 1$) with final set $\{s+u\}$, so all f_n are harmonic polylogarithms of $\frac{s}{u}$:

$$
\begin{aligned}
f_{-1}={}&-\tfrac{79}{70}\zeta_2^3 H_{-1}-\zeta_3\big(15\zeta_2 H_{-1,-1}-9\zeta_2 H_{-1,0}-H_{-1,-2,-1}+H_{-1,-1,-2}+6H_{-1,-1,0,0}\big)\\
&-6\zeta_3^2 H_{-1}-\tfrac{3}{2}\zeta_5(11H_{-1,-1}-5H_{-1,0})-\tfrac{3}{10}\zeta_2^2(H_{-1,-2}-17H_{-1,-1,0}-10H_{-1,-1,-1})\\
&-\zeta_2\Big(H_{-1,-2,0,0}-2H_{-1,-1,-2,0}+3H_{-1,-1,-2,-1}-H_{-1,-1,-1,0,0}+6H_{-1,-1,-3}\\
&\qquad-3H_{-1,-2,-1,-1}-2H_{-1,-1,0,0,0}\Big)+H_{-1,-2,-1,0,0,0}-H_{-1,-1,-2,-1,0,0}\\
&+H_{-1,-1,-2,0,0,0}-2H_{-1,-1,-3,0,0}+H_{-1,-2,-1,-1,0,0}
\end{aligned}
$$

Off-shell massless vertices

Consider zero internal masses $m_e = 0$ and three external momenta like in

$\Delta :=$ 

where

$$\psi = \alpha_1 + \alpha_2 + \alpha_3 \text{ and}$$
$$\varphi = p_1^2\alpha_2\alpha_3 + p_2^2\alpha_1\alpha_3 + p_3^3\alpha_1\alpha_2.$$

Off-shell massless vertices

Consider zero internal masses $m_e = 0$ and three external momenta like in

$\Delta :=$ 

where

$$\psi = \alpha_1 + \alpha_2 + \alpha_3 \text{ and}$$
$$\varphi = p_1^2\alpha_2\alpha_3 + p_2^2\alpha_1\alpha_3 + p_3^3\alpha_1\alpha_2.$$

To rationalize the square-root of the Källén function we parametrize as

$$p_2^2 = p_1^2 \cdot z\bar{z} \qquad \text{and} \qquad p_3^2 = p_1^2 \cdot (1-z)(1-\bar{z}), \qquad \text{such that}$$
$$z - \bar{z} = \sqrt{p_1^2 + p_2^2 + p_3^2 - 2p_1p_2 - 2p_1p_3 - 2p_2p_3}.$$

Off-shell massless vertices

Consider zero internal masses $m_e = 0$ and three external momenta like in

$\Delta :=$ 

where $\psi = \alpha_1 + \alpha_2 + \alpha_3$ and
$\varphi = p_1^2\alpha_2\alpha_3 + p_2^2\alpha_1\alpha_3 + p_3^3\alpha_1\alpha_2.$

To rationalize the square-root of the Källén function we parametrize as

$$p_2^2 = p_1^2 \cdot z\bar{z} \qquad \text{and} \qquad p_3^2 = p_1^2 \cdot (1-z)(1-\bar{z}), \qquad \text{such that}$$

$$z - \bar{z} = \sqrt{p_1^2 + p_2^2 + p_3^2 - 2p_1p_2 - 2p_1p_3 - 2p_2p_3}.$$

Then the triangle G_1 expands near $D = 4 - 2\varepsilon$ with $a_1 = a_2 = a_3 = 1$ as

$$\Phi(\Delta) = \frac{\Gamma(1+\varepsilon)}{z-\bar{z}} \cdot p_1^{-2(1+\varepsilon)} \cdot \sum_{n=0}^{\infty} f_n(z,\bar{z})\varepsilon^n, \qquad \text{at leading order}$$

$f_0 = 4i\Im\left\{\mathrm{Li}_2(z) + \ln|z| \cdot \ln(1-z)\right\}$ is the Bloch-Wigner dilogarithm.

Off-shell massless vertices

The functions f_n have symbols with letters in $\{z, \bar{z}, 1-z, 1-\bar{z}, z-\bar{z}\}$ and are generalized single-valued multiple polylogarithms (SVMP) [17], e.g.

$$f_1 = 4i\Im\left\{\mathrm{Li}_{1,2}\left(\frac{z}{\bar{z}}, \bar{z}\right) + 2\,\mathrm{Li}_3(z) + \mathrm{Li}_{2,1}(1,z) + \mathrm{Li}_{1,2}(1,z) \right.$$
$$\left. + \ln|z| \cdot \left[\mathrm{Li}_{1,1}\left(\frac{\bar{z}}{z}, z\right) - 2\,\mathrm{Li}_2(z) - \ln|z(1-z)| \cdot \ln(1-z)\right]\right\}.$$

Off-shell massless vertices

The functions f_n have symbols with letters in $\{z, \bar{z}, 1-z, 1-\bar{z}, z-\bar{z}\}$ and are generalized single-valued multiple polylogarithms (SVMP) [17], e.g.

$$f_1 = 4i\Im\left\{\mathrm{Li}_{1,2}\left(\frac{z}{\bar{z}}, \bar{z}\right) + 2\,\mathrm{Li}_3(z) + \mathrm{Li}_{2,1}(1,z) + \mathrm{Li}_{1,2}(1,z) \right.$$
$$\left. + \ln|z| \cdot \left[\mathrm{Li}_{1,1}\left(\frac{\bar{z}}{z}, z\right) - 2\,\mathrm{Li}_2(z) - \ln|z(1-z)| \cdot \ln(1-z)\right]\right\}.$$

Linear reducibility at two loops was observed in [10] and we checked

Theorem ([16])

All three-loop off-shell massless three-point functions are linearly reducible. Some require additional letters $\{z\bar{z}-1, z+\bar{z}-1, z\bar{z}-z-\bar{z}\}$.

Off-shell massless vertices

The functions f_n have symbols with letters in $\{z, \bar{z}, 1-z, 1-\bar{z}, z-\bar{z}\}$ and are generalized single-valued multiple polylogarithms (SVMP) [17], e.g.

$$f_1 = 4i\Im\left\{\mathrm{Li}_{1,2}\left(\frac{z}{\bar{z}}, \bar{z}\right) + 2\,\mathrm{Li}_3(z) + \mathrm{Li}_{2,1}(1,z) + \mathrm{Li}_{1,2}(1,z) \right.$$
$$\left. + \ln|z| \cdot \left[\mathrm{Li}_{1,1}\left(\frac{\bar{z}}{z}, z\right) - 2\,\mathrm{Li}_2(z) - \ln|z(1-z)| \cdot \ln(1-z)\right]\right\}.$$

Linear reducibility at two loops was observed in [10] and we checked

Theorem ([16])

All three-loop off-shell massless three-point functions are linearly reducible. Some require additional letters $\{z\bar{z}-1, z+\bar{z}-1, z\bar{z}-z-\bar{z}\}$.

Also at higher loop numbers we found linearly reducible graphs like



Non-trivial kinematics: up to 7 scales

Examples: massive integrals

Convention:

- thin lines: light-like momenta $p_e^2 = 0$, massless propagators $m_e = 0$
- thick lines: arbitrary masses m_e and external momenta p_e^2 (in particular the masses may be all different)

Then the following graphs are linearly reducible:



Non-trivial kinematics: up to 7 scales

Example: double triangle with three legs off-shell and two massive propagators

$$\Phi\left(\begin{array}{c}\\ \\ \end{array}\right.$$


$$\left.\begin{array}{c}\\ \\ \end{array}\right) = \Gamma(1+2\varepsilon)\int \frac{\Omega}{\varphi^{1+2\varepsilon}\psi^{1+\varepsilon}}$$

is linearly reducible.

Example: double triangle with three legs off-shell and two massive propagators



$$\Phi\left(\cdots \right) = \frac{\Gamma(1+2\varepsilon) q^{-1-2\varepsilon}}{1+p-s-u} \sum_{n=-1}^{\infty} f_n(s,u,p,m,M)\varepsilon^n$$

is linearly reducible. We simplify to $p_4^2 = 0$, let $q := p_3^2$ and set

$$p := \frac{p_1^2}{q}, s := \frac{(p_1+p_2)^2}{q}, u := \frac{(p_1+p_4)^2}{q}, M := \frac{m_3^2}{q}, m := \frac{m_4^2}{q} \text{ in}$$

Non-trivial kinematics: up to 7 scales

Example: double triangle with three legs off-shell and two massive propagators



$$\Phi\left(\text{graph}\right) = \frac{\Gamma(1+2\varepsilon) q^{-1-2\varepsilon}}{1+p-s-u} \sum_{n=-1}^{\infty} f_n(s,u,p,m,M)\varepsilon^n$$

is linearly reducible. We simplify to $p_4^2 = 0$, let $q := p_3^2$ and set

$$p := \frac{p_1^2}{q}, s := \frac{(p_1+p_2)^2}{q}, u := \frac{(p_1+p_4)^2}{q}, M := \frac{m_3^2}{q}, m := \frac{m_4^2}{q} \text{ in}$$

$$\begin{aligned}
f_{-1} =& \mathrm{Li}_{1,1,1}\left(\tfrac{1-u+p-s}{(1-s)(1-u)}, \tfrac{M(1-s)}{M-m}, \tfrac{M-m}{M}\right) - \mathrm{Li}_{1,1,1}\left(\tfrac{M(1-u+p-s)}{(M-m)(1-u)}, \tfrac{(u+M)(M-m)}{(-p-m+u+M)M}, \tfrac{-p-m+u+M}{u+M}\right) \\
&+ \mathrm{Li}_{1,1,1}\left(-\tfrac{u(1-u+p-s)}{(-u+p)(1-u)}, -\tfrac{(u+M)(-u+p)}{(-p-m+u+M)u}, \tfrac{-p-m+u+M}{u+M}\right) - \ln\left(\tfrac{m}{p+m}\right)\mathrm{Li}_{1,1}\left(-\tfrac{u(1-u+p-s)}{(-u+p)(1-u)}, -\tfrac{-u+p}{u}\right) \\
&- \mathrm{Li}_{1,1,1}\left(-\tfrac{u(1-u+p-s)}{(-u+p)(1-u)}, -\tfrac{M(-u+p)}{(M-m)u}, \tfrac{M-m}{M}\right) - \mathrm{Li}_{1,1,1}\left(\tfrac{1-u+p-s}{(1-s)(1-u)}, \tfrac{(1+M)(1-s)}{-m-s+1+M}, \tfrac{-m-s+1+M}{1+M}\right) \\
&+ \mathrm{Li}_{1,1,1}\left(\tfrac{M(1-u+p-s)}{(M-m)(1-u)}, \tfrac{(1+M)(M-m)}{(-m-s+1+M)M}, \tfrac{-m-s+1+M}{1+M}\right) - \ln\left(-\tfrac{p-s}{1-u}\right)\mathrm{Li}_2\left(\tfrac{p}{p+m}\right) + \ln\left(-\tfrac{p-s}{1-u}\right)\mathrm{Li}_2\left(\tfrac{s}{m+s}\right) \\
&+ \ln\left(\tfrac{m}{m+s}\right)\mathrm{Li}_{1,1}\left(\tfrac{1-u+p-s}{(1-s)(1-u)}, 1-s\right) - \ln\left(\tfrac{m}{m+s}\right)\mathrm{Li}_{1,1}\left(\tfrac{M(1-u+p-s)}{(M-m)(1-u)}, \tfrac{M-m}{M}\right) + \tfrac{1}{2}\ln\left(-\tfrac{p-s}{1-u}\right)\ln^2\left(\tfrac{m}{m+s}\right) \\
&- \tfrac{1}{2}\ln\left(-\tfrac{p-s}{1-u}\right)\ln^2\left(\tfrac{m}{p+m}\right) + \ln\left(\tfrac{m}{p+m}\right)\mathrm{Li}_{1,1}\left(\tfrac{M(1-u+p-s)}{(M-m)(1-u)}, \tfrac{M-m}{M}\right)
\end{aligned}$$

Analytic regularization of parametric integrals

Divergences in Schwinger parameters

Example: Vertex with one null momentum and one internal mass

Consider a vertex graph G with one internal mass and $p_3^2 = 0$:

$G =$ 

$$\psi = \alpha_1 + \alpha_2 + \alpha_3$$
$$\varphi = \alpha_3 \left(m^2 \psi + p_1^2 \alpha_2 + p_2^2 \alpha_1 \right)$$

Divergences in Schwinger parameters

Example: Vertex with one null momentum and one internal mass

Consider a vertex graph G with one internal mass and $p_3^2 = 0$:

$G =$ 

$$\psi = \alpha_1 + \alpha_2 + \alpha_3$$
$$\varphi = \alpha_3 \left(m^2 \psi + p_1^2 \alpha_2 + p_2^2 \alpha_1 \right)$$

It is linearly reducible, but the parametric integral in the representation

$$\Phi(G) = \int \frac{\mathrm{d}^D k}{\pi^{D/2}} \frac{1}{(k^2 + m^2)^{a_3} (k + p_2)^{2a_1} (k - p_1)^{2a_2}}$$
$$= \frac{\Gamma(\mathrm{sdd})}{\Gamma(a_1)\Gamma(a_2)\Gamma(a_3)} \int \frac{\alpha_1^{a_1 - 1} \alpha_2^{a_2 - 1} \alpha_3^{a_3 - 1}}{\psi^{D/2 - \mathrm{sdd}} \varphi^{\mathrm{sdd}}} \Omega$$

diverges at $\varepsilon \to 0$ for $D = 4 - 2\varepsilon$ and $a_1 = a_2 = a_3 = 1$.

Divergences in Schwinger parameters

Example: Vertex with one null momentum and one internal mass

Consider a vertex graph G with one internal mass and $p_3^2 = 0$:

$G =$ 

$$\psi = \alpha_1 + \alpha_2 + \alpha_3$$
$$\varphi = \alpha_3 \left(m^2\psi + p_1^2\alpha_2 + p_2^2\alpha_1\right)$$

It is linearly reducible, but the parametric integral in the representation

$$\Phi(G) = \int \frac{\mathrm{d}^D k}{\pi^{D/2}} \frac{1}{(k^2+m^2)^{a_3}(k+p_2)^{2a_1}(k-p_1)^{2a_2}}$$
$$= \frac{\Gamma(\mathrm{sdd})}{\Gamma(a_1)\Gamma(a_2)\Gamma(a_3)} \int \frac{\alpha_1^{a_1-1}\alpha_2^{a_2-1}\alpha_3^{a_3-1}}{\psi^{D/2-\mathrm{sdd}}\varphi^{\mathrm{sdd}}}\Omega$$

diverges at $\varepsilon \to 0$ for $D = 4 - 2\varepsilon$ and $a_1 = a_2 = a_3 = 1$. Then $\mathrm{sdd} = 1 + \varepsilon$ and the integral $\int_0^\infty \mathrm{d}\alpha_3$ diverges at the lower boundary:

$$\int \frac{\Omega}{(\alpha_1+\alpha_2+\alpha_3)^{1-2\varepsilon}\left[m^2(\alpha_1+\alpha_2+\alpha_3)+p_1^2\alpha_2+p_2^2\alpha_1\right]^{1+\varepsilon}\alpha_3^{1+\varepsilon}}$$

Divergences in Schwinger parameters: Power counting

$$F = \prod_{e \in E} \alpha_e^{a_e - 1} \cdot \psi^{\mathrm{sdd} - D/2} \cdot \varphi^{-\mathrm{sdd}}$$

Definition

For an integrand F and disjoint sets $J, K \subset E$ of edges, the rescaling

$$F_J^K := F \circ \left(\alpha_e \mapsto \begin{cases} \lambda \alpha_e, & e \in J \\ \lambda^{-1} \alpha_e, & e \in K \\ \alpha_e, & e \in E \setminus (J \cup K) \end{cases} \right)$$

defines a vanishing degree $\deg_J^K$ such that $F_J^K = \lambda^{\deg_J^K} \cdot \widetilde{F_J^K}$ where $\lim_{\lambda \to 0} \widetilde{F_J^K} \neq 0$ exists. Set $\omega_J^K := |J| - |K| + \deg_J^K$.

Divergences in Schwinger parameters: Power counting

$$F = \prod_{e \in E} \alpha_e^{a_e - 1} \cdot \psi^{\mathrm{sdd} - D/2} \cdot \varphi^{-\mathrm{sdd}}$$

Definition

For an integrand F and disjoint sets $J, K \subset E$ of edges, the rescaling

$$F_J^K := F \circ \left(\alpha_e \mapsto \begin{cases} \lambda \alpha_e, & e \in J \\ \lambda^{-1} \alpha_e, & e \in K \\ \alpha_e, & e \in E \setminus (J \cup K) \end{cases} \right)$$

defines a vanishing degree $\deg_J^K$ such that $F_J^K = \lambda^{\deg_J^K} \cdot \widetilde{F_J^K}$ where $\lim_{\lambda \to 0} \widetilde{F_J^K} \neq 0$ exists. Set $\omega_J^K := |J| - |K| + \deg_J^K$.

Theorem (Finiteness)

When all coefficients of φ are positive and $\omega_J^K > 0$ for all disjoint $J, K \subsetneq E$ with $0 \neq J \cup K \subsetneq E$, then $\int F \, \Omega$ converges absolutely.

Analytic regularization in Schwinger parameters

A partial integration yields the relation

$$\int_0^\infty \frac{\mathrm{d}\lambda}{\lambda}\lambda^{\omega_J^K}\cdot\widetilde{F_J^K}(\lambda)=\frac{\lambda^{\omega_J^K}}{\omega_J^K}\widetilde{F_J^K}(\lambda)\Big|_{\lambda=0}^{\infty}-\frac{1}{\omega_J^K}\int_0^\infty \mathrm{d}\lambda\cdot\lambda^{\omega_J^K}\frac{\partial}{\partial\lambda}\widetilde{F_J^K}(\lambda)$$

with vanishing boundary contribution in the convergent regime.
The analytically regularized functions associated to both integrals thus coincide (they are meromorphic in the regulators).

Analytic regularization in Schwinger parameters

A partial integration yields the relation

$$\int_0^\infty \frac{\mathrm{d}\lambda}{\lambda}\lambda^{\omega_J^K}\cdot\widetilde{F_J^K}(\lambda) = \frac{\lambda^{\omega_J^K}}{\omega_J^K}\widetilde{F_J^K}(\lambda)\Big|_{\lambda=0}^{\infty} - \frac{1}{\omega_J^K}\int_0^\infty \mathrm{d}\lambda\cdot\lambda^{\omega_J^K}\frac{\partial}{\partial\lambda}\widetilde{F_J^K}(\lambda)$$

with vanishing boundary contribution in the convergent regime.
The analytically regularized functions associated to both integrals thus coincide (they are meromorphic in the regulators).

Example (Vertex with one mass and $p_3^2 = 0$ in $D = 4 - 2\varepsilon$)

We have $\omega_{\{3\}}^{\emptyset} = -\varepsilon$ with $\widetilde{F_{\{3\}}^{\emptyset}} = \psi^{2\varepsilon-1}\cdot\left[m^2\psi + p_2^2\alpha_1 + p_1^2\alpha_2\right]^{-1-\varepsilon}$:

$$\int\frac{\Omega}{\psi^{1-2\varepsilon}\varphi^{1+\varepsilon}} = \frac{\alpha_3^{-\varepsilon}\widetilde{F_{\{3\}}^{\emptyset}}}{-\varepsilon}\Big|_{\alpha_3=0}^{\infty} + \frac{1}{\varepsilon}\cdot\int\frac{\Omega}{\alpha_3^{\varepsilon}}\frac{\partial}{\partial\alpha_3}\widetilde{F_{\{3\}}^{\emptyset}}$$

$$= \frac{1}{\varepsilon}\cdot\int\frac{\Omega\alpha_3}{\psi^{1-2\varepsilon}\varphi^{1+\varepsilon}}\left[\frac{2\varepsilon-1}{\psi} - \frac{(1+\varepsilon)\alpha_3 m^2}{\varphi}\right] \text{ when } \varepsilon < 0.$$

Regularization in Schwinger parameters

General construction

Proposition

For disjoint subsets $J, K \subsetneq E$, acting with the differential operator

$$\mathcal{D}_J^K := 1 - \frac{1}{\omega_J^K}\Big[\sum_{e\in J}\frac{\partial}{\partial\alpha_e}\alpha_e - \sum_{e\in K}\frac{\partial}{\partial\alpha_e}\alpha_e\Big]$$

on the integrand F yields a new integrand $\widetilde{F} := \mathcal{D}_J^K(F)$ such that

Proposition

For disjoint subsets $J, K \subsetneq E$, acting with the differential operator

$$\mathcal{D}_J^K := 1 - \frac{1}{\omega_J^K}\Big[\sum_{e\in J}\frac{\partial}{\partial\alpha_e}\alpha_e - \sum_{e\in K}\frac{\partial}{\partial\alpha_e}\alpha_e\Big]$$

on the integrand F yields a new integrand $\widetilde{F} := \mathcal{D}_J^K(F)$ such that

1. $\int F\ \Omega = \int \widetilde{F}\ \Omega$ *as analytically regularized integrals,*

Regularization in Schwinger parameters

General construction

Proposition

For disjoint subsets $J, K \subsetneq E$, acting with the differential operator

$$\mathcal{D}_J^K := 1 - \frac{1}{\omega_J^K}\Big[\sum_{e\in J}\frac{\partial}{\partial\alpha_e}\alpha_e - \sum_{e\in K}\frac{\partial}{\partial\alpha_e}\alpha_e\Big]$$

on the integrand F yields a new integrand $\widetilde{F} := \mathcal{D}_J^K(F)$ such that

1. $\int F\ \Omega = \int \widetilde{F}\ \Omega$ *as analytically regularized integrals,*
2. $\omega_{J'}^{K'}(\widetilde{F}) \geq \omega_{J'}^{K'}(F)$ *for all* $J' \cap K' = \emptyset$ *and*

Regularization in Schwinger parameters

General construction

Proposition

For disjoint subsets $J, K \subsetneq E$, acting with the differential operator

$$\mathcal{D}_J^K := 1 - \frac{1}{\omega_J^K}\Big[\sum_{e\in J}\frac{\partial}{\partial\alpha_e}\alpha_e - \sum_{e\in K}\frac{\partial}{\partial\alpha_e}\alpha_e\Big]$$

on the integrand F yields a new integrand $\widetilde{F} := \mathcal{D}_J^K(F)$ such that

1. $\int F\ \Omega = \int \widetilde{F}\ \Omega$ *as analytically regularized integrals,*
2. $\omega_{J'}^{K'}(\widetilde{F}) \geq \omega_{J'}^{K'}(F)$ *for all* $J' \cap K' = \emptyset$ *and*
3. $\omega_J^K\left(\widetilde{F}\right) \geq 1 + \omega_J^K(F)$ *increases at least by one.*

Regularization in Schwinger parameters

General construction

Proposition

For disjoint subsets $J, K \subsetneq E$, acting with the differential operator

$$\mathcal{D}_J^K := 1 - \frac{1}{\omega_J^K}\Big[\sum_{e\in J}\frac{\partial}{\partial\alpha_e}\alpha_e - \sum_{e\in K}\frac{\partial}{\partial\alpha_e}\alpha_e\Big]$$

on the integrand F yields a new integrand $\widetilde{F} := \mathcal{D}_J^K(F)$ such that

1. $\int F\ \Omega = \int \widetilde{F}\ \Omega$ *as analytically regularized integrals,*
2. $\omega_{J'}^{K'}(\widetilde{F}) \geq \omega_{J'}^{K'}(F)$ *for all* $J' \cap K' = \emptyset$ *and*
3. $\omega_J^K\left(\widetilde{F}\right) \geq 1 + \omega_J^K(F)$ *increases at least by one.*

Thus finitely many partial integrations suffice to make all ω_J^K positive, thereby yielding a convergent integral representation (suitable for hyperlogarithmic integration).

Proposition

For disjoint subsets $J, K \subsetneq E$, acting with the differential operator

$$\mathcal{D}_J^K := 1 - \frac{1}{\omega_J^K}\Big[\sum_{e\in J}\frac{\partial}{\partial\alpha_e}\alpha_e - \sum_{e\in K}\frac{\partial}{\partial\alpha_e}\alpha_e\Big]$$

on the integrand F yields a new integrand $\widetilde{F} := \mathcal{D}_J^K(F)$ such that

1. $\int F\ \Omega = \int \widetilde{F}\ \Omega$ *as analytically regularized integrals,*
2. $\omega_{J'}^{K'}(\widetilde{F}) \geq \omega_{J'}^{K'}(F)$ *for all* $J' \cap K' = \emptyset$ *and*
3. $\omega_J^K\left(\widetilde{F}\right) \geq 1 + \omega_J^K(F)$ *increases at least by one.*

Thus finitely many partial integrations suffice to make all ω_J^K positive, thereby yielding a convergent integral representation (suitable for hyperlogarithmic integration).

This procedure does not introduce new singularities, i.e. the polynomial reduction and linear reducibility is not affected.

Summary: Hyperlogarithmic integration

1. If a graph G (with specification of kinematics) is *linearly reducible*, it can be integrated with hyperlogarithms:
   - to any order in ε, expansion near arbitrary even dimension $D|_{\varepsilon=0} \in 2\mathbb{N}$
   - with any tensor structures
   - for arbitrary powers $a_e = n_e + \varepsilon\nu_e$ of propagators ($n_e \in \mathbb{Z}, \nu_e \in \mathbb{C}$)

Summary: Hyperlogarithmic integration

1. If a graph G (with specification of kinematics) is *linearly reducible*, it can be integrated with hyperlogarithms:
   - to any order in ε, expansion near arbitrary even dimension $D|_{\varepsilon=0} \in 2\mathbb{N}$
   - with any tensor structures
   - for arbitrary powers $a_e = n_e + \varepsilon\nu_e$ of propagators ($n_e \in \mathbb{Z}, \nu_e \in \mathbb{C}$)
2. Applications also to other parametric integrals, e.g. [4, 3].

Summary: Hyperlogarithmic integration

1. If a graph G (with specification of kinematics) is *linearly reducible*, it can be integrated with hyperlogarithms:
   - to any order in ε, expansion near arbitrary even dimension $D|_{\varepsilon=0} \in 2\mathbb{N}$
   - with any tensor structures
   - for arbitrary powers $a_e = n_e + \varepsilon\nu_e$ of propagators $(n_e \in \mathbb{Z}, \nu_e \in \mathbb{C})$
2. Applications also to other parametric integrals, e.g. [4, 3].
3. A convergent representation is needed and can always be constructed in analytic (dimensional) regularization using partial integrations.
   But for highly divergent integrals, this generates huge expressions and a more economic way of subtracting divergences seems desireable.

Summary: Hyperlogarithmic integration

1. If a graph G (with specification of kinematics) is *linearly reducible*, it can be integrated with hyperlogarithms:
   - to any order in ε, expansion near arbitrary even dimension $D|_{\varepsilon=0} \in 2\mathbb{N}$
   - with any tensor structures
   - for arbitrary powers $a_e = n_e + \varepsilon\nu_e$ of propagators $(n_e \in \mathbb{Z}, \nu_e \in \mathbb{C})$
2. Applications also to other parametric integrals, e.g. [4, 3].
3. A convergent representation is needed and can always be constructed in analytic (dimensional) regularization using partial integrations. But for highly divergent integrals, this generates huge expressions and a more economic way of subtracting divergences seems desireable.
4. Sometimes a graph can become linearly reducible by a change of variables (in particular for massive integrals). A systematic study of such transformations on Schwinger parameters is still lacking.

Summary: Hyperlogarithmic integration

1. If a graph G (with specification of kinematics) is *linearly reducible*, it can be integrated with hyperlogarithms:
   - to any order in ε, expansion near arbitrary even dimension $D|_{\varepsilon=0} \in 2\mathbb{N}$
   - with any tensor structures
   - for arbitrary powers $a_e = n_e + \varepsilon\nu_e$ of propagators ($n_e \in \mathbb{Z}, \nu_e \in \mathbb{C}$)
2. Applications also to other parametric integrals, e.g. [4, 3].
3. A convergent representation is needed and can always be constructed in analytic (dimensional) regularization using partial integrations. But for highly divergent integrals, this generates huge expressions and a more economic way of subtracting divergences seems desireable.
4. Sometimes a graph can become linearly reducible by a change of variables (in particular for massive integrals). A systematic study of such transformations on Schwinger parameters is still lacking.
5. Linear reducibility is not fully understood combinatorially, but tests on individual graphs/integrands are automated.

Summary: Hyperlogarithmic integration

1. If a graph G (with specification of kinematics) is *linearly reducible*, it can be integrated with hyperlogarithms:
   - to any order in ε, expansion near arbitrary even dimension $D|_{\varepsilon=0} \in 2\mathbb{N}$
   - with any tensor structures
   - for arbitrary powers $a_e = n_e + \varepsilon\nu_e$ of propagators ($n_e \in \mathbb{Z}, \nu_e \in \mathbb{C}$)
2. Applications also to other parametric integrals, e.g. [4, 3].
3. A convergent representation is needed and can always be constructed in analytic (dimensional) regularization using partial integrations. But for highly divergent integrals, this generates huge expressions and a more economic way of subtracting divergences seems desireable.
4. Sometimes a graph can become linearly reducible by a change of variables (in particular for massive integrals). A systematic study of such transformations on Schwinger parameters is still lacking.
5. Linear reducibility is not fully understood combinatorially, but tests on individual graphs/integrands are automated.
6. The package `HyperInt` [15] implements the algorithms in `Maple`$^{\text{TM}}$.

Additional slides

Single-scale integrals

Massive on-shell operator elements

Theorem ([2, 5, 1])

The following on-shell ($p^2 = 0$) propagators with equally massive fermions are linearly reducible:



Single-scale integrals

Massive on-shell operator elements

Theorem ([2, 5, 1])

The following on-shell ($p^2 = 0$) propagators with equally massive fermions are linearly reducible:

$\hat{I}_4 =$ 

In fact, the calculation used a two-scale setup to get all Mellin moments:

$$\tilde{I}_4(x) := \int \frac{\Omega}{\varphi^2} \sum_{0=j\leq N}^{\infty} x^N \cdot \frac{T_{4a}^{N-j} T_{4b}^{j}}{\psi^N} = \int \frac{\Omega}{(\psi - xT_{4a})(\psi - xT_{4b})M^2},$$

where $T_{4a}, T_{4b}, \psi \in \mathbb{Z}[\alpha_1, \ldots, \alpha_8]$ are cubic polynomials and $M := \sum_{m_e \neq 0} \alpha_e$.

Single-scale integrals

Massive on-shell operator elements

Theorem ([2, 5, 1])

The following on-shell ($p^2 = 0$) propagators with equally massive fermions are linearly reducible:

$\hat{I}_4 =$ 

In fact, the calculation used a two-scale setup to get all Mellin moments:

$$\tilde{I}_4(x) := \int \frac{\Omega}{\varphi^2} \sum_{0=j\leq N}^{\infty} x^N \cdot \frac{T_{4a}^{N-j} T_{4b}^{j}}{\psi^N} = \int \frac{\Omega}{(\psi - xT_{4a})(\psi - xT_{4b})M^2},$$

where $T_{4a}, T_{4b}, \psi \in \mathbb{Z}[\alpha_1, \ldots, \alpha_8]$ are cubic polynomials and $M := \sum_{m_e \neq 0} \alpha_e$.

$$\tilde{I}_4(x) = \left[-\tfrac{1+x}{x} L_{\omega_{-1}} - \tfrac{1-2x-x^2}{(1-x)x^3} L_{\omega_0\omega_{-1}} + \cdots \right] \zeta(3) + 3\tfrac{1+x}{2x^3} L_{\omega_{-1}\omega_0\omega_0\omega_1} + \cdots$$
$$\in L\left(\left\{-1, 0, \tfrac{1}{2}, 1\right\}\right) \otimes \mathrm{lin}_{\mathbb{Q}} \{1, \zeta(3)\}.$$

Non-trivial kinematics: 3 variables

Examples: Conformal four-point integrals

The same functions with $z\bar{z} = \frac{x_{12}^2 x_{34}^2}{x_{13}^2 x_{24}^2}$ and $(1-z)(1-\bar{z}) = \frac{x_{14}^2 x_{23}^2}{x_{13}^2 x_{24}^2}$ describe conformal four-point integrals [11]. Many of them are linearly reducible, e.g.

$$E_{12;34} = \quad = \frac{x_{23}^2 x_{24}^2}{\pi^6} \int_{\mathbb{R}^{12}} \frac{\mathrm{d}^4 x_5\, \mathrm{d}^4 x_6\, \mathrm{d}^4 x_7 \cdot x_{16}^2}{(x_{15}^2 x_{25}^2 x_{35}^2) x_{56}^2 (x_{26}^2 x_{36}^2 x_{46}^2) x_{67}^2 (x_{17}^2 x_{27}^2 x_{47}^2)}$$

$$= H_{12;34} = \frac{x_{34}^2}{\pi^6} \int_{\mathbb{R}^{12}} \frac{\mathrm{d}^4 x_5\, \mathrm{d}^4 x_6\, \mathrm{d}^4 x_7 \cdot x_{57}^2}{(x_{15}^2 x_{25}^2 x_{35}^2 x_{45}^2) x_{56}^2 (x_{36}^2 x_{46}^2) x_{67}^2 (x_{17}^2 x_{27}^2 x_{37}^2 x_{47}^2)}$$

can be integrated parametrically to reproduce results of [11]. The result is a rational function in x_{ij}^2 multiplied with a polylogarithm in the variables z and $\bar{z}$.

Non-trivial kinematics: 3 variables

Examples: Conformal four-point integrals

Without inverse (numerator) propagators, there are only three four-point integrals with 4 internal vertices ("loops") that are not linearly reducible:







Non-trivial kinematics: 3 variables

Examples: Conformal four-point integrals

Without inverse (numerator) propagators, there are only three four-point integrals with 4 internal vertices ("loops") that are not linearly reducible:







Linearly reducible examples:







Non-trivial kinematics: 3 variables

Examples: Conformal four-point integrals

graph	denominator	weight	additional symbol letters	terms
	$x_{34}^2 x_{13}^4 x_{24}^4$ $\cdot(z-\bar{z})$ $\cdot(z+\bar{z}-2)$	8	$z-\bar{z}$ $z\bar{z}-1$ $z+\bar{z}-1$	4235
	$x_{34}^2 x_{13}^4 x_{24}^4$ $\cdot(z-\bar{z})^2$	8	$\emptyset$	107
	$x_{13}^4 x_{24}^4$ $\cdot z\bar{z}(z-\bar{z})$	7	$z-\bar{z}$	146

Extending linear reducibility

Box with two masses vis-à-vis



The on-shell massive box $p_1^2 = p_2^2 = p_3^2 = p_4^2 = -m^2$ with $m_2 = m_4 = 0$ and $m_1 = m_3 = m \neq 0$ is not linearly reducible.

Extending linear reducibility

Box with two masses vis-à-vis



The on-shell massive box $p_1^2 = p_2^2 = p_3^2 = p_4^2 = -m^2$ with $m_2 = m_4 = 0$ and $m_1 = m_3 = m \neq 0$ is not linearly reducible. The graph polynomials are

$$\psi = \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 \quad \text{and} \quad \varphi = s\alpha_1\alpha_3 + t\alpha_2\alpha_4 + m^2\left(\alpha_1 + \alpha_3\right)^2$$

and φ is only linear in α_2 and α_4.

Extending linear reducibility

Box with two masses vis-à-vis



The on-shell massive box $p_1^2 = p_2^2 = p_3^2 = p_4^2 = -m^2$ with $m_2 = m_4 = 0$ and $m_1 = m_3 = m \neq 0$ is not linearly reducible. The graph polynomials are

$$\psi = \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 \quad \text{and} \quad \varphi = s\alpha_1\alpha_3 + t\alpha_2\alpha_4 + m^2\left(\alpha_1 + \alpha_3\right)^2$$

and φ is only linear in α_2 and α_4. Reducing (integrating) α_4 we obtain

$$S_{\{4\}} = \left\{\alpha_1 + \alpha_2 + \alpha_3, s\alpha_1\alpha_3 + m^2\left(\alpha_1 + \alpha_3\right)^2, R\right\} \quad \text{where the resultant}$$
$$R := [\psi, \varphi]_{\alpha_4} = s\alpha_1\alpha_3 + m^2\left(\alpha_1 + \alpha_3\right)^2 - t\alpha_2(\alpha_1 + \alpha_2 + \alpha_3)$$

is irreducible and quadratic in all remaining Schwinger parameters, therefore prohibiting any further integration.

Extending linear reducibility

Box with two masses vis-à-vis

To proceed we change variables according to

$$\frac{s}{m^2} = \frac{(1-x)^2}{x} \quad \frac{t}{m^2} = \frac{(1-y)^2}{y} \quad \alpha_2 = \widetilde{\alpha_2}\,(\alpha_1 + x\alpha_3) \quad \alpha_4 = \widetilde{\alpha_4}\,(x\alpha_1 + \alpha_3)\,.$$

On one hand we reparametrized the kinematics via x and y to rationalize roots that would otherwise appear in the result, while afterwards

$$\varphi = \frac{m^2}{x}(\alpha_1 + x\alpha_3)(x\alpha_1 + \alpha_3) + \frac{m^2}{y}(1-y)^2\alpha_2\alpha_4$$

suggests to introduce the variables $\widetilde{\alpha_2}$ and $\widetilde{\alpha_4}$ with the effect that

$$\varphi = \frac{m^2}{xy}(\alpha_1 + x\alpha_3)(x\alpha_1 + \alpha_3)\left[y + x(1-y)^2\widetilde{\alpha}_2\widetilde{\alpha}_4\right]$$

factors linearly in these new parameters. Now $\left[\psi, y + x(1-y)^2\widetilde{\alpha}_2\widetilde{\alpha}_4\right]_{\widetilde{\alpha}_4}$ is linear in α_1 and α_3 allowing for a further integration.

Extending linear reducibility

Box with two masses vis-à-vis

Indeed we now observe linear reducibility in the new variables and obtain

$$S_{\{\widetilde{\alpha}_2,\alpha_3,\widetilde{\alpha}_4\}} = \{x+1, x-1, y+1, y-1, xy+1, x+y\}$$

as the final set in the polynomial reduction. Together with $\{x, y\}$ it defines the alphabet of the symbol of the resulting function of x and y in agreement with [13].

Extending linear reducibility

K_4

After integrating α_2 we are stuck, again because the resultant $R := [\psi, \varphi]_2 \in S_{\{2\}}$ is irreducible and quadratic in all Schwinger parameters. But its discriminant

$$D_{\alpha_3}(R) = \alpha_1^2 (\alpha_1\alpha_6 + \alpha_6\alpha_4 + \alpha_4\alpha_5 + \alpha_6\alpha_5)^2$$
$$\times \left[\alpha_6^2 t^2 \alpha_4^2 - 2t\alpha_4\alpha_6 (-t\alpha_6 + s\alpha_4 - s\alpha_6)\alpha_5 + (s\alpha_6 + s\alpha_4 + t\alpha_6)^2 \alpha_5^2\right]$$

becomes a perfect square if we introduce a variable ξ and reparametrize

$$\alpha_5 = \frac{t\alpha_4\alpha_6 (s\alpha_6 + t\alpha_6 + \xi)}{\xi (s\alpha_6 + s\alpha_4 + t\alpha_6 + \xi)}.$$

Extending linear reducibility

K_4

After integrating α_2 we are stuck, again because the resultant $R := [\psi, \varphi]_2 \in S_{\{2\}}$ is irreducible and quadratic in all Schwinger parameters. But its discriminant

$$D_{\alpha_3}(R) = \alpha_1^2 (\alpha_1\alpha_6 + \alpha_6\alpha_4 + \alpha_4\alpha_5 + \alpha_6\alpha_5)^2$$
$$\times \left[\alpha_6^2 t^2 \alpha_4^2 - 2t\alpha_4\alpha_6 (-t\alpha_6 + s\alpha_4 - s\alpha_6)\alpha_5 + (s\alpha_6 + s\alpha_4 + t\alpha_6)^2 \alpha_5^2\right]$$

becomes a perfect square if we introduce a variable ξ and reparametrize

$$\alpha_5 = \frac{t\alpha_4\alpha_6 (s\alpha_6 + t\alpha_6 + \xi)}{\xi (s\alpha_6 + s\alpha_4 + t\alpha_6 + \xi)}.$$

After this transformation, R factorizes linearly in α_3 and polynomial reduction reveals linear reducibility along the sequence $\alpha_2, \alpha_3, \alpha_1, \alpha_4, \xi$ ($\alpha_6 = 1$) of integrations.

Extending linear reducibility

K_4

After integrating α_2 we are stuck, again because the resultant $R := [\psi, \varphi]_2 \in S_{\{2\}}$ is irreducible and quadratic in all Schwinger parameters. But its discriminant

$$D_{\alpha_3}(R) = \alpha_1^2 (\alpha_1\alpha_6 + \alpha_6\alpha_4 + \alpha_4\alpha_5 + \alpha_6\alpha_5)^2$$
$$\times \left[\alpha_6^2 t^2 \alpha_4^2 - 2t\alpha_4\alpha_6 (-t\alpha_6 + s\alpha_4 - s\alpha_6)\alpha_5 + (s\alpha_6 + s\alpha_4 + t\alpha_6)^2 \alpha_5^2\right]$$

becomes a perfect square if we introduce a variable ξ and reparametrize

$$\alpha_5 = \frac{t\alpha_4\alpha_6 (s\alpha_6 + t\alpha_6 + \xi)}{\xi (s\alpha_6 + s\alpha_4 + t\alpha_6 + \xi)}.$$

After this transformation, R factorizes linearly in α_3 and polynomial reduction reveals linear reducibility along the sequence $\alpha_2, \alpha_3, \alpha_1, \alpha_4, \xi$ ($\alpha_6 = 1$) of integrations.
The final set is $\{s + t\}$ and proves that to all orders we obtain harmonic polylogarithms of $\frac{t}{s}$ which we calculated explicitly up to order ε^2 and checked against [12].

Soft phase space integrals

Using Mellin-Barnes representations in an intermediate step, a technology developed in [3, 4] allowed to rewrite soft phase space integrals in terms of parametric integrals like

$$\begin{aligned}\mathcal{I}_{9,1}(\varepsilon) = &- \int_0^\infty \mathrm{d}t_1 \mathrm{d}t_2 \int_0^1 \mathrm{d}x_1 \mathrm{d}x_2 \mathrm{d}x_3 \; t_1^{2-4\varepsilon} (1+t_1)^{\varepsilon-1} t_2^{1-2\varepsilon} \\ &\times x_1^{-\varepsilon} (1-x_1)^{2-4\varepsilon} x_2^{1-3\varepsilon} (1-x_2)^{-\varepsilon} x_3^{-\varepsilon} (1+t_2 x_3)^{1-3\varepsilon} (1+t_2 x_2 x_3)^{\varepsilon} \\ &\times \Big(t_1 t_2^2 x_1 x_2 x_3 + t_2^2 x_2 x_3 + t_1 t_2 x_1 x_2 + t_1 t_2 x_3 + t_2 x_2 x_3 + t_2 + t_1 + 1 \Big)^{3\varepsilon-3}.\end{aligned}$$

These are linearly reducible and could be integrated along the sequence t_1, x_1, x_2, x_3, t_2 of variables and evaluate to MZV.

References

Bibliography I

J. Ablinger, J. Blümlein, A. De Freitas, A. Hasselhuhn, S. Klein, C. Schneider, and F. Wißbrock.
New Results on the 3-Loop Heavy Flavor Wilson Coefficients in Deep-Inelastic Scattering.
ArXiv e-prints, January 2013.
arXiv:1212.5950v2.

J. Ablinger, J. Blümlein, A. Hasselhuhn, S. Klein, C. Schneider, and F. Wißbrock.
Massive 3-loop ladder diagrams for quarkonic local operator matrix elements.
Nuclear Physics B, 864(1):52–84, November 2012.
URL: http://www.sciencedirect.com/science/article/pii/S0550321312003355, arXiv:1206.2252.

Bibliography II

C. Anastasiou, C. Duhr, F. Dulat, F. Herzog, and B. Mistlberger.
Real-virtual contributions to the inclusive Higgs cross-section at N^3LO.
JHEP, 2013(12):88, December 2013.
arXiv:1311.1425.

C. Anastasiou, C. Duhr, F. Dulat, and B. Mistlberger.
Soft triple-real radiation for Higgs production at N_3LO.
JHEP, 2013(7):1–78, jul 2013.
arXiv:1302.4379.

J. Blümlein, A. De Freitas, C. Raab, F. Wißbrock, J. Ablinger, A. Hasselhuhn, M. Round, C. Schneider, and A. von Manteuffel.
Recent Results on the 3-Loop Heavy Flavor Wilson Coefficients in Deep-Inelastic Scattering.
ArXiv e-prints, July 2013.
arXiv:1307.7548.

Bibliography III

C. Bogner and M. Lüders.
Multiple polylogarithms and linearly reducible Feynman graphs.
ArXiv e-prints, February 2013.
arXiv:1302.6215.

F. C. S. Brown.
Multiple zeta values and periods of moduli spaces $\overline{\mathfrak{M}}_{0,n}(\mathbb{R})$.
Annales scientifiques de l'École Normale Supérieure, 42(3):371–489, June 2009.
arXiv:math/0606419.

F. C. S. Brown.
On the periods of some Feynman integrals.
ArXiv e-prints, October 2009.
arXiv:0910.0114.

Bibliography IV

F. C. S. Brown.
The Massless Higher-Loop Two-Point Function.
Communications in Mathematical Physics, 287:925–958, May 2009.
arXiv:0804.1660.

F. Chavez and C. Duhr.
Three-mass triangle integrals and single-valued polylogarithms.
Journal of High Energy Physics, 11:114, November 2012.
arXiv:1209.2722.

J. Drummond, C. Duhr, B. Eden, P. Heslop, J. Pennington, and V. A. Smirnov.
Leading singularities and off-shell conformal integrals.
Journal of High Energy Physics, 8:133, August 2013.
arXiv:1303.6909.

Bibliography V

J. M. Henn, A. V. Smirnov, and V. A. Smirnov.
Evaluating single-scale and/or non-planar diagrams by differential equations.
ArXiv e-prints, December 2013.
arXiv:1312.2588.

J. M. Henn and V. A. Smirnov.
Analytic results for two-loop master integrals for Bhabha scattering I.
JHEP, 1311:041, 2013.
arXiv:1307.4083.

E. Panzer.
On the analytic computation of massless propagators in dimensional regularization.
Nuclear Physics B, 874(2):567–593, September 2013.
URL: http://www.sciencedirect.com/science/article/pii/S0550321313003003, arXiv:1305.2161.

Bibliography VI

E. Panzer.
Algorithms for the symbolic integration of hyperlogarithms with applications to Feynman integrals.
ArXiv e-prints, March 2014.
arXiv:1403.3385.

E. Panzer.
On hyperlogarithms and Feynman integrals with divergences and many scales.
ArXiv e-prints, January 2014.
arXiv:1401.4361.

O. Schnetz.
Graphical functions and single-valued multiple polylogarithms.
ArXiv e-prints, February 2013.
arXiv:1302.6445.