20090701

Counting rational points on a certain exponential-algebraic surface

Jonathan Pila

1. Introduction

This paper is devoted to giving an upper estimate for the number of *non-trivial* rational points (or algebraic points over a given real numberfield) up to a given height on the surface $X \subset \mathbb{R}^3$ defined by

$$X = \{(x, y, z) \in (0, \infty)^3 : \log x \log y = \log z\}.$$

The half-lines $L_x = \{(x, 1, 1) : x > 0\}$ and $L_y = \{(1, y, 1) : y > 0\}$ contained in X evidently contain rational (or algebraic) points $(r, 1, 1), (1, s, 1) \in X$, where $r, s \in \mathbb{Q}_{>0}$ (or $r, s \in \overline{\mathbb{Q}} \cap \mathbb{R}_{>0}$), and these algebraic points we call *trivial*. Schanuel's conjecture implies (as we elaborate in §4) that there are no non-trivial algebraic points on X, and hence that there are no rational points on $X^0 = X - (L_x \cup L_y)$. Our result is that this conjecturally empty set is fairly sparse.

For a set $Y \subset \mathbb{R}^n$ put $Y(\mathbb{Q}) = Y \cap \mathbb{Q}^n$ and define, for $T \geq e$ (which we assume throughout),

$$Y(\mathbb{Q}, T) = \{x = (x_1, \ldots, x_n) \in Y \cap \mathbb{Q}^n : H(x_1), \ldots, H(x_n) \leq T\}$$

where $H(a/b) = \max(|a|, |b|)$ for a rational number a/b in lowest terms. The cardinality of a set A will be denoted $\#A$. Note that $\#(L_x \cup L_y)(\mathbb{Q}, T) \geq cT^2$, where c is some positive constant. In the sequel, $c(\alpha, \beta, \ldots), C(\alpha, \beta, \ldots)$ denote positive constants that depend only on $\alpha, \beta, \ldots$, and that may differ at each occurrence.

1.1. Theorem. *Let $\epsilon > 0$. Then*

$$\#X^0(\mathbb{Q}, T) \leq c(\epsilon)\Big(\log T\Big)^{44+\epsilon}.$$

This result may be viewed as a statement about the set of points $(x, y) \in (0, \infty)^2$ at which the three algebraically independent real-analytic functions $x, y, \exp(\log x \log y)$ are simultaneously rational, or alternatively about the points $(u, v) \in \mathbb{R}^2$ at which the functions e^u, e^v, e^{uv} are simultaneously rational. The set of points at which algebraically independent meromorphic functions of several complex variables simultaneously assume values in a number field has been quite extensively studied in connection with transcendental number theory, especially functions generating rings closed under partial differentiation [8, 1]. Without such assumptions, results of Lang [9], systematizing methods going back to Schneider, have been improved and extended by Waldschmidt [17] and others (see e.g. [20, 19]), and are intimately connected to interpolation problems and Schwarz Lemmas in several variables, see e.g. papers of Roy [16]. See also [18]. Note that we do not assume any hypotheses on the points (u, v), such as lying in a Cartesian product, nor is the ring of functions $\mathbb{C}[e^u, e^v, e^{uv}]$ closed under partial differentiation, while the function $\exp(\log x \log y)$ is not meromorphic in $\mathbb{C}^2$. Nevertheless, complex variable methods may well yield results along the lines of 1.1, although I am not aware of any explicit statements in the literature that imply such a result. We will employ real variable methods and draw on the theory of *o-minimal structures*.

To contextualise our result, we review the background results and conjectures. An *o-minimal structure* over $\mathbb{R}$ is, informally speaking, a sequence $\mathcal{S} = (\mathcal{S}_n), n = 1, 2, \ldots$with each $\mathcal{S}_n$ a collection of subsets of $\mathbb{R}^n$ such that $\cup_n \mathcal{S}_n$ contains all semi-algebraic sets and is closed under certain operations (boolean operations, products and projections), but nevertheless has strong finiteness properties (the boundary of every set in $\mathcal{S}_1$ is finite). A formal definition is given in the Appendix (§7), or see [5]. If $\mathcal{S}$ is an o-minimal structure over $\mathbb{R}$, a set $Y \subset \mathbb{R}^n$ belonging to $\mathcal{S}_n$ is said to be *definable* in $\mathcal{S}$. A set $Y \subset \mathbb{R}^n$ will be called *definable* if it is definable in some o-minimal structure over $\mathbb{R}$.

The paradigm example of an o-minimal structure is the collection of semi-algebraic sets. Another example is provided by the collection $\mathbb{R}_{\mathrm{an}}$ of *globally subanalytic* sets (see [6]), and the crucial example for this paper is the collection $\mathbb{R}_{\exp}$ of sets definable using the exponential function (see §7). The o-minimality of $\mathbb{R}_{\exp}$ is due to Wilkie [21], whose result yields the elegant description of the sets definable in $\mathbb{R}_{\exp}$ given in 7.2. The set X is definable in $\mathbb{R}_{\exp}$ (see 7.3).

Suppose then that $Y \subset \mathbb{R}^n$ is definable, and consider the counting function $\#Y(\mathbb{Q}, T)$. If Y contains semialgebraic sets of positive dimension (such as rational curves, as is the case for the set X), then one can certainly have

$$\#Y(\mathbb{Q}, T) \geq c(Y)T^{\delta}$$

for some positive δ. If on the other hand Y contains no semialgebraic sets of positive dimension then, according to [15] , one has

$$\#Y(\mathbb{Q}, T) \leq c(Y, \epsilon)T^{\epsilon}$$

for every $\epsilon > 0$. Indeed if we define, for any $Y \subset \mathbb{R}^n$, the *algebraic part* Y^{alg} of Y to be the union of all connected semialgebraic subsets of Y of positive dimension, then an estimate as above holds for the rational points of the *transcendental part* $Y^{\mathrm{trans}} = Y - Y^{\mathrm{alg}}$ of any definable set Y.

1.2. Theorem. ([15]) *Let Y be definable in an o-minimal structure over $\mathbb{R}$ and $\epsilon > 0$. Then*

$$\#Y^{\mathrm{trans}}(\mathbb{Q}, T) \leq c(Y, \epsilon)T^{\epsilon}.$$

Examples show (see [10, 7.5 and 7.6], elaborating a remark from [3]) that this estimate cannot be much improved in general. For example one can construct sets definable in $\mathbb{R}_{\mathrm{an}}$ such that no estimate of the form

$$\#Y^{\mathrm{trans}}(\mathbb{Q}, T) \leq c(Y)(\log T)^{C(Y)}$$

holds. However, Wilkie conjectured in [15] that such an estimate always holds for a set definable in $\mathbb{R}_{\exp}$.

1.3. Conjecture. Suppose Y is definable in $\mathbb{R}_{\exp}$. Then

$$\#Y^{\mathrm{trans}}(\mathbb{Q}, T) \leq c(Y)(\log T)^{C(Y)}.$$

Thus Theorem 1.1 affirms this conjecture for the particular set X. In fact X^{alg} consists of L_x and L_y together with infinitely many other rational curves defined over $\mathbb{R}$ (see 4.1). However these other rational curves do not contain any algebraic points (see 4.3).

Consider now the question of estimating the number of points of a definable set Y up to a given height defined over a real numberfield. Set $Y(F) = Y \cap F^n$ for a field $F \subset \mathbb{R}$ and put (again for $T \geq e$),

$$Y(F, T) = \{(x_1, \ldots, x_n) \in Y(F) : H(x_1), \ldots, H(x_n) \leq T\}.$$

where $H(x)$ is the absolute multiplicative height of an algebraic number, as defined in [2], which agrees with the previous definition of $H(x)$ for rational x. Theorem 1.2 may be extended quite straightforwardly to an estimate of the same form for $\#Y^{\mathrm{trans}}(F, T)$ when F is a numberfield (i.e. $[F : \mathbb{Q}] < \infty$), in which the implicit constant depends on Y, ϵ, and $[F : \mathbb{Q}]$.

Less straightforwardly, a much stronger result holds. For an integer $k \geq 1$, denote by

$$Y(k) = \{(x_1, \ldots, x_n) \in Y : [\mathbb{Q}(x_1) : \mathbb{Q}], \ldots, [\mathbb{Q}(x_n) : \mathbb{Q}] \leq k\}$$

the set of algebraic points of Y of degree $\leq k$. Observe that the definition permits the coordinates of a point in $Y(k)$ to be defined over different fields. Put (for $T \geq e$)

$$Y(k, T) = \{(x_1, \ldots, x_n) \in Y(k) : H(x_1), \ldots, H(x_n) \leq T\}.$$

Then for a definable set $Y \subset \mathbb{R}^n, k \geq 1$, and $\epsilon > 0$ we have ([14])

$$\#Y^{\text{trans}}(k,T) \leq c(Y,k,\epsilon)T^{\epsilon}.$$

To obtain this result one studies the rational points of a suitable definable set Y_k of higher dimension than Y whose rational points correspond to points of Y of degree $\leq k$. However Y_k^{trans} is empty, and a closer study of the proof structure of 1.2 is required.

In view of the above results for $Y(F,T)$ and $Y(k,T)$, it seems likely that if Conjecture 1.3 is affirmed, then the following stronger versions will also be affirmed. First, a version for varying number field with exponent independent of the number field.

1.4. Conjecture. Let $Y \subset \mathbb{R}^n$ be definable in $\mathbb{R}_{\text{exp}}$ and $F \subset \mathbb{R}$ a numberfield of degree $f = [F : \mathbb{Q}] < \infty$. Then

$$\#Y^{\text{trans}}(F,T) \leq c(Y,f)\Big(\log T\Big)^{C(Y)}.$$

Second, a version for algebraic points of bounded degree.

1.5. Conjecture. Let $Y \subset \mathbb{R}^n$ be definable in $\mathbb{R}_{\text{exp}}$ and $k \geq 1$. Then

$$\#Y^{\text{trans}}(k,T) \leq c(Y,k)\Big(\log T\Big)^{C(Y,k)}.$$

The following theorem affirms 1.4 for X. For the time being I cannot establish 1.5 for X. However I frame in Section 3 a conjecture (3.4) that would imply 1.4 and 1.5 in general.

1.6. Theorem. *Let $F \subset \mathbb{R}$ be a numberfield of degree f over $\mathbb{Q}$, and let $\epsilon > 0$. Then*

$$\#X^{\text{trans}}(F,T) \leq c(f,\epsilon)\Big(\log T\Big)^{44+\epsilon}.$$

That the exponent of $\log T$ in 1.6 is independent of F is a feature related to transcendence theory. In [13] I affirmed Wilkie's conjecture for *pfaff curves* (see 5.2). (This class of plane curves does not contain all plane curves definable in $\mathbb{R}_{\text{exp}}$, but on the other hand there are pfaff curves that are not definable in $\mathbb{R}_{\text{exp}}$.) In [14] I observed that the result held for the points of a pfaff curve defined over a real number field F, and with an exponent of $\log T$ independent of F. This result applies in particular to the graph $W_\alpha : y = x^\alpha, x \in (0,\infty)$, for positive irrational α, though it gives a result weaker than previously known results in that case. According to [13] and (for algebraic points) [14], if $F \subset \mathbb{R}$ is a numberfield with $[F : \mathbb{Q}] = f$ then

$$\#W_\alpha(F,T) \leq C(f)\Big(\log T\Big)^{20}.$$

This estimate directly implies a weak form of the "Six Exponentials Theorem" as follows. Suppose there were 21 algebraic points (x_i, y_i) on W_α with the x_i multiplicatively independent. Then, considering the points $(\Pi x_i^{a_i}, \Pi y_i^{a_i})$ for 21-tuples of integers a_i, we would have $\#W_\alpha(F,T) \geq c(W_\alpha,F)\Big(\log T\Big)^{21}$ for suitable F, giving a contradiction. Therefore, we conclude that if w_i are 21 real numbers, linearly independent over $\mathbb{Q}$, then at least one of the 42 exponentials $\exp w_i, \exp(\alpha w_i)$ must be transcendental.

In fact the same conclusion holds if there are just 3 linearly independent w_i, namely that at least one of the *six* exponentials $\exp w_i, \exp(\alpha w_i)$ is transcendental. This is the Six Exponentials Theorem, and our "Forty-Two Exponentials Theorem" is rather weak. However the point I wish to observe is that any estimate $\#W_\alpha(F,T) \leq c(W_\alpha,F)\Big(\log T\Big)^{C(W_\alpha)}$ with $C(W_\alpha)$ independent of F entails a transcendence result because the curve W_α is a group (with suitable height growth in the group law), so that finitely many independent points generate an infinite set of a certain log-power density. The surface X is not a group, and so our $\#X^{\text{trans}}(F,T) \leq c(f)\Big(\log T\Big)^{C}$ estimate does not yield a transcendence result, even though it is – qualitatively speaking – of the same quality.

Thus a uniform version of Wilkie's conjecture i.e. that $\#Y^{\mathrm{trans}}(F,T) \le c(Y,F)\Big(\log T\Big)^{C(Y)}$ for a set Y definable in $\mathbb{R}_{\exp}$ and a real numberfield F with an exponent $C(Y)$ indepenent of F (just as we affirm for X in 1.6) can be viewed as a qualitative transcendence-type statement, and for suitable sets Y it would indeed imply a transcendence result.

Our strategy combines elements of the approaches of several previous papers. The key to the method of [15] is the possibility of parameterizing a definable subset of $(0,1)^n$ of dimension k by finitely many functions $(0,1)^k \to (0,1)^n$ all of whose partial derivatives up to a prescribed order are bounded in absolute value by 1. In [12] I showed that Wilkie's conjecture holds for pfaff curves that are *mild*, i.e. admit a parameterization in which derivatives to *all* orders are suitably controlled (see §2). Later, I established Wilkie's conjecture in the form 1.3 for all pfaff curves by a different method in [13], and in the form 1.4 in [14]. Here, a mild parameterization of X is used to show that $X(F,T)$ is contained in $<< (\log T)^C$ intersections of X with hypersurfaces of degree $<< (\log T)^2$. These intersection curves are treated by adapting the methods of [13]. Here, as in [12, 13], a crucial role is played by results of Gabrielov and Vorobjov [7] estimating the topological complexity of *Pfaffian sets* (see §5). As it stands, this combination of methods — mild parameterization for the initial set and Pfaffian bounds for the intersection curves — is applicable only to surfaces. Our surface X was selected as being related to the threefold $\log x \log y = \log z \log t$ associated with the Four Exponentials Conjecture (see [17]). The present method is generalized by Butler [4] to further surfaces definable in $\mathbb{R}_{\exp}$.

Acknowledgements. My thanks to Lee Butler for detailed corrections to a previous version of this paper, to Eric Descheemaeker for assistance, and to the referee for helpful comments and suggestions. I am grateful to Roger Heath-Brown and the Mathematical Institute, Oxford, for affording me hospitality as an Academic Visitor, and to the Leverhulme Trust for supporting my work through a Research Fellowship.

2. Mild functions

We write $x = (x_1, \dots, x_k)$ etc as variables in $\mathbb{R}^k$. For a function $\phi : U \to \mathbb{R}$ defined on some domain $U \subset \mathbb{R}^k$ and $\mu = (\mu_1, \dots, \mu_k) \in \mathbb{N}^k$ we set $|\mu| = \sum \mu_i$ and denote by $\partial^\mu \phi$ the partial derivative

$$\partial^\mu \phi = \phi^{(\mu)} = \frac{\partial^{|\mu|}\phi}{\partial x_1^{\mu_1} \dots \partial x_k^{\mu_k}}$$

of order $|\mu|$. We denote by x^μ the monomial $\prod_i x_i^{\mu_i}$ of degree $|\mu|$. We set $\mu! = \prod_i \mu_i!$ and $\overline{\mu} = \max_i \mu_i$.

2.1. Definition. A function $\phi : (0,1)^k \to (0,1)$ is called (A,C)-*mild* if it is C^∞ and, for all $\mu \in \mathbb{N}^k$ and all $z \in (0,1)^k$,

$$|\partial^\mu \phi(z)| \le \mu!(A|\mu|^C)^{|\mu|}.$$

2.2. Remark. One could define a finer notion (A,B,C)-*mild* with a term $(|\mu|+1)^B$ to enable finer estimates. However only the parameter C survives to influence the exponent of $\log T$ in the density estimate, so the above notion was preferred for simplicity.

2.3. Definition. A function $\theta : (0,1)^k \to (0,1)^n, \theta(x) = (\theta_1(x), \dots, \theta_n(x))$ is called (A,C)-*mild* if each of its coordinate functions θ_i is (A,C)-mild.

2.4. Definition. A set $Y \subset (0,1)^n$ of dimension k is called (J,A,C)-*mild* if there exists a collection Θ of (A,C)-mild maps $\theta : (0,1)^k \to (0,1)^n$ such that $\#\Theta = J$ and

$$\bigcup_{\theta \in \Theta} \theta\Big((0,1)^k\Big) = Y.$$

A set $Y \subset (0,1)^n$ is called *mild* if it is (J,A,C)-mild for some J, A, C.

2.5. Conjecture. Every set $Y \subset (0,1)^n$ definable in $\mathbb{R}_{\exp}$ is mild.

A more precise version of this conjecture is formulated in 3.4. A more optimistic version would require a fixed value of C. The following property of mild functions will be used in the sequel.

2.6. Proposition. *Suppose $\phi_1, \ldots, \phi_\ell : (0,1)^k \to (0,1)$ are (A,C)-mild, $\mu \in \mathbb{N}^k$ and $z \in (0,1)^k$. Then*

$$|\partial^\mu \phi_1 \ldots \phi_\ell(z)| \le \mu!(\overline{\mu}+1)^{(\ell-1)k} \left(A|\mu|^C\right)^{|\mu|}.$$

Proof. We have

$$\partial^\mu \phi_1 \ldots \phi_\ell = \sum_{\mu_1+\ldots+\mu_\ell=\mu} \mathrm{Ch}(\mu_1, \ldots, \mu_\ell) \prod_{i=1}^{\ell} \partial^{\mu_i} \phi_i$$

where, for $\alpha = (\alpha_1, \ldots, \alpha_k), \beta = (\beta_1, \ldots, \beta_k)$, etc

$$\mathrm{Ch}(\alpha, \beta, \ldots, \zeta) = \prod_{j=1}^{k} \frac{(\alpha_j + \beta_j + \ldots + \zeta_j)!}{\alpha_j! \beta_j! \ldots \zeta_j!}.$$

Therefore

$$\frac{|\partial^\mu \phi_1 \ldots \phi_\ell(z)|}{\mu!} \le \sum_{\mu_1+\ldots+\mu_\ell=\mu} \prod_{i=1}^{k} \frac{|\partial^{\mu_i} \phi_i|}{\mu_i!} \le (\overline{\mu}+1)^{(\ell-1)k} \prod \left(A|\mu_i|^C\right)^{|\mu_i|} \le (\overline{\mu}+1)^{(\ell-1)k} \left(A|\mu_|^C\right)^{|\mu_|}$$

as required. □

We next establish that certain functions that we will use in our parameterizations are mild. First observe that the function

$$\psi(r) = r^r e^{1-r} = \exp\Big(r \log r + 1 - r\Big)$$

is increasing for $r \ge 1$, as the derivative $\log r$ of the exponent is positive for $r > 1$, and has $\psi(1) = 1$. We define $\psi(0) = 1$.

2.7. Lemma. *Let $m = (m_1, \ldots, m_k) \in (0,\infty)^k, a = (a_1, \ldots, a_k) \in [0,\infty)^k$ and suppose that, for each i, either $a_i = 0$ or $a_i \ge m_i$. Define $E_{m,a} : (0,1)^k \to \mathbb{R}$ by*

$$E_{m,a}(z) = \exp\left(1 - \frac{1}{z^m}\right) \frac{1}{z^a}.$$

Then

$$\sup_{z \in (0,1)^k} |E_{m,a}(z)| = \psi\Big(\max_i (a_i/m_i)\Big).$$

Proof. If all $a_j = 0$ then the supremum is clearly 1, which agrees with our definition of $\psi(0) = 1$. So we can assume that some $a_j > 0$, so that $a_j \ge m_j$ by our hypothesis, and then $\max_i(a_i/m_i) \ge a_j/m_j \ge 1$.

We proceed by induction on k. If $k = 1$ we have $E_{m,a}(z) = E(t) = \exp(1 - t^{-1})t^{-r}$ where $t = z^m$, $t \in (0,1)$, $r \ge 1$. The maximum of the function for $t \in [0,\infty)$ occurs at $t = 1/r \in (0,1]$ and has the value $\psi(r)$.

Suppose the result true for $k-1$ variables, $k \ge 2$. We have

$$\partial^{x_i} E_{m,a}(z) = \frac{E_{m,a}(z)}{z_i} \left(m_i z^{-m} - a_i\right).$$

If all $a_i/m_i = r$, the function again reduces to a function of one variable, $E_{m,a}(z) = \exp(1 - t^{-1})t^{-r}$, where $t = z^m, r \ge 1$, $t \in (0,1)$. As before the maximum of the function for $t \in [0,\infty)$ occurs at $t = 1/r$ and has the value $\psi(r)$, affirming the conclusion.

If the a_i/m_i are not all equal, then there is no stationary point inside $(0,1)^k$ and the supremum is given by the maximum of the function on $[0,1]^k$, which is attained on a boundary, and moreover on a boundary where some $x_i = 1$, as the function is flat at the $x_i = 0$ boundaries.

By induction, the supremum on a boundary $x_j = 1$ is $\psi(\max(r_j, j \neq i))$. As the function ψ is increasing for arguments ≥ 1, we get the desired conclusion in this case too, and complete the induction and the proof. □

2.8. Proposition. *For $m = (m_1, \ldots, m_k) \in (0,\infty)^k$ define $E_m : (0,1)^k \to \mathbb{R}$ by*

$$E_m(z) = \exp\left(1 - \frac{1}{z^m}\right).$$

Then E_m is (A,C)-mild with $C = \max\left((m_i+1)/m_i\right)$ and $A = (\overline{m}+1)C^C e^{-C}$.

Proof. Write $E = E_m$. For $\mu \in \mathbb{N}^k$ we have

$$\partial^\mu E = E \sum_{m'} a_{m'}^{(\mu)} z^{-m'}$$

over suitable $m' \in (0,\infty)^k$. The m' that appear all have, for each i, $m'_i = 0$ or $m'_i > m_i$. Furthermore, for each i, the largest m'_i occuring is $\mu_i(m_i+1)$.

Set, for $\mu \in \mathbb{N}^k$,

$$\alpha_\mu = \sum_{m'} |a_{m'}^{(\mu)}|$$

and, for $\ell \in \mathbb{N}$,

$$\alpha_\ell = \max_{|\mu|=\ell} \alpha_\mu.$$

Denote by e_i the element of $\mathbb{N}^k$ that has zero entries except for an entry 1 in the i-th place, so that $\partial^{e_i} = \partial^{z_i}$. Observe that

$$\partial^{e_i}\partial^\mu E = \partial^{e_i}\left(E \sum_{m'} a_{m'}^{(\mu)} z^{-m'}\right) = E \sum_{m'} a_{m'}^{(\mu)} \left(z^{-m'} \frac{m_i}{z^{m+e_i}} - \frac{m'_i}{z^{m'+e_i}}\right).$$

Therefore

$$\alpha_{\mu+e_i} \leq m_i\alpha_\mu + \max_{m'}(m'_i)\alpha_\mu = m_i\alpha_\mu + \mu_i(m_i+1)\alpha_\mu \leq (\mu_i+1)(\overline{m}+1)\alpha_\mu,$$

and so, by induction on $|\mu|$,

$$\alpha_\mu \leq \mu!(\overline{m}+1)^{|\mu|}.$$

The largest "a/m" occuring is

$$\max_i \frac{\mu_i(m_i+1)}{m_i} \leq \overline{\mu}\lambda$$

where

$$\lambda = \max_i \frac{m_i+1}{m_i}.$$

By Lemma 2.7,

$$\frac{|\partial^\mu E(z)|}{\mu!} \leq (\overline{m}+1)^{|\mu|}\left(\frac{\overline{\mu}\lambda}{e}\right)^{\overline{\mu}\lambda}.$$

This establishes that E_m is (A,C)-mild with

$$A = (\overline{m}+1)\left(\frac{\lambda}{e}\right)^\lambda, \quad C = \lambda$$

as required. □

3. Exploring mild sets with algebraic hypersurfaces

3.1. Proposition. *For integers $a > 0, x \geq a(a+1)/2$,*

$$\frac{x^a}{a!} \leq \binom{a+x}{a} \leq \frac{x^a}{a!}\left(1 + \frac{a(a+1)}{x}\right).$$

Proof. We have

$$\binom{a+x}{a} = \frac{(a+x)!}{a!x!} = \frac{x^a}{a!}(1+\frac{a}{x})(1+\frac{a-1}{x})\dots(1+\frac{1}{x}).$$

So the left-hand inequality of the Proposition is immediate provided only a, x are positive, while

$$\log\left((1+\frac{a}{x})(1+\frac{a-1}{x})\dots(1+\frac{1}{x})\right) \leq \frac{a}{x} + \dots + \frac{1}{x} = \frac{a(a+1)}{x}.$$

Since $e^y \leq 1 + 2y$ for $0 \leq y \leq 1$, the assumption $x \geq a(a+1)/2$ implies

$$(1+\frac{a}{x})(1+\frac{a-1}{x})\dots(1+\frac{1}{x}) \leq \exp\left(\frac{a(a+1)}{2x}\right) \leq 1 + \frac{a(a+1)}{x}$$

giving the right-hand inequality provided $x \geq a(a+1)/2$. □

We observe the following consequences of this Lemma, in which the expression "$1+o(1)$" is to apply for $d \to \infty$ with k, n fixed.

Let $\Lambda_k(d)$ denote the set of monomials of exact degree d in k variables, and $L_k(d) = \#\Lambda_k(d)$. Then

$$L_k(d) = \binom{k-1+d}{k-1} = \frac{d^{k-1}}{(k-1)!}(1+o(1)).$$

Let $\Delta_k(d)$ denote the set of monomials of degree $\leq d$ in k variables, and $D_k(d) = \#\Delta_k(d)$. Then

$$D_k(d) = L_{k+1}(d) = \frac{d^k}{k!}(1+o(1)).$$

Let $b(k,n,d)$ be the unique positive integer b with

$$D_k(b) \leq D_n(d) < D_k(b+1).$$

Then

$$b(k,n,d) = \left(\frac{k!d^n}{n!}\right)^{1/k}(1+o(1)).$$

Let

$$B(k,n,d) = \sum_{\beta+0}^{b} L_k(\beta)\beta + \left(D_n(d) - \sum_{\beta=0}^{b} L_k(\beta)\right)(b+1).$$

Then

$$B(k,n,d) = \frac{1}{(k+1)!(k-1)!}\left(\frac{k!}{n!}\right)^{(k+1)/k} d^{n(k+1)/k}(1+o(1)).$$

Finally, let

$$V(n,d) = \sum_{\beta=0}^{d} L_n(\beta)\beta.$$

Then

$$V(n,d) = \frac{1}{(n+1)(n-1)!}d^{n+1}(1+o(1)).$$

The following are the results showing that, for a mild set $Y \subset (0,1)^n$ of dimension k, $Y(F,T)$ is contained in "few" algebraic hypersurfaces. It is convenient to establish the result first using a different height function.

For an algebraic number α we denote by $\mathrm{den}(\alpha)$ the *denominator* of α, namely, the least positive integer m such that $m\alpha$ is an algebraic integer. If $\alpha_i \in \mathbb{C}$ are the conjugates of α we set

$$H^{\mathrm{size}}(\alpha) = \max\{\mathrm{den}(\alpha), |\alpha_i|\}.$$

Suppose α, of degree f, with minimal polynomial (over $\mathbb{Z}$) $a_f(t-\alpha_1)\dots(t-\alpha_f)$. Then [2, 1.6.5, 1.6.6]

$$H^{\mathrm{size}}(\alpha) \le |a_f| \prod \max(1, |\alpha_i|) = H(\alpha)^f.$$

For $Y \subset \mathbb{R}^n$ we set

$$Y^{\mathrm{size}}(F,T) = \{(x_1,\dots,x_n) \in Y(F) : H^{\mathrm{size}}(x_1),\dots,H^{\mathrm{size}}(x_n) \le T\}.$$

For $\alpha \in \mathbb{R}$ we let $[\alpha]$ denote the integer part (least integer not exceeding α).

3.2. Theorem. *Suppose $Y \subset (0,1)^n$ of dimension k has a (J,A,C)-mild parameterization. Let f be a positive integer and $F \subset \mathbb{R}$ a numberfield of degree f over $\mathbb{Q}$. Then $Y^{\mathrm{size}}(F,T)$ is contained in at most*

$$Jc(k,n)^f A^{(k+1)(1+o(1))} \Big(\log T\Big)^{C\left(\frac{n(k+1)}{n-k}\right)(1+o(1))}$$

intersections of Y with hypersurfaces (possibly reducible) of degree

$$d = \left[\Big(\log T\Big)^{\frac{k}{n-k}}\right]$$

where "$1+o(1)$" is taken as $T \to \infty$ with implicit constants depending only on k,n.

Proof. Since Y is the union of J images of mild maps, it suffices (given the factor J in the conclusion) to suppose that Y is the image of a single (A,C)-mild map $\theta : (0,1)^k \to (0,1)^n$.

Consider a $D_n(d) \times D_n(d)$ determinant Δ of the form

$$\Delta = \det\Big((x^{(i)})^j\Big)$$

where $j \in \mathbb{N}^n$ with $|j| \le d$ indexes the columns, $x^{(i)} \in Y(F,T)$, $i = 1,\dots,D_n(d)$, and x^j denotes as usual the monomial $\prod_\ell x_\ell^{j_\ell}$.

Each coordinate of each $x^{(i)}$ has denominator $\le T$. The entries in row i consist of monomials in which each coordinate is raised to power $\le d$. Therefore $K\Delta$ is an algebraic integer for some positive integer K with

$$K \le T^{ndD_n(d)},$$

and then

$$\prod_\sigma \Big(K\Delta\Big)^\sigma \in \mathbb{Z}$$

where σ runs over the embeddings $F \to \mathbb{C}$.

Let us estimate $|\Delta^\sigma|$ (later we will use the mild parameterization to get a better estimate for Δ itself, i.e. when $\sigma = \mathrm{id}$). Expand Δ^σ into a sum of $D_n(d)!$ terms. Since Δ has $L_n(\beta)$ columns of degree β, for $\beta = 0,\dots,d$, and in each column the entries have absolute value at most T^β, the largest term in the expansion has complex absolute value

$$\le T^{\sum L_n(\beta)\beta} = T^{V(n,d)}$$

so that, for any σ,

$$|\Delta^\sigma| \le D_n(d)! T^{V(n,d)}.$$

Therefore, if $\Delta \neq 0$ then $\prod_\sigma (K\Delta)^\sigma$ is a non-zero integer and

$$1 \le |K\Delta| \prod_{\sigma \neq \mathrm{id}} |K\Delta^\sigma| \le |\Delta| T^{fndD_n(d)+(f-1)V(n,d)} \Big(D_n(d)!\Big)^{f-1}.$$

To estimate $|\Delta|$, suppose that the points $x^{(i)}$ are the images of some points $z^{(i)} \in (0,1)^k$ under θ where the $z^{(i)}$ in fact belong to some cube of side $\le r \le 1$, and so are at a distance $\le r$ in each coordinate from the centre $z^{(0)}$ of the cube, which contains also all the lines segments from $z^{(0)}$ to $z^{(i)}$. We have then that

$$\Delta = \det\Big(\phi_j(z^{(i)})\Big)$$

where ϕ_j is the monomial function indexed by j, namely

$$\phi_j(z^{(i)}) = \Big(\theta_1(z^{(i)}), \dots, \theta_n(z^{(i)})\Big)^j = \Big(x_1^{(i)}, \dots, x_n^{(i)}\Big)^j.$$

We expand each entry of Δ as a Taylor series about $z^{(0)}$ of order $b = b(k,n,d)$ with remainder terms of order $b+1$:

$$\phi_j(z^{(i)}) = \sum_{\mu \in \Delta_k(b)} \frac{\partial^\mu \phi_j(z^{(0)})}{\mu!} \Big(z^{(i)} - z^{(0)}\Big)^\mu + \sum_{\mu \in \Lambda_k(b+1)} \frac{\partial^\mu \phi_j(\zeta)}{\mu!} \Big(z^{(i)} - z^{(0)}\Big)^\mu$$

where $\zeta = \zeta_{ij}$ is a suitable intermediate point on the line segment from $z^{(0)}$ to $z^{(i)}$.

Now we expand out the determinant. In doing so, terms of low degree as products of terms of the form $(z_\ell^{(i)} - z_\ell^{(0)})$ cancel out, as observed in [10, Proof of 3.1]. Specifically, consider the totality of terms corresponding to a particular specification of the number of multiplicands of each order of derivative. Consider a minor of size $h \times h$ of $\det(\phi_j(z^{(i)})$ comprising the expansion terms of degree $\beta \le b$ only. That is, select h points $\zeta^{(i)}$ from among the $z^{(i)}$, and h functions ψ_j from among the ϕ_j and consider

$$\det\left(\sum_{\mu \in \Lambda_k(\beta)} \frac{\partial^\mu \psi_j(z^{(0)}}{\mu!} \Big(\zeta^{(i)} - z^{(0)}\Big)^\mu\right).$$

If $h > L_k(\beta)$ then the columns are dependent and the minor vanishes. Thus if, for a particular specification of orders, there are more than $L_k(\beta)$ multiplicands of order β for some β, then the totality of terms corresponding to this choice vanishes.

Therefore, all surviving terms are products of $B(k,n,d)$ or more terms of the form $(z_\ell^{(i)} - z_\ell^{(0)})$. The number of surviving terms is estimated by the number of terms assuming no cancellation, i.e. for each term we consider which row the multiplicand from column j came from, for which there are $D_n(d)!$ possibilities, and given this choice we can then choose, for each column, one of the $D_k(b+1)$ terms in the Taylor expansion, giving an estimate for the number of terms of at most

$$D_n(d)! D_k(b+1)^{D_n(d)}.$$

Finally, each term is a product of $D_n(d)$ terms, each one of the summands in the Taylor formula for ϕ_j which, neglecting the terms $(z_\ell^{(i)} - z_\ell^{(0)})$, takes the form

$$\frac{\partial^\mu \left(\theta^j\right)(\zeta)}{\mu!}$$

for some suitable ζ, and some μ with $|\mu| \le b+1$. By Proposition 2.6, as θ is (A,C)-mild and $|\mu| \le b+1$,

$$\frac{|\partial^\mu \left(\theta^j\right)(\zeta)|}{\mu!} \le (\overline{\mu}+1)^{(|j|-1)k} \Big(A(b+1)^C\Big)^{b+1} \le (b+2)^{|j|k} \Big(A(b+1)^C\Big)^{b+1}.$$

Now

$$\sum_{j\in\mathbb{N}^n:|j|\le d} |j| = \sum_{\beta=0}^{d} \beta L_n(\beta) = V(n,d)$$

so that

$$\prod_{j\in\mathbb{N}^n:|j|\le d} (b+2)^{|j|k} \le (b+2)^{kV(n,d)}.$$

Therefore, since $|z_\ell^{(i)} - z_\ell^{(0)}| \le r \le 1$,

$$|\Delta| \le D_n(d)! D_k (b+1)^{D_n(d)} (b+2)^{kV(n,d)} \left(\left(A(b+1)^C\right)^{b+1}\right)^{D_n(d)} r^{B(k,n,d)},$$

and if the points $x^{(i)}$ **do not** lie on any hypersurface in $\mathbb{R}^n$ of degree d then $\Delta \neq 0$ and

$$1 \le (D_n(d)!)^f\, D_k (b+1)^{D_n(d)} (b+2)^{kV(n,d)} T^{fndD_n(d)+fV(n,d)} \left(\left(A(b+1)^C\right)^{b+1}\right)^{D_n(d)} r^{B(k,n,d)}.$$

Now we take the $B(k,n,d)$-th root of this inequality. In the following discussion, the expression "$1+o(1)$" is to be taken as $d\to\infty$ with k,n fixed, while $c(k,n)$ is a positive constant that may differ at each occurence.

First we observe that

$$\frac{D_n(d)}{B(k,n,d)} = \frac{d^n}{n!}\frac{(k+1)(k-1)!}{d^{n(k+1)/k}}\left(\frac{n!}{k!}\right)^{\frac{k+1}{k}}(1+o(1)) = \frac{c(k,n)}{d^{n/k}}$$

where

$$c(k,n) = \frac{k+1}{k}\left(\frac{n!}{k!}\right)^{1/k}(1+o(1)),$$

and that

$$\frac{V(n,d)}{B(k,n,d)} = c(k,n)(1+o(1))\frac{d^{n+1}}{d^{n(k+1)/k}} = \frac{c(k,n)}{d^{n/k-1}}.$$

So

$$\left(D_n(d)!\right)^{f/B(k,n,d)} \le D_n(d)^{\frac{fD_n(d)}{B(k,n,d)}} = \left(c(k,n)(1+o(1)d^n)\right)^{fc(k,n)/d^{n/k}} = \left(1+o(1)\right)^f,$$

and similarly

$$D_k(b+1)^{\frac{D_n(d)}{B(k,n,d)}} = \left(\frac{(b+1)^k}{k!}(1+o(1))\right)^{\frac{c(k,n)(1+o(1))}{d^{n/k}}} = \left(c(k,n)(1+o(1))d^n)\right)^{\frac{c(k,n)(1+o(1))}{d^{n/k}}} = 1+o(1).$$

Next,

$$(b+2)^{\frac{kV(n,d)}{B(k,n,d)}} = \left(c(k,n)(1+o(1))d^{n/k}\right)^{\frac{c(k,n)}{d^{n/k-1}}} = 1+o(1).$$

We have

$$T^{\frac{fnD_n(d)+fV(n,d)}{B(k,n,d)}} = c(k,n)^f$$

provided

$$d = \left[\left(\log T\right)^{\frac{k}{n-k}}\right].$$

Finally

$$\frac{(b+1)D_n(d)}{B(k,n,d)} = \frac{k+1}{k}(1+o(1)),$$

so that

$$\left(A(b+1)^C\right)^{\frac{(b+1)D_n(d)}{B(k,n,d)}} = \left(Ac(k,n)(1+o(1))d^{Cn/k}\right)^{\frac{k+1}{k}(1+o(1))} = c(k,n)A^P d^{nCP/k}.$$

where

$$P = \frac{k+1}{k}(1+o(1)).$$

Thus if $\Delta \neq 0$ we find that

$$1 \leq c(k,n)^f A^P d^{nCP/k} r$$

where

$$d = \left[\left(\log T\right)^{\frac{k}{n-k}}\right]$$

and all the preimages $z^{(i)}$ of the points $x^{(i)}$ lie in a cube of side r in $(0,1)^k$. The points $x^{(i)}$ whose coordinates have $H^{\text{size}}(x_j^{(i)}) \leq T$ and whose preimages lie in such a cube must therefore all lie on **one** hypersurface (possibly reducible) of degree d provided

$$r < c(k,n)^f A^{-P} d^{-nCP/k},$$

and since $(0,1)^k$ may be covered by at most

$$c(k,n)^f A^{kP} d^{nCP}$$

such cubes, and T, d go to infinity together, the proof is complete. □

3.3. Corollary. *Suppose $Y \subset (0,1)^n$ of dimension k has a (J,A,C)-mild parameterization. Let f be a positive integer and $F \subset \mathbb{R}$ a numberfield of degree f over $\mathbb{Q}$. Then $Y(F,T)$ is contained in at most*

$$Jc(k,n)^f A^{(k+1)(1+o(1))} \left(f \log T\right)^{C\left(\frac{n(k+1)}{n-k}\right)(1+o(1))}$$

intersections of Y with hypersurfaces (possibly reducible) of degree

$$d = \left[\left(f \log T\right)^{\frac{k}{n-k}}\right]$$

where "$1+o(1)$" is taken as $T \to \infty$ with implicit constants depending only on k, n.

Proof. We have $Y(F,T)$ contained in $Y^{\text{size}}(F, T^f)$. □

3.4. Conjecture. Let $Y \subset (0,1)^n$ be definable in $\mathbb{R}_{\exp}$. There exist constants C_1, C_2, C_3, C_4, C_5 depending only on Y with the following property. Let $\mathcal{F}$ be an algebraic family of closed algebraic sets in $\mathbb{R}^n$ of degree $d = d(\mathcal{F})$, and suppose $V \in \mathcal{F}$. Then $Y \cap V$ is $(C_2 d^{C_3}, C_4 d^{C_5}, C_1)$-mild.

3.5. Corollary. *Conjecture 3.4 implies Conjectures 1.4 and 1.5.*

Proof. It suffices to work with H^{size}. By maps $x \to \pm x^{\pm 1}$ it suffices, as in [15], to consider sets $Y \subset (0,1)^n$. Then one iteratively intersects with hypersurfaces. Assuming 3.4, all the sets involved are (J,A,C_1)-mild with C_1 fixed and J, A depending polynomially on the degree of the family. For 1.5, imitate the proof of Theorem 5.3 in [14] using 3.3 to estimate the number of intersections required at each stage, rather than the appeal in [14] (via [15]) to [10, Lemma 4.4]. For 1.4, use 3.3 on Y and then on the intersections given by the conclusion of 3.3 repeatedly. In both cases the degrees of the families are polynomial in $(\log T)$ at each stage. □

4. The algebraic part, Schanuel's conjecture and algebraic points

4.1. Proposition. *Let*

$$X = \{(x,y,z) \in (0,\infty)^3 : \log x \log y = \log z\}$$

Then X^{alg} consists of the lines $L_x = \{(x,1,1) : x \in (0,\infty)\}$ and $L_y = \{(1,y,1) : y \in (0,\infty)\}$ and, for $q \in \mathbb{Q}^*$, the curves $\Gamma_{x,q} = \{(x, e^q, z) : z = x^q, x > 0\}$ and $\Gamma_{y,q} = \{(e^q, y, z) : z = y^q, y > 0\}$.

Proof. Suppose that Γ is an arc of an algebraic curve contained in X. Suppose x is constant on Γ. If $x = 1$ then also $z = 1$ and Γ is an arc of the line L_y. If x is constant but not equal to 1 then $q = \log x$ must be rational, and Γ is contained in the curve $\Gamma_{y,q}$. Similarly, if y is constant we find Γ contained in L_x or $\Gamma_{x,q}$. If z is constant, we get no algebraic curves unless $z = 1$ and we find that either $x = 1$ or $y = 1$ identically on Γ and revert to the previous cases. Otherwise, x, y, z are non-constant and further y, z are algebraic functions of x. We then have

$$x = \exp\left(\frac{\log z(x)}{\log y(x)}\right)$$

on Γ and, by analytic continuation, this relation holds also for large (possibly complex) x. Then $y(x), z(x)$ are given by some convergent Puiseaux series,

$$z(x) = z_0 x^{\zeta} + \ldots, \quad y(x) = y_0 x^{\eta} + \ldots$$

and we have

$$x = \exp\left(\frac{\zeta \log x + \log z_0 + \log(1 + \ldots)}{\eta \log x + \log y_0 + \log(1 + \ldots)}\right)$$

which is clearly untenable for large $|x|$ as the right hand side tends to a finite limit. □

We now elaborate the implications of Schanuel's conjecture for algebraic points on X. Schanuel's conjecture implies that the logarithms of multiplicatively independent algebraic numbers are algebraically independent over $\mathbb{Q}$ (see e.g. [19]).

4.2. Proposition. *Assume Schanuel's conjecture (or just that the logarithms of multiplicatively independent algebraic numbers are algebraically independent). Then if $x, y, z \in (0, \infty)$ are algebraic with $\log x \log y = \log z$ then either $(x, y, z) = (x, 1, 1)$ for some $x \in \overline{\mathbb{Q}}$, or $(x, y, z) = (1, y, 1)$ for some $y \in \overline{\mathbb{Q}}$.*

Proof. Suppose x, y, z are algebraic numbers in $(0, \infty)$ with $\log x \log y = \log z$. Then x, y, z are multiplicatively dependent, and we have

$$x^a y^b z^c = 1$$

for certain integers a, b, c. If two of a, b, c equal 0 then one of $x, y, z = 1$ and then we have either $x = z = 1$ and y arbitrary or $y = z = 1$ and x arbitrary.

Suppose that just one of a, b, c is zero, assuming $x, y, z \neq 1$. If $c = 0$ we have $y = x^r$ for some rational $r \neq 0$ and $r(\log x)^2 = \log z$. Then x, z must (by Schanuel) be multiplicatively related, say $z = x^s$ for some $s \in \mathbb{Q}^*$ and $r(\log x)^2 = s \log x$ implies $\log x = 0$ (contrary to our assumptions) or $\log x \in \mathbb{Q}^*$, whence x is non-algebraic. If $a = 0$, then $z = y^r$ for some $r \in \mathbb{Q}^*$ and $\log x \log y = r \log y$ implies (as $\log y \neq 0$) that $\log x \in \mathbb{Q}^*$ and is not algebraic.

Suppose then that none of a, b, c is zero. Then z depends multiplicatively on x and y and we get a relation $r \log x + s \log y = \log x \log y$ with r, s non-zero rational numbers. Then x, y must be multiplicatively related, and we find that $\log x$ is algebraic and hence $x = 1$. □

4.3. Summary. The set X^{alg} consists of infinitely many real semi-algebraic curves: the lines L_x, L_y and, for each $q \in \mathbb{Q}^*$, the curves $\Gamma_{x,q}, \Gamma_{y,q}$. By the Hermite-Lindemann theorem the curves $\Gamma_{x,q}, \Gamma_{y,q}$ contain no algebraic points. The lines L_x, L_y evidently contain algebraic points. Under Schanuel's Conjecture, $X^0 (\supset X^{\text{trans}})$ contains no algebraic points.

5. Pfaffian sets and Gabrielov-Vorobjov bounds

Definition 5.1 and the key result Theorem 5.3 are taken from the paper [7] of Gabrielov and Vorobjov.

5.1. Definition. ([7, Definition 2.1]) A *pfaffian chain* of order $r \geq 0$ and degree $\alpha \geq 1$ in an open domain $G \subset \mathbb{R}^n$ is a sequence of analytic functions $f_1, \ldots, f_r$ in G satisfying differential equations

$$df_j = \sum_{i=1}^{n} g_{ij}(x, f_1(x), \ldots, f_j(x)) dx_i$$

for $1 \le j \le r$, were $g_{ij} \in \mathbb{R}[x_1, \ldots, x_n, y_1, \ldots, y_j]$ are polynomials of degree not exceeding α. A function

$$f = P(x_1, \ldots, x_n, f_1, \ldots, f_r)$$

where $P \in \mathbb{R}[x_1, \ldots, x_n, y_1, \ldots, y_r]$ is a polynomial of degree not exceeding $\beta \ge 1$ is called a *pfaffian function* of order r and degree (α, β).

5.2. Definition. By a *pfaffian set* we will mean the set of common zeros of some pfaffian functions. By a *pfaff curve* we mean the graph of a pfaffian function of one variable on a connected subset of $\mathbb{R}$.

In the above definition no restriction is placed on the domain G. To obtain complexity bounds on pfaffian sets, one must impose restrictions on G (as we will do, following [7]), or allow more complicated domains whose complexity contributes to the complexity of the pfaffian sets. By a *simple domain* $G \subset \mathbb{R}^n$ we mean, as in [7], that G is a domain of the form $\mathbb{R}^n, [-1,1]^n, (0,\infty)^n$ or $\{x : ||x||^2 < 1\}$. The number of connected components of a set Y is denoted $\mathrm{cc}(Y)$.

5.3. Theorem. ([7, Corollary 3.3]) *Let $h_1, \ldots, h_\ell$ be pfaffian functions in a simple domain $G \subset \mathbb{R}^n$ having a common pfaffian chain of order r and degrees (α, β_i) respectively. Put $\beta = \max_i \beta_i$. Let Y be the pfaffian set*

$$Y = \{x \in G : h_1(x) = \ldots = h_\ell(x) = 0\}$$

Then

$$\mathrm{cc}(Y) \le 2^{r(r-1)/2+1}\beta(\alpha + 2\beta - 1)^{n-1}\left((2n-1)(\alpha+\beta) - 2n + 2\right)^r. \ \square$$

Observe that the bound on $\mathrm{cc}(Y)$ does not depend on ℓ. When the ambient space $\mathbb{R}^n$ and the pfaffian chain are fixed, as they will be, this fixes n, r, α and then we have

$$\mathrm{cc}(Y) \le c(n, r, \alpha)\beta^{n+r}.$$

6. Proof of Theorems 1.1 and 1.6

Theorems 1.1 and 1.6 concern the surface

$$X = \{(x, y, z) \in (0, \infty)^3 : \log x \log y = \log z\}.$$

If $\log x = 0$ then $\log z = 0$ also, so $X \cap \{x = 1\} = \{(x, y, z) : x = z = 1\} \subset X^{\mathrm{alg}}$. Likewise $X \cap \{y = 1\} \subset X^{\mathrm{alg}}$, while if $\log z = 0$ we must have $\log x = 0$ or $\log y = 0$, so that $X \cap \{z = 1\} \subset X^{\mathrm{alg}}$ too. In studying $(X - X^{\mathrm{alg}})(F, T)$ we may therefore assume that $x, y, z \ne 1$. Let

$$\mathcal{X} = \{(x, y, z) \in (0, 1)^3 : \log x \log y = -\log z\}.$$

The surface $\mathcal{X}$ contains semi-algebraic curves corresponding to fixing a rational negative value for $\log x$ or $\log y$. However, these curves contain no algebraic points (the corresponding x or y is transcendental by the Hermite-Lindemann Theorem). Thus $\mathcal{X}^{\mathrm{alg}}(\overline{\mathbb{Q}})$ is empty, and we need not restrict our counting to $\mathcal{X}^{\mathrm{trans}}$.

If $(x, y, z) \in X(F, T)$ with $x > 1, y > 1$ then $z > 1$ also. Since $H(\alpha) = H(1/\alpha)$ for any nonzero algebraic number α, we see that $(1/x, 1/y, 1/z) \in \mathcal{X}(F, T)$. If $(x, y, z) \in X(F, T)$ with $x < 1, y > 1$ then $z < 1$ and now $(x, 1/y, z) \in \mathcal{X}(F, T)$. The cases $x > 1, y < 1$ and $x, y < 1$ are similar and we see that, up to a finite multiplicative factor, Theorems 1.1 and 1.6 follow from the following result concerning $\mathcal{X}$.

6.1. Theorem. *Let $F \subset \mathbb{R}$ be a numberfield of degree f over $\mathbb{Q}$ and let $\epsilon > 0$. Then*

$$\#\mathcal{X}(F, T) \le c(\mathcal{X}, f, \epsilon)\Big(\log T\Big)^{44+\epsilon}.$$

Proof. It suffices to prove a bound of the stated form for $\#\mathcal{X}^{\text{size}}(F,T)$. For each integer $g>1$ we have a $(J(g),A(g),1+1/g)$-mild parameterization (with $J(g)=1$) of $\mathcal{X}$ given by

$$\theta:(0,1)^2\to(0,1)^3,$$

$$\theta(s,t)=(\exp\left(1-\frac{1}{s^g}\right),\exp\left(1-\frac{1}{t^g}\right),\exp\left(-\left(1-\frac{1}{s^g}\right)\left(1-\frac{1}{t^g}\right)\right)).$$

By Theorem 3.2, $\mathcal{X}^{\text{size}}(F,T)$ is contained in

$$\leq c(g,f)\Big(\log T\Big)^{9(1+1/g)(1+o(1))}$$

intersections of $\mathcal{X}$ with hypersurfaces of degree

$$\left[\Big(\log T\Big)^2\right]$$

with the $1+o(1)$ as $T\to\infty$ (and implicit constants depending only on g,f). These intersections all have dimension 1, since $\mathcal{X}$ is not semi-algebraic, and we may ignore any semi-algebraic components, as the semi-algebraic curves in $\mathcal{X}$ contain no algebraic points.

The mild parameterization plays no further role in the study of these hypersurface intersections. In applying the Gabrielov-Vorobjov bounds it is advantageous to define them as pfaffian sets with as low degree as possible. For the remainder of the proof we therefore consider $\mathcal{X}$ to be parameterized by

$$(0,\infty)^2\to(0,1)^3,$$

$$(p,q)\mapsto\left(e^{-p},e^{-q},e^{-pq}\right)=(x,y,z)\in\mathcal{X}.$$

If $H\in\mathbb{R}[x,y,z]$ defines the hypersurface $V_H:H(x,y,z)=0$ then the intersection $\mathcal{X}\cap V_H$ is the image of the exponential-algebraic curve (not necessarily connected) in the (p,q)-plane defined by

$$K(p,q)=H(e^{-p},e^{-q},e^{-pq})=0,\quad p,q>0.$$

We observe that, for $H\neq 0$, the equation $K(p,q)=0$ defines a curve $V=V_K$, i.e. a set of dimension 1, again because $\mathcal{X}$ is not semi-algebraic. The set of singular points V_s of V is defined by

$$K=0,\quad K_p=-H_xe^{-p}-qH_ze^{-pq}=0,\quad K_q=-H_ye^{-q}-pH_ze^{-pq}=0.$$

It is a finite set (definable of dimension zero).

We now follow the procedure of [10, 11], substituting Gabrielov-Vorobjov bounds for the appeals made in [10, 11] to Gabrielov's Theorem for subanalytic sets.

Let then Π be a coordinate plane in $\mathbb{R}^3$ whose coordinates we denote (u,v). Projection of $\mathbb{R}^3$ onto Π takes the curve V defined by $K(p,q)=0$ into some curve in Π. At a point $P=(p,q)$ of $V-V_s$, V is locally an analytic curve. If $K_q\neq 0$ at P then we may use q as a local parameter and we find that u is nonconstant at P unless

$$u_pK_q-u_qK_p=0.$$

Similarly, v is nonconstant at P unless

$$v_pK_q-v_qK_p=0.$$

Let V_u be the subset of $V-V_s$ where one or more of these quantities vanish. At points of $V-V_s-V_u$ the slope du/dv is well defined, and the image of V in Π is locally the graph of a function. We proceed to derive an expression for its derivatives. We have, locally,

$$u=u(p(v),q(v)),\quad v=v(p(v),q(v)),\quad K(p(v),q(v))=0.$$

Differentiating the second and third equations implicitly,

$$1 = v_p p' + v_q q', \quad K_p p' + K_q q' = 0$$

which we may write as a matrix equation

$$\begin{pmatrix} v_p & v_q \\ K_p & K_q \end{pmatrix} \begin{pmatrix} p' \\ q' \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

giving

$$\begin{pmatrix} p' \\ q' \end{pmatrix} = \frac{1}{v_p K_q - v_q K_p} \begin{pmatrix} K_q & -v_q \\ -K_p & v_p \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \frac{1}{v_p K_q - v_q K_p} \begin{pmatrix} K_q \\ -K_p \end{pmatrix}.$$

We have then

$$\frac{du}{dv} = u_p p' + u_q q' = \frac{u_p K_q - u_q K_p}{v_p K_q - v_q K_p}.$$

To get expressions for higher derivatives, we differentiate this expression with respect to v and use the expressions we have for p', q'. For points (u, v) with $v_p K_q - v_q K_p \neq 0$ and a positive integer m we will have

$$\frac{d^m u}{dv^m} = \frac{R_m(u, v, K)}{(v_p K_q - v_q K_p)^{2m-1}}$$

for a suitable differential polynomial R_m.

We want to estimate the number of zeros of R_m which we will do by controlling its order and degree as a pfaffian function. Let us write

$$\Delta = v_p K_q - v_q K_p$$

(no confusion should arise with the previous use of Δ), which we consider as a function of v, so that

$$p' = \frac{K_q}{\Delta}, \quad q' = \frac{-K_p}{\Delta}$$

and

$$\Delta' = \frac{v_{pp} K_q^2 - v_{pq} K_p K_q + v_p K_{qp} K_q - v_p K_{qq} K_p - v_{qp} K_q K_p + v_{qq} K_p^2 - v_q K_{pp} K_q + v_q K_{pq} K_p}{\Delta} = \frac{\Gamma}{\Delta}.$$

If we now write

$$\frac{d^m u}{dv^m} = \frac{R_m}{\Delta^{2m-1}}, \quad R_m' = \frac{S_m}{\Delta}$$

then

$$\frac{d^{m+1} u}{dv^{m+1}} = \frac{\Delta^{2m-2} \frac{\Delta S_m}{\Delta} - (2m-1)\Delta^{2m-2} \frac{\Gamma}{\Delta} R_m}{\Delta^{4m-2}} = \frac{R_{m+1}}{\Delta^{2(m+1)-1}}$$

gives a recurrence for R_m (and validates the asserted form for $d^m u / dv^m$), starting with

$$R_1 = u_p K_q - u_q K_p.$$

Consider the pfaffian chain of functions on $(0, \infty)^2$

$$f_1 = e^{-p}, f_2 = e^{-q}, f_3 = e^{-pq}$$

where we have $\partial_p f_3 = -q f_3, \partial_q f_3 = -p f_3$. This is then a pfaffian chain of order $r = 3$ and degree $\alpha = 2$. The function u, v, K and their partial derivatives with respect to p, q are pfaffian with this chain, i.e. they are polynomials in p, q, f_1, f_2, f_3, and therefore so are all the functions R_m and S_m, and they therefore have order 3 and degree $(2, \beta)$, where $\beta \geq 1$ is their degree as a polynomial in p, q, f_1, f_2, f_3.

Claim. u_μ has degree $(2, |\mu| + 1)$.

Proof of Claim. By induction. It holds for $|\mu| = 1$, the "worst case" being $u = f_3 = e^{-pq}$ for which $u_p = -qf_3$ is a polynomial of degree 2. Suppose the Claim is true for all μ with $|\mu| \le m$. Then, with some polynomial P of degree $\le |\mu| + 1$,

$$\partial_p \partial_\mu u = \partial_p P(p, q, f_1, f_2, f_3) = P_p + P_{f_1}(f_1)_p + P_{f_3}(f_3)_p = P_p - P_{f_1} f_1 - qP_{f_3} f_3$$

having degree $\le |\mu| + 2$. The $\partial_q \partial_\mu$ calculation is similar. □

If H has degree d then $K = H(f_1, f_2, f_3)$ is pfaffian with the chain f_1, f_2, f_3 and degree $(2, d)$. Generalizing the previous Claim we find:

Claim. K_μ has degree $(2, d + |\mu|)$. □

Returning to our functions R_m and S_m, we have that $R_1 = u_p K_q - u_q K_p$ has degree $(2, d + 3)$. Suppose R_m has degree $(2, \rho_m)$, so $R_m = P(p, q, f_1, f_2, f_3)$ for suitable polynomial P of degree $\le \rho_m$. Then

$$R'_m = P_p p' + P_q q' + P_{f_1} f'_1 + P_{f_2} f'_2 + P_{f_3} f'_3 =$$
$$\frac{P_p K_q - P_q K_p - P_{f_1} f_1 K_q + P_{f_2} f_2 K_p - q f_3 P_{f_3} K_q + p f_3 P_{f_3} K_p}{\Delta}.$$

Thus the degree $(2, \sigma_m)$ of S_m where $R'_m = S_m/\Delta$ is

$$\sigma_m = (\rho_m - 1) + 2 + d + 1 = \rho_m + d + 2.$$

Since $\deg \Gamma = (2, 2d + 5)$ by applying the above Claims to the exhibited expression for Γ and $\deg(\Delta) = (2, d + 3)$ we find that

$$\rho_{m+1} = \max(d + 3 + \rho_m + d + 2, 2d + 5 + \rho_m) = \rho_m + 2d + 5$$

and therefore

$$\rho_m = m(2d + 5) - (d + 2).$$

With these degrees in hand, we consider the decomposition of the curve V_K defined by $K(p, q) = 0$ into "good" curves, where a "good" curve is a connected subset whose projection into each coordinate plane Π is a "good" graph with respect to one or other of the axes, namely, the graph of a function ϕ which is smooth (indeed analytic) on an interval, has slope of absolute value at most 1 at each point, and such that the derivative of $\phi^{(m)}$ of each order $m = 1, \ldots, M$ is either non-vanishing in the interior of the interval or identically zero.

In the following, constants in $<<$ depend on a pfaffian chain on a simple domain G. This will always be the chain f_1, f_2, f_3 of order 3 and degree 2 in the simple domain $p, q > 0$ in $\mathbb{R}^2$, so that the implicit constant is then absolute and explicit from Theorem 5.3.

The set $V = V_K$ has $<< d^5$ connected components. Its singular set V_s is defined by $K = 0, K_p = K_q = 0$ where K_p, K_q have degree at most $d + 1$. So

$$\mathrm{cc}(V_s) << d^5$$

and therefore also

$$\mathrm{cc}(V - V_s) << d^5.$$

Let V_u be the subset of $V - V_s$ where du/dv is undefined. Considering the conditions exhibited above for such points, and also for the set V_a where the slope of the graph in Π is ± 1 we have again

$$\mathrm{cc}(V - V_s - V_u - V_a) << d^5.$$

Now take one such component, fix a coordinate plane Π, and consider the points where some $R_m = 0$. Since $\deg(R_m) \le (2, (2d + 5)m)$, we have at most $m^5(2d + 5)^5$ points where $R_m = 0$, unless it vanishes

identically on the component. In this case the image in Π is the graph of a polynomial with respect to one of the axes. If the graph is not a polynomial than, summing over $m = 1, 2, \ldots, M$, we have at most $<< M^6 d^5$ further components, whose slope lies in $[-1, 1]$, and for which no derivative up to order M vanishes. Taking the isolated points where some $R_m = 0$ for $m = 1, 2, \ldots, M$ for each of the 3 coordinate planes Π, we find that V_K decomposes into $<< M^6 d^5$ connected components whose image in each coordinate plane is a graph with respect to one of the axes with slope in $[-1, 1]$ and such that, for each $m = 1, 2, \ldots, M$, R_m is nonzero in the interior or identically zero on the component, i.e. "good" components.

If such a connected component of V_K is semi-algebraic then its projection in each coordinate plane Π will be algebraic, and conversely if all the projections are semi-algebraic then the component is semi-algebraic. Now we need not consider algebraic components, therefore we can assume that every component has a non-algebraic (and hence non-polynomial) projection into one of the planes Π.

Let W be a "good" component of V_K, and Y its non-semi-algebraic image in some Π. If we intersect Y with a plane algebraic curve (in Π) defined by $L(u, v) = 0$ of degree b, then since the function $L(u(p, q), v(p, q))$ is pfaffian of degree $(2, b)$, intersecting with Y gives again at most

$$<< \max(b, d)^5$$

connected components. So $Y \cap \{L = 0\}$ consists of at most this many isolated points.

Since Y is a "good" graph then, by [13] (for rational points) and [14, 6.7] (for F-points), $Y^{\text{size}}(F, T)$ is contained in

$$c(f) M \log T$$

plane algebraic curves of degree b where $M = (b + 1)(b + 2)/2$. So we get

$$\#Y^{\text{size}}(F, T) \leq c(f) \max(b, d)^5 M \log T$$

and the same estimate holds for the corresponding component of V_H, where having a point of $\mathcal{X}^{\text{size}}(F, T)$ requires that the other coordinate be also in F with its H^{size} bounded by T.

Putting all the above together, we find

$$\#\mathcal{X}^{\text{size}}(F, T) \leq c(\mathcal{X}, f, g) \Big(\log T \Big)^{9(1+1/g)(1+o(1))} M^6 d^5 M \log T \max(b, d)^5$$

where $d = \Big[\Big(\log T \Big)^2 \Big]$, $M = (b + 1)(b + 2)/2$, and $b = \Big[\log T \Big]$, giving

$$\#\mathcal{X}^{\text{size}}(F, T) \leq c(\mathcal{X}, f, g) \Big(\log T \Big)^{9(1+1/g)(1+o(1))+35}.$$

This completes the poof of Theorem 6.1, and thereby establishes Theorems 1.1 and 1.6 as well. □

7. Appendix: O-minimal structures

We give the basic definitions, following [22], referring the reader to [5, 6, 21, 22] for more information.

7.1. Definition. A *pre-structure* is a sequence $\mathcal{S} = (\mathcal{S}_n : n \geq 1)$ where each $\mathcal{S}_n$ is a collection of subsets of $\mathbb{R}^n$. A pre-structure $\mathcal{S}$ is called a *structure (over the real field)* if, for all $n, m \geq 1$, the following conditions are satisfied:

(1) $\mathcal{S}_n$ is a boolean algebra (under the usual set-theoretic operations)
(2) $\mathcal{S}_n$ contains every semi-algebraic subset of $\mathbb{R}^n$
(3) if $A \in \mathcal{S}_n$ and $B \in \mathcal{S}_m$ then $A \times B \in \mathcal{S}_{n+m}$
(4) if $m \geq n$ and $A \in \mathcal{S}_m$ then $\pi(A) \in \mathcal{S}_n$, where $\pi : \mathbb{R}^m \to \mathbb{R}^n$ is projection onto the first n coordinates.

If $\mathcal{S}$ is a structure and $X \subset \mathbb{R}^n$, we say X is *definable in* $\mathcal{S}$ if $X \in \mathcal{S}_n$. If $\mathcal{S}$ is a structure and, in addition,

(5) the boundary of every set in $\mathcal{S}_1$ is finite

then $\mathcal{S}$ is called an *o-minimal* structure (over the real field).

7.2. Definition. [5, p3] We denote by $\mathbb{R}_{\exp}$ the prestructure consisting of those sets in $\mathbb{R}^n$ arising as the image under projection maps $\mathbb{R}^{n+k} \to \mathbb{R}^n$ of sets of the form $\{(x,y) \in \mathbb{R}^{n+k} : P(x,y,e^x,e^y) = 0\}$ where P is a real polynomial in $2(n+k)$ variables, and where $x = (x_1,\ldots,x_n), y = (y_1,\ldots,y_k), e^x = (e^{x_1},\ldots,e^{x_n}), e^y = (e^{y_1},\ldots,e^{y_k})$.

7.3. Example. The set X is the image under the projection $\mathbb{R}^6 \to \mathbb{R}^3$ of

$$Y = \{(x,y,z,u,v,w) : (x-e^u)^2 + (y-e^v)^2 + (z-e^w)^2 + (uv-w)^2 = 0\}.$$

7.4. Theorem. (Wilkie [21]) $\mathbb{R}_{\exp}$ *is an o-minimal structure.* □

References

1. E. Bombieri, Algebraic values of meromorphic maps, *Inventiones* **10** (1970), 267–287.
2. E. Bombieri and W. Gubler, *Heights in diophantine geometry,* CUP 2006.
3. E. Bombieri and J. Pila, The number of integral points on arcs and ovals, *DMJ* **59** (1989), 337–357.
4. L. Butler, Some cases of Wilkie's conjecture, working paper, April 2009.
5. L. van den Dries, *Tame topology and o-minimal structures,* CUP 1998.
6. L. van den Dries and C. Miller, Geometric categories and o-minimal structures, *DMJ* **84** (1996), 497–540.
7. A. Gabrielov and N. Vorobjov, Complexity of computations with Pfaffian and Noetherian functions, in *Normal Forms, Bifurcations and Finiteness problems in Differential Equations,* Kluwer, 2004.
8. S. Lang, Algebraic values of meromorphic functions, *Topology* **3** (1965), 183–191.
9. S. Lang, *Introduction to transcendental numbers,* Addison-Wesley, Reading Mass., 1966.
10. J. Pila, Integer points on the dilation of a subanalytic surface, *QJM* **55** (2004), 207–223.
11. J. Pila, Rational points on a subanalytic surface, *AIF* **55** (2005), 1501–1516.
12. J. Pila, Mild parameterization and the rational points of a pfaff curve, *Commentarii Mathematici Universitatis Sancti Pauli* **55** (2006), 1–8 (and Erratum p231).
13. J. Pila, The density of rational points on a pfaff curve, *Ann. Fac. Sci. Toulouse,* **16** (2007), 635–645.
14. J. Pila, On the algebraic points of a definable set, *Selecta Math. N.S.* **15** (2009), 151–170.
15. J. Pila and A. J. Wilkie, The rational points of a definable set, *DMJ* **133** (2006), 591–616.
16. D. Roy, Interpolation formulas and auxiliary functions, *J. Number Theory* **94** (2002), 248–285.
17. M. Waldschmidt, Proprietés arithmétiques de fonctions de plusieurs variables. II. *Séminaire Pierre Lelong (Analyse), Année 1974/5,* Lecture Notes in Mathematics **524** (1976), 108–135.
18. M. Waldschmidt, Integer values entire functions on Cartesian products, *Number theory in progress, Vol. 1 (Zakopane-Koscieliko, 1997),* 553–576, de Gruyter, Berlin, 1999.
19. M. Waldschmidt, *Diophantine approximation on linear algebraic groups*, Grundlehren der mathematischen Wissenschaften **326**, Springer-Verlag, Berlin 2000.
20. M. Waldschmidt, Algebraic values of analytic functions, *J. Comput. Appl. Math.* **160** (2003), 323–333.
21. A. J. Wilkie, Model completeness results for expansions of the ordered field of real numbers by restricted Pfaffian functions and the exponential function, *J. Amer. Math. Soc.* **9** (1996), 1051–1094.
22. A. J. Wilkie, A theorem of the complement and some new o-minimal structures, *Selecta Math. N.S.* **5** (1999), 397–421.

School of Mathematics
University of Bristol
Bristol, BS8 1TW
UK

j.pila@bristol.ac.uk

Counting rational points on a certain exponential-algebraic surface

Jonathan Pila

Abstract

We study the distribution of rational points on a certain exponential-algebraic surface and we prove, for this surface, a conjecture of A. J. Wilkie.

2000 Mathematics Subject Classification: 11G99, 03C64
Keywords: O-minimal structure, rational points, transcendental numbers.

Running title:

Counting rational points

Comptage des points rationnels sur une surface exponentielle-algébrique particulière

Jonathan Pila

Résumé

Nous étudions la repartition des points rationnels sur une certaine surface exponentielle-algébrique et prouvons, pour cette surface, une conjecture de A. J. Wilkie.

Classification math.: 11G99, 03C64
Mots-clés: Structure o-minimale, points rationnels, nombres transcendants.