

Convergence of an Euler Scheme for a Hybrid Stochastic-Local Volatility Model with Stochastic Rates in Foreign Exchange Markets*

Andrei Cozma[†], Matthieu Mariapragassam[†], and Christoph Reisinger[†]

Abstract. We study the Heston–Cox–Ingersoll–Ross++ stochastic-local volatility model in the context of foreign exchange markets and propose a Monte Carlo simulation scheme which combines the full truncation Euler scheme for the stochastic volatility component and the stochastic domestic and foreign short interest rates with the log-Euler scheme for the exchange rate. We establish the exponential integrability of full truncation Euler approximations for the Cox–Ingersoll–Ross process and find a lower bound on the explosion time of these exponential moments. Under a full correlation structure and a realistic set of assumptions on the so-called leverage function, we prove the strong convergence of the exchange rate approximations and deduce the convergence of Monte Carlo estimators for a number of vanilla and path-dependent options. Then we perform a series of numerical experiments for an autocallable barrier dual currency note.

Key words. Heston model, shifted square-root process, calibration, Monte Carlo simulation, exponential integrability, strong convergence

AMS subject classifications. 60H35, 65C05, 65C30, 65N21

DOI. 10.1137/17M1114569

1. Introduction. The class of stochastic-local volatility (SLV) models, introduced by Jex, Henderson, and Wang [30], Lipton [34], and Lipton and McGhee [36], among others, has become very popular in the financial sector in recent years. They contain a stochastic volatility component as well as a local volatility component (called the leverage function) and combine advantages of the two. According to Ren, Madan, and Qian [40], Tian et al. [42], and van der Stoep, Grzelak, and Oosterlee [43], SLV models allow for a better calibration to European options and improve the pricing and risk-management performance when compared to pure local volatility (LV) or pure stochastic volatility (SV) models, and Lipton, Gal, and Lasis [35] recently referred to them as the de facto standard for pricing foreign exchange (FX) options. We focus on the Heston SLV model because the Cox–Ingersoll–Ross (CIR) process for the variance is widely used in the industry due to its desirable properties, such as mean-reversion and nonnegativity, and since semianalytic formulae are available for calls and puts under Heston’s model and can help calibrate the parameters easily. The local volatility component allows a perfect calibration to the market prices of vanilla options. At the same time, the stochastic volatility component already provides built-in smiles and skews which give a rough calibration, so that a relatively flat leverage function suffices for a perfect calibration.

In order to improve the pricing and hedging of FX options, we introduce stochastic do-

*Received by the editors February 1, 2017; accepted for publication November 3, 2017; published electronically January 24, 2018.

<http://www.siam.org/journals/sifin/9-1/M111456.html>

[†]Mathematical Institute, University of Oxford, Oxford OX2 6GG, UK (andrei.cozma@maths.ox.ac.uk, matthieu.mariapragassam@maths.ox.ac.uk, christoph.reisinger@maths.ox.ac.uk).

mestic and foreign short interest rates into the model. Assuming constant interest rates is appealing due to its simplicity and does not lead to a serious mispricing of options with short maturities. However, empirical results [45] have confirmed that the constant interest rate assumption is inappropriate for long-dated FX products, and the effect of interest rate volatility can be as relevant as that of the FX rate volatility for longer maturities. There has been a great deal of research carried out in the area of option pricing with stochastic volatility and interest rates in the past couple of years. For instance, Van Haastrecht et al. [45] extended the model of Schöbel and Zhu [41] to currency derivatives by including stochastic interest rates, a model that benefits from analytical tractability even in a full correlation setting due to the processes being Gaussian, whereas Ahlip and Rutkowski [1], Grzelak and Oosterlee [19], and Van Haastrecht and Pelsler [46] examined the Heston–2CIR/Vasicek hybrid models and concluded that a full correlation structure gives rise to a nonaffine model even under a partial correlation of the driving Brownian motions.

There has also been an increasing interest in the calibration to vanilla options of models with stochastic and local volatility and stochastic interest rate dynamics. For instance, Guyon and Henry-Labordère [21] examined Monte Carlo–based calibration methods for a 3-factor SLV equity model with a stochastic domestic rate and discrete dividends, while Mariapragassam, Cozma, and Reisinger [11] and Deelstra and Rayee [13] examined 4-factor hybrid SLV models with stochastic domestic and foreign rates and proposed different approaches to calibrate the leverage function. Hambly, Mariapragassam, and Reisinger [22] took a different path and discussed the calibration of a 2-factor SLV model to barriers. As an aside, in this paper, the spot FX rate is defined as the number of units of domestic currency per unit of foreign currency.

The model of Cox, Ingersoll, and Ross [10] is very popular when modeling short rates because the square-root (CIR) process admits a unique strong solution, is mean-reverting, and is analytically tractable. Cox et al. found the conditional distribution to be noncentral chi-squared, and Broadie and Kaya [8] proposed an efficient exact simulation scheme for the square-root process based on acceptance-rejection sampling. However, their algorithm presents a number of disadvantages such as complexity and lack of speed, and it is not fit to price strongly path-dependent options that require the value of the FX rate at a large number of time points. Furthermore, in the context of an SLV model, the correlations between the underlying processes make it difficult to simulate a noncentral chi-squared increment together with a correlated increment for the FX rate and the short rates, if applicable.

As of late, the nonnegativity of the CIR process is considered to be less desirable when modeling short rates. On the one hand, the central banks have significantly reduced the interest rates since the 2008 financial crisis and it is now commonly accepted that interest rates need not be positive. On the other hand, if interest rates dropped too far below zero, then large amounts of money would be withdrawn from banks and government bonds, putting a severe squeeze on deposits. Hence, we model the domestic and foreign short rates using the shifted CIR (CIR++) process of Brigo and Mercurio [7]. The CIR++ model allows the short rates to become negative and can fit any observed term structure exactly while preserving the analytical tractability of the original model for bonds, caps, swaptions, and other basic interest rate products.

In the present work, we put forward the Heston–Cox–Ingersoll–Ross++ stochastic-local

volatility (Heston–2CIR++ SLV) model to price FX options. Independent of the correlation structure, the model is nonaffine and hence a closed-form solution to the European option valuation problem is not available. Finite-difference methods are popular in finance and when the evolution of the exchange rate is governed by a complex system of stochastic differential equations (SDEs), it all comes down to solving a higher-dimensional partial differential equation (PDE). This can prove to be difficult due to the curse of dimensionality, because the number of grid points required increases exponentially with the number of dimensions. Monte Carlo algorithms are often preferred due to their ability to handle path-dependent features easily and there are numerous discretization schemes available, like the simple Euler–Maruyama scheme; see, e.g., Glasserman [18]. However, there are several disadvantages of this discretization, such as the fact that the approximation process can become negative with nonzero probability. In practice, one can set the process equal to zero when it turns negative—called an absorption fix—or reflect it in the origin—referred to as a reflection fix. An overview of the Euler schemes considered thus far in the literature, including the full truncation scheme, can be found in Lord, Koekkoek, and van Dijk [37].

The usual theorems in Kloeden and Platen [33] on the convergence of numerical simulations require the drift and diffusion coefficients to be globally Lipschitz and satisfy a linear growth condition, whereas Higham, Mao, and Stuart [24] extended the analysis to a simple Euler scheme for a locally Lipschitz SDE. The standard convergence theory does not apply to the CIR process since the square-root is not locally Lipschitz around zero. Consequently, alternative approaches have been employed to prove the weak or strong convergence of various discretizations for the square-root process. Alfonsi [2], Deelstra and Delbaen [12], Higham and Mao [23], and Lord, Koekkoek, and van Dijk [37] examined the strong global approximation and found either a logarithmic convergence rate or none at all. Strong convergence of order 1/2 of the symmetrized and the backward (drift-implicit) Euler–Maruyama (BEM) schemes was established in Berkaoui, Bossy, and Diop [6] and Dereich, Neuenkirch, and Szpruch [14], respectively, albeit in a very restricted parameter regime for the symmetrized scheme. Alfonsi [3] and Neuenkirch and Szpruch [38] recently showed that the BEM scheme for the SDE obtained through a Lamperti transformation is strongly convergent with rate one in the case of an inaccessible boundary point, while Hutzenthaler, Jentzen, and Noll [28] established a positive strong order of convergence in the case of an accessible boundary point.

Hutzenthaler, Jentzen, and Wang [29] identified a class of stopped increment-tamed Euler approximations for nonlinear systems of SDEs and proved that they preserve the exponential integrability of the exact solution under some mild assumptions, unlike the explicit, the linear-implicit, and some tamed Euler schemes, which rarely do. In this work, we consider a different exponential functional of the CIR process and prove that an explicit Euler scheme retains exponential integrability properties, a result which plays a key role in establishing the boundedness of moments of Euler approximations for SDE systems with CIR dynamics in one or more dimensions.

To the best of our knowledge, the convergence of Monte Carlo algorithms in a stochastic-local volatility context has yet to be established. Higham and Mao [23] considered an Euler simulation of the Heston model with a reflection fix in the diffusion coefficient to avoid negative values. They studied convergence properties of the stopped approximation process and used the boundedness of payoffs to prove weak convergence for a European put and an up-and-out

call option. However, Higham and Mao mentioned that the arguments cannot be extended to cope with unbounded payoffs. We work under a different Euler scheme and overcome this problem by proving the uniform boundedness of moments of the true solution and its approximation, and then the strong convergence of the latter. The existence of moment bounds for Euler approximations of the Heston model is important for deriving strong convergence [24] and it has been a long-standing open problem until now. Furthermore, the existence of moment bounds for hybrid Heston-type SLV models plays an important role in the calibration routine [11].

In this paper, we focus on the Heston SLV model with CIR++ short interest rates and examine convergence properties of the Monte Carlo algorithm with the full truncation Euler (FTE) discretization for the squared volatility and the two short rates, and the log-Euler discretization for the exchange rate. We prefer the full truncation scheme introduced by Lord, Koekkoek, and van Dijk [37] because it preserves the positivity of the original process, is easy to implement, and is found empirically to produce the smallest bias of all explicit Euler schemes and to outperform the quasi-second order schemes of Kahl and Jäckel [31] and Ninomiya and Victoir [39]. We also choose the FTE scheme over the BEM scheme since the latter works only in a restricted parameter regime that is often unrealistic, since the BEM scheme is equivalent to the implicit Milstein method employed by Kahl and Jäckel to first order in the time step, and since the quanto correction term in the dynamics of the foreign short rate would lead to technical challenges in the convergence analysis.

The quadratic exponential (QE) scheme of Andersen [4] uses moment-matching techniques to approximate the noncentral chi-squared distributed square-root process and is an efficient alternative to the FTE scheme that typically produces a smaller bias at the expense of a more complex implementation. Empirical results suggest that the QE scheme outperforms the FTE scheme in the case of a severely violated Feller condition [44], whereas when the Feller condition is satisfied, the two schemes perform comparably well because the bias with the FTE scheme is small even when only a few number of time steps are used per year. Since there is no natural extension to the QE scheme to multidimensional Heston-type models that involve several correlated square-root processes such as the ones studied here, we prefer the FTE scheme in the subsequent convergence analysis.

In summary, to the best of our knowledge, this paper is the first to establish (1) the exponential integrability of the FTE scheme for the CIR process; (2) the boundedness of moments of order greater than 1 of approximation schemes for Heston-type models; (3) the strong convergence of approximation schemes for models with SLV (with a bounded and Lipschitz leverage function) and stochastic CIR++ short rates; and (4) the weak convergence for options with unbounded payoffs, in particular, for European, Asian, and barrier contracts (up to a critical time).

The remainder of this paper is structured as follows. In section 2, we introduce the model and discuss the postulated assumptions and an efficient calibration. In section 3, we first define the simulation scheme and discuss the main theorem. Then we investigate the uniform exponential integrability of the full truncation scheme for the square-root process and prove convergence of the exchange rate approximations. We conclude the section by establishing the convergence of Monte Carlo simulations for computing the expected discounted payoffs of European, Asian, and barrier options. In section 4, we carry out numerical experiments

to justify our choice of model and to demonstrate convergence. Finally, [section 5](#) contains a short discussion.

2. Preliminaries.

2.1. Model definition. In its most general form, we have in mind a model in an FX market, for the spot FX rate S , the squared volatility of the FX rate v , the domestic short interest rate r^d , and the foreign short interest rate r^f . Unless otherwise stated, in this paper, the subscripts and superscripts d and f indicate domestic and foreign, respectively. Consider a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ and suppose that the dynamics of the underlying processes are governed by the system of SDEs under the domestic risk-neutral measure \mathbb{Q} :

$$(2.1) \quad \begin{cases} dS_t = (r_t^d - r_t^f)S_t dt + \sigma(t, S_t)\sqrt{v_t}S_t dW_t^s, & S_0 > 0, \\ dv_t = k(\theta - v_t)dt + \xi\sqrt{v_t}dW_t^v, & v_0 > 0, \\ r_t^d = g_t^d + h_d(t), \\ dg_t^d = k_d(\theta_d - g_t^d)dt + \xi_d\sqrt{g_t^d}dW_t^d, & g_0^d > 0, \\ r_t^f = g_t^f + h_f(t), \\ dg_t^f = (k_f\theta_f - k_f g_t^f - \rho_{sf}\xi_f\sigma(t, S_t)\sqrt{v_t g_t^f})dt + \xi_f\sqrt{g_t^f}dW_t^f, & g_0^f > 0, \end{cases}$$

where W^s , W^v , W^d , and W^f are one-dimensional standard \mathcal{F}_t -adapted Brownian motions that are possibly correlated, σ is the leverage function, and h_d and h_f are deterministic functions of time. The mean-reversion parameters k , k_d , and k_f , the long-term mean parameters θ , θ_d , and θ_f , and the volatility parameters ξ , ξ_d , and ξ_f are positive real numbers. The quanto correction term in the drift of the foreign short rate in (2.1) comes from changing from the foreign to the domestic risk-neutral measure [9]. As an aside, for Hull–White short rate processes, changing from the domestic spot measure to the domestic T -forward measure leads to a dimension reduction of the problem because the diffusion coefficients of the short rates do not depend on the level of the rates [20]. Since this is not the case for CIR short rate processes, we prefer to work under the domestic spot measure. Note that the above system can collapse to the Heston–2CIR++ model if we set $\sigma = 1$, or to a local volatility model with stochastic short rates if we set $k = \xi = 0$. The standard Heston SLV model is the special case $k_d = \xi_d = k_f = \xi_f = h_{d,f} = 0$. We can also think of (2.1) as a model in an equity market with stock price process S , stochastic interest rate r^d , and stochastic dividend yield r^f , in which case the quanto correction term vanishes. We consider a full correlation structure between the Brownian drivers W^s , W^v , W^d , and W^f , i.e., no assumptions on the constant correlation matrix Σ are made, where

$$(2.2) \quad \Sigma = \begin{bmatrix} 1 & \rho_{sv} & \rho_{sd} & \rho_{sf} \\ \rho_{sv} & 1 & \rho_{vd} & \rho_{vf} \\ \rho_{sd} & \rho_{vd} & 1 & \rho_{df} \\ \rho_{sf} & \rho_{vf} & \rho_{df} & 1 \end{bmatrix}.$$

Fix a time horizon $T > 0$. We work under the following assumptions.

(A1) There exists a nonnegative constant σ_{max} such that, for all $t \in [0, T]$ and $x \in [0, \infty)$, we have

$$(2.3) \quad 0 \leq \sigma(t, x) \leq \sigma_{max}.$$

(A2) There exist $N_T \in \mathbb{N}$, a positive real number α , and nonnegative constants A, B , and $(C_j)_{1 \leq j \leq N_T}$ such that, for all $t, u \in [0, T]$ and $x, y \in [0, \infty)$, we have

$$(2.4) \quad |\sigma(t, x) - \sigma(u, y)| \leq A|t - u|^\alpha + B|x - y| + \sum_{j=1}^{N_T} C_j \mathbb{1}_{t \wedge u < \frac{jT}{N_T} \leq t \vee u}.$$

(A3) There exists a nonnegative constant h_{max} such that, for all $t \in [0, T]$ and $i \in \{d, f\}$, we have

$$(2.5) \quad |h_i(t)| \leq h_{max}.$$

Hence, we assume that σ is bounded, piecewise Hölder continuous in time, and Lipschitz continuous in spot. According to [11], for the leverage function to be consistent with call and put prices, it has to be given by the formula (2.8), which depends on the calibrated Dupire local volatility. In practice, the leverage function is defined on a grid of points (20 points per year in time and 30 points in space usually suffice for an acceptable calibration error [11, 21]), interpolated flat-forward in time and using cubic splines in spot, and extrapolated flat outside an interval. Therefore, there is no loss of generality from a practical point of view in making these assumptions. We also assume that $h_{d,f}$ are bounded. According to [7], for a perfect fit to the initial term structure of interest rates, each of the two shift functions must be given by the difference between the (flat-forward) market instantaneous forward rate and a continuous function of time. Then $h_{d,f}$ are piecewise continuous and the third assumption holds.

2.2. Model calibration. The calibration of the 4-factor Heston-2CIR++ SLV model (2.1) is of paramount importance and represents a mandatory step for the efficient pricing of derivative contracts. In [11], we consider the 4-factor SLV model and propose a new calibration approach that builds on the particle method of [21], combined with a novel and efficient variance reduction technique that takes advantage of PDE calibration to increase its stability and accuracy. The numerical experiments in [11] suggest that this method almost recovers the calibration speed from the associated 2-factor SLV model with deterministic rates. We assume here a partial correlation structure where only ρ_{sv} , ρ_{sd} , and ρ_{sf} may be nonzero, and we denote by D^d and D^f the domestic and foreign discount factors associated with their respective money market accounts; i.e., for any $t \in [0, T]$,

$$(2.6) \quad D_t^d = e^{-\int_0^t r_u^d du} \quad \text{and} \quad D_t^f = e^{-\int_0^t r_u^f du}.$$

In addition to (2.1), we consider a pure local volatility (LV) model,

$$(2.7) \quad dS_t^{LV} = (f_t^d - f_t^f) S_t^{LV} dt + \sigma_{LV}(t, S_t^{LV}) S_t^{LV} dW_t^s, \quad S_0^{LV} = S_0,$$

where for $i \in \{d, f\}$, $f_t^i = -\frac{\partial}{\partial t} \log P^i(0, t)$ is the market instantaneous forward rate at time 0 for a maturity t and $P^i(0, t)$ is the market zero-coupon bond price at time 0 for a maturity t .

Suppose that the LV model (2.7) has been calibrated and that σ_{LV} has been determined. The main building block of the calibration routine is expressing the leverage function σ in terms of the local volatility function σ_{LV} . We state the necessary and sufficient condition for a perfect calibration to vanilla options in the following theorem, proved in [11].

Theorem 2.1 (see [11, Theorem 1]). *Suppose that assumptions (A1) and (A3) hold and that both the marginal density of S_T in (2.1) and $\mathbb{E}[D_T^d v_T | S_T = \cdot]$ are continuous. Then the call price under model (2.1) matches the call price C_{LV} under model (2.7) for any maturity up to T^* and any strike if and only if, for all $T, K > 0$,*

$$(2.8) \quad \sigma^2(T, K) = \frac{\mathbb{E}[D_T^d | S_T = K]}{\mathbb{E}[D_T^d v_T | S_T = K]} \left\{ \sigma_{LV}^2(T, K) + \frac{2}{K^2 \frac{\partial^2 C_{LV}}{\partial K^2}} \left(\mathbb{E}[D_T^d (r_T^f - f_T^f)(S_T - K)^+] \right) \right. \\ \left. - K \mathbb{E}[D_T^d (r_T^d - f_T^d) \mathbf{1}_{S_T \geq K}] + K \mathbb{E}[D_T^d (r_T^f - f_T^f) \mathbf{1}_{S_T \geq K}] \right\},$$

where all expectations are under the domestic risk-neutral measure, σ_{LV} is a local volatility as in (2.7), and, denoting $\varphi = 2 + \sqrt{2}$ and $\zeta = \xi \sigma_{max}$, T^* is as given below.

(1) When $k < \varphi \zeta$,

$$(2.9) \quad T^* = \frac{2}{\sqrt{\varphi^2 \zeta^2 - k^2}} \left[\frac{\pi}{2} + \arctan \left(\frac{k}{\sqrt{\varphi^2 \zeta^2 - k^2}} \right) \right].$$

(2) When $k \geq \varphi \zeta$,

$$(2.10) \quad T^* = \infty.$$

This result generalizes the formula in [21] to a stochastic foreign short rate. Deelstra and Rayee [13] obtained a similar formula for a 4-factor SLV model with Hull–White short rate processes. An important step in the derivation is that the stochastic integral

$$(2.11) \quad \int_0^T \mathbf{1}_{S_t \geq K} \sigma(t, S_t) \sqrt{v_t} D_t^d S_t dW_t^s$$

is a true martingale. On the one hand, T^* is a lower bound on the explosion time of the second moment of the discounted spot process $D_t^d S_t$ (see Proposition 3.12). On the other hand, moments of the Heston model can explode in finite time [5], a property that is inherited by the Heston-type model (2.1). Note also Theorems 3.1 and 3.6 for possible moment explosions of the numerical approximations. Therefore, the formula (2.8) may not hold for some values of the model parameters and for large maturities T . However, in practice, T^* is very large. For instance, following our calibration routine to EURUSD market data from [11], we found $T^* = 28.6$.

As an aside, note that if the volatility in model (2.1) is purely local, i.e., if $v = 1$, we recover the formula in [9], whereas if the two short rates are deterministic, i.e., if $g^d = g^f = 0$, we recover the formula in [21, 40], i.e.,

$$(2.12) \quad \sigma(T, K) = \frac{\sigma_{LV}(T, K)}{\sqrt{\mathbb{E}[v_T | S_T = K]}}.$$

In the computations in [section 4](#), we assume a simple correlation structure where only ρ_{sv} may be nonzero. We use a five-step calibration routine which is explained in detail in [\[11\]](#). First, we calibrate the LV model [\(2.7\)](#) and find σ_{LV} . We also calibrate the associated 2-factor SLV model with deterministic rates via PDE methods, using formula [\(2.12\)](#) and results from the literature [\[9, 40\]](#). Independently, we calibrate the two CIR++ processes under their respective markets. Then, we find the Heston parameters by calibrating the associated 4-factor SV model, i.e., the Heston–2CIR++ SV model, using the closed-form solution for the call price in [\[1\]](#), extended to CIR++ processes, with other parameters from the above calibration. Finally, we calibrate σ in the Heston–2CIR++ SLV model [\(2.1\)](#) by working with the particle method of [\[21\]](#) and extending it using the calibrated 2-factor SLV model with deterministic rates as a control variate. For a partial correlation structure with possibly nonzero ρ_{sd} and ρ_{sf} , we may use the historical values of the correlations between the spot FX rate and the short-term zero-coupon bonds, and extend the approximation formulae of [\[19\]](#) for the call price from a 3-factor Heston–CIR model to a 4-factor Heston–2CIR++ model.

3. Convergence analysis.

3.1. The simulation scheme. We employ the full truncation Euler (FTE) scheme from [\[37\]](#) to discretize the variance and the two short rate processes. Consider the CIR process

$$(3.1) \quad dy_t = k_y(\theta_y - y_t)dt + \xi_y \sqrt{y_t} dW_t^y.$$

Let $N \in \mathbb{N}$ be a multiple of N_T and create an evenly spaced grid

$$(3.2) \quad T = N\delta t, \quad t_n = n\delta t \quad \forall n \in \{0, 1, \dots, N\}.$$

First, we introduce the discrete-time auxiliary process

$$(3.3) \quad \tilde{y}_{t_{n+1}} = \tilde{y}_{t_n} + k_y(\theta_y - \tilde{y}_{t_n}^+) \delta t + \xi_y \sqrt{\tilde{y}_{t_n}^+} \delta W_{t_n}^y,$$

where $y^+ = \max(0, y)$ and $\delta W_{t_n}^y = W_{t_{n+1}}^y - W_{t_n}^y$, and its continuous-time interpolation

$$(3.4) \quad \tilde{y}_t = \tilde{y}_{t_n} + k_y(\theta_y - \tilde{y}_{t_n}^+)(t - t_n) + \xi_y \sqrt{\tilde{y}_{t_n}^+} (W_t^y - W_{t_n}^y),$$

for any $t \in [t_n, t_{n+1})$, as suggested in [\[23\]](#). Then, we define the nonnegative processes

$$(3.5) \quad \begin{cases} \hat{y}_t = \tilde{y}_t^+, \\ \bar{y}_t = \tilde{y}_{t_n}^+ \end{cases}$$

$$(3.6)$$

whenever $t \in [t_n, t_{n+1})$. Let \bar{v} and \bar{g}^d be the FTE discretizations of v and g^d , respectively. Taking into account the presence of the quanto correction term in the drift of the foreign short rate, we similarly define

$$(3.7) \quad \begin{aligned} \tilde{g}_t^f &= \tilde{g}_{t_n}^f + \left[k_f \theta_f - k_f (\tilde{g}_{t_n}^f)^+ - \rho_{sf} \xi_f \sigma(t_n, \bar{S}_{t_n}) \sqrt{\tilde{v}_{t_n}^+ (\tilde{g}_{t_n}^f)^+} \right] (t - t_n) \\ &+ \xi_f \sqrt{(\tilde{g}_{t_n}^f)^+} (W_t^f - W_{t_n}^f), \end{aligned}$$

where \bar{S} is the continuous-time approximation of S defined below, as well as

$$(3.8) \quad \begin{cases} \hat{g}_t^f = (\tilde{g}_t^f)^+, \\ \bar{g}_t^f = (\tilde{g}_{t_n}^f)^+ \end{cases}$$

whenever $t \in [t_n, t_{n+1})$. For $i \in \{d, f\}$, the domestic and foreign short rate discretizations are

$$(3.10) \quad \bar{r}_t^i = \bar{g}_t^i + h_i(t).$$

Finally, we use an Euler–Maruyama scheme to discretize the log-exchange rate. Let x and \bar{x} be the actual and the approximated log-processes, and let $\bar{S} = e^{\bar{x}}$ be the continuous-time approximation of S . For brevity, define $h = h_d - h_f$. Then the discrete method reads

$$(3.11) \quad \bar{x}_{t_{n+1}} = \bar{x}_{t_n} + \int_{t_n}^{t_{n+1}} h(u) du + \left(\bar{g}_{t_n}^d - \bar{g}_{t_n}^f - \frac{1}{2} \sigma^2(t_n, \bar{S}_{t_n}) \bar{v}_{t_n} \right) \delta t + \sigma(t_n, \bar{S}_{t_n}) \sqrt{\bar{v}_{t_n}} \delta W_{t_n}^s.$$

However, we find it convenient to work with the continuous-time approximation

$$(3.12) \quad \bar{x}_t = \bar{x}_{t_n} + \int_{t_n}^t h(u) du + \left(\bar{g}_t^d - \bar{g}_t^f - \frac{1}{2} \bar{\sigma}^2(t, \bar{S}_t) \bar{v}_t \right) (t - t_n) + \bar{\sigma}(t, \bar{S}_t) \sqrt{\bar{v}_t} \Delta W_t^s,$$

where $\Delta W_t^s = W_t^s - W_{t_n}^s$ and $\bar{\sigma}(t, \bar{S}_t) = \sigma(t_n, \bar{S}_{t_n})$ whenever $t \in [t_n, t_{n+1})$. Hence,

$$(3.13) \quad \bar{x}_t = x_0 + \int_0^t \left(\bar{r}_u^d - \bar{r}_u^f - \frac{1}{2} \bar{\sigma}^2(u, \bar{S}_u) \bar{v}_u \right) du + \int_0^t \bar{\sigma}(u, \bar{S}_u) \sqrt{\bar{v}_u} dW_u^s.$$

Note that the convergence of the continuous-time approximation ensures that the discrete method approximates the true solution accurately at the gridpoints. Using Itô's formula, we obtain

$$(3.14) \quad \bar{S}_t = S_0 + \int_0^t (\bar{r}_u^d - \bar{r}_u^f) \bar{S}_u du + \int_0^t \bar{\sigma}(u, \bar{S}_u) \sqrt{\bar{v}_u} \bar{S}_u dW_u^s.$$

Furthermore, we define \hat{S} to be the piecewise constant continuous-time approximation of S , i.e.,

$$(3.15) \quad \hat{S}_t = \bar{S}_{t_n}$$

whenever $t \in [t_n, t_{n+1})$, an approximation which will be of use when pricing path-dependent options. We prefer the log-Euler scheme to the standard Euler scheme to discretize the exchange rate process because the former preserves positivity and produces no discretization bias in the S -direction when σ is constant. Moreover, if v , g^d , and g^f are also constant, then the log-Euler scheme is exact.

3.2. The main theorem. Define the arbitrage-free price of an option and its approximation under (3.15),

$$(3.16) \quad U = \mathbb{E} \left[e^{-\int_0^T r_t^d dt} f(S) \right],$$

$$(3.17) \quad \bar{U} = \mathbb{E} \left[e^{-\int_0^T \bar{r}_t^d dt} f(\hat{S}) \right].$$

The payoff function f may depend on the entire path of the underlying process and the expectation is under the domestic risk-neutral measure. The following theorem combines the statements of Theorems 3.15–3.20, and the proof is postponed to subsection 3.5.

Theorem 3.1. *If $2k_f\theta_f > \xi_f^2$, and under assumptions (A1)–(A3), the following statements hold.*

- (1) *The approximations to the values of the European put, the up-and-out barrier call, and any barrier put option defined in (3.17) converge as $\delta t \rightarrow 0$.*
- (2) *Denote $\zeta = \xi\sigma_{max}$, with σ_{max} from (2.3). If*

$$(3.18) \quad T < T^* \equiv \frac{4k}{\zeta^2} \mathbb{1}_{\zeta < 2k} + \frac{1}{\zeta - k} \mathbb{1}_{\zeta \geq 2k},$$

then the approximations to the values of the European call, Asian options, the down-and-in/out, and the up-and-in barrier call option defined in (3.17) converge as $\delta t \rightarrow 0$.

Remark 3.2. If assumption (A1) holds, then for the purpose of this paper we choose the smallest upper bound on the leverage function, namely

$$(3.19) \quad \sigma_{max} = \sup \{ \sigma(t, x) \mid t \in [0, T], x \in [0, \infty) \}.$$

Remark 3.3. If the exchange rate and the foreign short rate dynamics are independent of each other, i.e., if $\rho_{sf} = 0$, then the quanto correction term in the drift of the foreign short rate vanishes and Theorem 3.1 holds independently of the condition $2k_f\theta_f > \xi_f^2$, albeit with a simpler proof. If the domestic and foreign short rates are constant throughout the lifetime of the option and if, moreover, $\sigma(t, x) = 1$ for all $t \in [0, T]$ and $x \in [0, \infty)$, then the system of equations (2.1) collapses to the Heston model and Theorem 3.1 applies with $\zeta = \xi$. This extends the convergence results of Higham and Mao [23] to options with unbounded payoffs.

Remark 3.4. Tian et al. [42] calibrated the term-structure Heston SLV model parameters to the market implied volatility data on EURUSD using 10 term structures and maturities up to 5 years. The data in Table 3.1 suggest that the condition in (3.18) is typically satisfied in practice, both for shorter and longer maturities.

Table 3.1

The calibrated Heston SLV parameters for EURUSD market data from August 23, 2012 [42].

T	k	ξ	σ_{max}	ζ	T^*
1 month	0.885	0.342	1.600	0.547	11.8 years
5 years	0.978	0.499	1.300	0.649	9.3 years

In [11], the Heston–2CIR++ SLV model is calibrated to the EURUSD market data from March 18, 2016, for maturities up to 5 years. The data in Table 3.2 indicate that the condition in (3.18) is satisfied even for very long maturities. Furthermore, the following parameter values for the foreign short rate process are recovered: $k_f = 0.011$, $\theta_f = 1.166$, $\xi_f = 0.037$, so that the other condition also holds, i.e., $2k_f\theta_f = 0.0257 \gg 0.0014 = \xi_f^2$.

Table 3.2

The calibrated Heston–2CIR++ SLV parameters for EURUSD market data from March 18, 2016 [11].

T	k	ξ	σ_{max}	ζ	T^*
5 years	1.412	0.299	1.399	0.418	32.3 years

In equity markets, the mean-reversion speed is usually several times greater than the volatility of volatility. For instance, Hurn, Lindsay, and McClelland [26] calibrated the Heston model for the S&P 500 index from January 1990 to December 2011 using a combination of two out-of-the-money options and found that $k = 1.977 \gg 0.456 = \xi$. Furthermore, we typically have $\zeta < 1$, so that the condition in (3.18) holds even for longer maturities.

3.3. The square-root process. In order to prove the convergence of the approximation scheme in (3.14), we first need to examine the stability of the moments of order higher than 1 of the actual and the discretized processes. However, this problem is directly related to the exponential integrability of the CIR process and its approximation. Let y be the CIR process in (3.1) and \bar{y} be the piecewise constant FTE interpolant as per (3.6).

Proposition 3.5. Let $\lambda > 0$ and define the stochastic process

$$(3.20) \quad \Theta_t \equiv \exp \left\{ \lambda \int_0^t y_u du \right\} \quad \forall t \geq 0.$$

If $T < T^*$, then the first moment of Θ_T is bounded, i.e.,

$$(3.21) \quad \mathbb{E} [\Theta_T] < \infty,$$

where T^* is as given below.

(1) When $k_y^2 < 2\lambda\xi_y^2$,

$$(3.22) \quad T^* = \frac{2}{\sqrt{2\lambda\xi_y^2 - k_y^2}} \left[\frac{\pi}{2} + \arctan \left(\frac{k_y}{\sqrt{2\lambda\xi_y^2 - k_y^2}} \right) \right].$$

(2) When $k_y^2 \geq 2\lambda\xi_y^2$,

$$(3.23) \quad T^* = \infty.$$

Proof. The proof follows directly from Proposition 3.1 in [5]. ■

Theorem 3.6. Let $\lambda > 0$ and define the stochastic process

$$(3.24) \quad \bar{\Theta}_t \equiv \exp \left\{ \lambda \int_0^t \bar{y}_u du \right\} \quad \forall t \geq 0.$$

If $T \leq T^*$ and $\delta_T < k_y^{-1}$, then the first moment of $\bar{\Theta}_T$ is uniformly bounded, i.e.,

$$(3.25) \quad \sup_{\delta t \in (0, \delta_T)} \mathbb{E} [\bar{\Theta}_T] < \infty,$$

where T^* is as given below.

(1) When $2k_y^2 \leq \lambda\xi_y^2$,

$$(3.26) \quad T^* = \frac{1}{\sqrt{2\lambda\xi_y} - k_y}.$$

(2) When $2k_y^2 > \lambda\xi_y^2$,

$$(3.27) \quad T^* = \frac{2k_y}{\lambda\xi_y^2}.$$

Proof. First, we prove that there exists $\eta \geq 1$ independent of δt such that, for all $\omega \in [0, 1]$,

$$(3.28) \quad \eta^2 \omega^2 \lambda \xi_y^2 T^2 - 2\eta \omega k_y T - 2\eta + 2 \leq 0.$$

Fix any $\eta \geq 1$ and let $f_\eta : [0, 1] \rightarrow \mathbb{R}$ be defined by

$$(3.29) \quad f_\eta(\omega) = \omega^2 \eta^2 \lambda \xi_y^2 T^2 - 2\omega \eta k_y T - 2(\eta - 1),$$

with two distinct real roots

$$(3.30) \quad \omega_\pm = \frac{k_y \pm \sqrt{k_y^2 + 2(\eta - 1)\lambda\xi_y^2}}{\eta\lambda\xi_y^2 T}.$$

Since $\omega_- \leq 0 < \omega_+$, we have that $f_\eta([0, 1]) \subseteq (-\infty, 0]$ if and only if $f_\eta(1) \leq 0$, i.e.,

$$(3.31) \quad \eta^2 \lambda \xi_y^2 T^2 - 2\eta(1 + k_y T) + 2 \leq 0.$$

This holds for some $\eta \geq 1$ if and only if the quadratic polynomial in η on the left-hand side has a real root greater than or equal to one. Hence, we find the necessary and sufficient conditions

$$(3.32) \quad (\sqrt{2\lambda\xi_y} - k_y)T \leq 1$$

and

$$(3.33) \quad 2\lambda\xi_y^2 T \leq k_y + \sqrt{k_y^2 + 4\lambda\xi_y^2} \quad \text{or} \quad k_y + \sqrt{k_y^2 + 4\lambda\xi_y^2} < 2\lambda\xi_y^2 T \leq 4k_y,$$

which are equivalent to $T \leq T^*$, with T^* defined in (3.26)–(3.27). Fix any η satisfying (3.28).

Next, we prove by induction on $0 \leq m \leq N$ that, for all $\delta t < \delta_T$,

$$(3.34) \quad \mathbb{E} [\bar{\Theta}_T] \leq \exp \left\{ \frac{\eta}{2} (k_y \theta_y + \nu_y \xi_y) \lambda (\delta t)^2 (m-1)m \right\} \\ \times \mathbb{E} \left[\exp \left\{ \lambda \delta t \sum_{i=0}^{N-m-1} \bar{y}_{t_i} + \eta m \lambda \delta t \bar{y}_{t_{N-m}} \right\} \right],$$

where

$$(3.35) \quad \nu_y = \sqrt{\frac{\xi_y^2}{4\pi(1-k_y\delta_T)^2} + \frac{1}{2\pi} \sqrt{\frac{\xi_y^4}{4(1-k_y\delta_T)^4} + \frac{k_y^2\theta_y^2}{(1-k_y\delta_T)^2}}.$$

Let $\{\mathcal{G}_t^y, 0 \leq t \leq T\}$ be the natural filtration generated by the Brownian motion W^y and consider the shorthand notation $\mathbb{E}_t^y[\cdot] = \mathbb{E}[\cdot | \mathcal{G}_t^y]$ for the conditional expectation. Note that (3.34) clearly holds when $m \in \{0, 1\}$. Next, let us assume that (3.34) holds for $1 \leq m < N$ and prove the inductive step. Conditioning on the σ -algebra $\mathcal{G}_{t_{N-m-1}}^y$, we obtain

$$(3.36) \quad \begin{aligned} \mathbb{E}[\bar{\Theta}_T] &\leq \exp\left\{\frac{1}{2}\eta(k_y\theta_y + \nu_y\xi_y)\lambda(\delta t)^2(m-1)m\right\} \\ &\times \mathbb{E}\left[\exp\left\{\lambda\delta t \sum_{i=0}^{N-m-1} \bar{y}_{t_i}\right\} \mathbb{E}_{t_{N-m-1}}^y\left[\exp\left\{\eta m\lambda\delta t \bar{y}_{t_{N-m}}\right\}\right]\right]. \end{aligned}$$

For convenience, define $w = \bar{y}_{t_{N-m-1}}$. If $Z \sim \mathcal{N}(0, 1)$ independent of $\mathcal{G}_{t_{N-m-1}}^y$, then we have $\delta W_{t_{N-m-1}}^y \stackrel{\text{law}}{=} \sqrt{\delta t}Z$. Let \mathcal{I} be the inner expectation in (3.36); then

$$(3.37) \quad \mathcal{I} \leq \mathbb{E}_{0,w}\left[\exp\left\{\eta m\lambda\delta t \max\left[0, w + k_y(\theta_y - w)\delta t + \xi_y\sqrt{w\delta t}Z\right]\right\}\right].$$

There are two possible outcomes, namely $w = 0$, in which case

$$(3.38) \quad \mathcal{I} \leq \exp\left\{\eta m k_y \theta_y \lambda (\delta t)^2\right\},$$

and $w > 0$, which is treated now:

$$(3.39) \quad \begin{aligned} \mathcal{I} &\leq \int_{z_0}^{\infty} \frac{1}{\sqrt{2\pi}} \exp\left\{-\frac{1}{2}z^2 + \eta m \xi_y \lambda \sqrt{w} (\delta t)^{3/2} z + \eta m \lambda \delta t [w + k_y(\theta_y - w)\delta t]\right\} dz \\ &+ \int_{-\infty}^{z_0} \frac{1}{\sqrt{2\pi}} \exp\left\{-\frac{1}{2}z^2\right\} dz, \end{aligned}$$

where

$$(3.40) \quad z_0 = -\frac{k_y\theta_y\delta t + (1-k_y\delta t)w}{\xi_y\sqrt{w\delta t}}.$$

Recall that $\delta_T < k_y^{-1}$ and define

$$(3.41) \quad z_1 = z_0 - \eta m \xi_y \lambda \sqrt{w} (\delta t)^{3/2}.$$

Then $z_1 < z_0 < 0$. If ϕ and Φ are the standard normal probability density function (PDF) and cumulative distribution function (CDF), then

$$(3.42) \quad \mathcal{I} \leq \Phi(z_0) + \exp\left\{\eta m \lambda \delta t [k_y\theta_y\delta t + (1-k_y\delta t)w] + \frac{1}{2}\eta^2 m^2 \xi_y^2 \lambda^2 w (\delta t)^3\right\} \{1 - \Phi(z_1)\},$$

and hence

$$(3.43) \quad \mathcal{I} \leq \exp \left\{ \eta m k_y \theta_y \lambda (\delta t)^2 + \frac{1}{2} a \lambda w \delta t \right\} \left\{ 1 + \Phi(z_0) - \Phi(z_1) \right\},$$

where

$$(3.44) \quad a = 2\eta m(1 - k_y \delta t) + \eta^2 m^2 \xi_y^2 \lambda (\delta t)^2 > 0.$$

Applying the mean value (MV) theorem to $\Phi \in C^1$, we can find $z \in [z_1, z_0]$ such that

$$(3.45) \quad \Phi(z_0) - \Phi(z_1) = (z_0 - z_1) \phi(z) \leq (z_0 - z_1) \phi(z_0).$$

Hence, from (3.40) and (3.41),

$$(3.46) \quad \Phi(z_0) - \Phi(z_1) \leq \underbrace{\frac{1}{\sqrt{2\pi}} \eta m \xi_y \lambda (\delta t)^{3/2}}_{= b, \text{ constant w.r.t. } w} \cdot \sqrt{w} \exp \left\{ -\frac{[k_y \theta_y \delta t + (1 - k_y \delta t)w]^2}{2\xi_y^2 w \delta t} \right\}.$$

We can think of the right-hand side as a function of w , say $f : (0, \infty) \rightarrow \mathbb{R}$. Next, we show that

$$(3.47) \quad f(w) \leq \eta m \nu_y \xi_y \lambda (\delta t)^2 \quad \forall w > 0.$$

In order to find its global maximum, we need to compute the first derivative.

$$(3.48) \quad f'(w) = b e^{-z_0^2/2} \left\{ \frac{1}{2\sqrt{w}} + \sqrt{w} \left[\frac{k_y^2 \theta_y^2 \delta t}{2\xi_y^2 w^2} - \frac{(1 - k_y \delta t)^2}{2\xi_y^2 \delta t} \right] \right\}.$$

Therefore,

$$(3.49) \quad f'(w) = 0 \Leftrightarrow -(1 - k_y \delta t)^2 w^2 + \xi_y^2 w \delta t + k_y^2 \theta_y^2 (\delta t)^2 = 0.$$

To solve this quadratic, we divide throughout by $(\delta t)^2$ and introduce a new variable, $\hat{w} = w/\delta t$. Then there exists a unique, positive solution \hat{w}_0 for which the derivative is zero, namely

$$(3.50) \quad \hat{w}_0 = \frac{\xi_y^2}{2(1 - k_y \delta t)^2} + \sqrt{\frac{\xi_y^4}{4(1 - k_y \delta t)^4} + \frac{k_y^2 \theta_y^2}{(1 - k_y \delta t)^2}}.$$

From (3.35), we have that

$$(3.51) \quad \hat{w}_0 < 2\pi \nu_y^2 \Rightarrow w_0 < 2\pi \nu_y^2 \delta t.$$

Since the second root is negative, $f(w)$ must be increasing up to w_0 and decreasing after this point, so the function attains its global maximum at w_0 . Hence,

$$(3.52) \quad f(w) \leq f(w_0) = b \sqrt{w_0} e^{-z_0^2/2} \leq b \sqrt{2\pi \nu_y^2 \delta t} = \eta m \nu_y \xi_y \lambda (\delta t)^2.$$

Making use of the upper bound in (3.47), we derive the following inequality:

$$(3.53) \quad 1 + \Phi(z_0) - \Phi(z_1) \leq \exp \left\{ \eta m \nu_y \xi_y \lambda (\delta t)^2 \right\}.$$

Substituting back into (3.43) with (3.53), we get

$$(3.54) \quad \mathcal{I} \leq \exp \left\{ \eta m (k_y \theta_y + \nu_y \xi_y) \lambda (\delta t)^2 + \frac{1}{2} a \lambda w \delta t \right\}.$$

Note from (3.38) that this holds when $w = 0$ as well. Applying (3.28) with $\omega = \frac{m}{N}$ leads to

$$(3.55) \quad \eta^2 m^2 \xi_y^2 \lambda (\delta t)^2 - 2 \eta m k_y \delta t - 2 \eta + 2 \leq 0.$$

Hence, from (3.44),

$$(3.56) \quad a \leq 2 \eta (m + 1) - 2.$$

Therefore,

$$(3.57) \quad \mathcal{I} \leq \exp \left\{ \eta m (k_y \theta_y + \nu_y \xi_y) \lambda (\delta t)^2 - \lambda w \delta t + \eta (m + 1) \lambda w \delta t \right\}.$$

Substituting back into (3.36) gives the inductive step. Finally, taking $m = N$ in (3.34) leads to

$$(3.58) \quad \mathbb{E} [\bar{\Theta}_T] < \exp \left\{ \frac{1}{2} \eta \lambda T^2 (k_y \theta_y + \nu_y \xi_y) + \eta \lambda T y_0 \right\}.$$

The right-hand side is finite and independent of δt , and the conclusion follows. \blacksquare

The next two results establish uniform moment bounds for both the original and the discretized variance and short rate processes. Note that under assumption (A3), the moment boundedness of g^i and \hat{g}^i , for $i \in \{d, f\}$, extends naturally to $|r^i|$ and $|\bar{r}^i|$.

Proposition 3.7. *The following two statements hold.*

(1) *The square-root process in (3.1) has uniformly bounded moments, i.e.,*

$$(3.59) \quad \mathbb{E} \left[\sup_{t \in [0, T]} y_t^p \right] < \infty \quad \forall p \geq 1.$$

(2) *The FTE scheme in (3.5) for the square-root process has uniformly bounded moments, i.e.,*

$$(3.60) \quad \sup_{\delta t \in (0, \delta_T)} \mathbb{E} \left[\sup_{t \in [0, T]} \hat{y}_t^p \right] < \infty \quad \forall p \geq 1, \quad \forall \delta_T > 0.$$

Proof. (1) The polynomial moments of the square-root process can be expressed in terms of the confluent hypergeometric function and, according to [14] or to Theorem 3.1 in [25], they are uniformly bounded, i.e.,

$$(3.61) \quad \sup_{t \in [0, T]} \mathbb{E} [y_t^p] < \infty \quad \forall p > -\frac{2k_y \theta_y}{\xi_y^2}.$$

On the other hand, using Hölder's inequality, we deduce that

$$(3.62) \quad \sup_{t \in [0, T]} y_t^p \leq 3^{p-1} (y_0 + k_y \theta_y T)^p + 3^{p-1} k_y^p T^{p-1} \int_0^T y_u^p du + 3^{p-1} \xi_y^p \sup_{t \in [0, T]} \left| \int_0^t \sqrt{y_u} dW_u^y \right|^p.$$

Using the Burkholder–Davis–Gundy (BDG) inequality, we can find a constant $C_p > 0$ such that

$$(3.63) \quad \begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} \left| \int_0^t \sqrt{y_u} dW_u^y \right|^p \right] &\leq C_p \mathbb{E} \left[\left(\int_0^T y_u du \right)^{p/2} \right] \\ &\leq \frac{1}{2} C_p + \frac{1}{2} C_p T^{p-1} \mathbb{E} \left[\int_0^T y_u^p du \right]. \end{aligned}$$

Taking expectations in (3.62) and employing Fubini's theorem, we get

$$(3.64) \quad \mathbb{E} \left[\sup_{t \in [0, T]} y_t^p \right] \leq 3^{p-1} (y_0 + k_y \theta_y T)^p + \frac{1}{2} 3^{p-1} \xi_y^p C_p + 3^{p-1} \left(k_y^p + \frac{1}{2} \xi_y^p C_p \right) T^p \sup_{t \in [0, T]} \mathbb{E} [y_t^p].$$

The right-hand side is finite by (3.61), whence the conclusion.

(2) Integrating the auxiliary process \tilde{y} defined in (3.4), we deduce that

$$(3.65) \quad \tilde{y}_t = y_0 + k_y \int_0^t (\theta_y - \bar{y}_u) du + \xi_y \int_0^t \sqrt{\bar{y}_u} dW_u^y.$$

For any $p, \epsilon > 0$, there exists a constant $c(p, \epsilon) > 0$ such that $\max(0, x)^p \leq c(p, \epsilon) e^{\epsilon x}$ for all $x \in \mathbb{R}$. In particular, this implies that $\hat{y}_t^p \leq c(p, \epsilon) e^{\epsilon \tilde{y}_t}$ for all $t \in [0, T]$. Hence,

$$(3.66) \quad \sup_{t \in [0, T]} \mathbb{E} [\hat{y}_t^p] \leq c(p, \epsilon) e^{\epsilon(y_0 + k_y \theta_y T)} \sup_{t \in [0, T]} \mathbb{E} [\Gamma_t],$$

where

$$(3.67) \quad \Gamma_t \equiv \exp \left\{ -\epsilon k_y \int_0^t \bar{y}_u du + \epsilon \xi_y \int_0^t \sqrt{\bar{y}_u} dW_u^y \right\}.$$

Assuming that $t \in [t_n, t_{n+1}]$ and conditioning on the σ -algebra $\mathcal{G}_{t_n}^y$, we get

$$(3.68) \quad \mathbb{E}_{t_n}^y [\Gamma_t] = \Gamma_{t_n} \exp \left\{ \left(\frac{1}{2} \epsilon^2 \xi_y^2 - \epsilon k_y \right) (t - t_n) \bar{y}_{t_n} \right\}.$$

However, we can find ϵ sufficiently small such that $2k_y \geq \epsilon \xi_y^2$. Then $\mathbb{E}_{t_n}^y [\Gamma_t] \leq \mathbb{E}_{t_n}^y [\Gamma_{t_n}]$ and the law of iterated expectations ensures that

$$(3.69) \quad \mathbb{E} [\Gamma_t] \leq \mathbb{E} [\Gamma_{t_n}] \leq \mathbb{E} [\Gamma_{t_{n-1}}] \leq \dots \leq \mathbb{E} [\Gamma_0] \Rightarrow \sup_{t \in [0, T]} \mathbb{E} [\Gamma_t] = 1.$$

Substituting back into (3.66), we deduce that

$$(3.70) \quad \sup_{t \in [0, T]} \mathbb{E} [\hat{y}_t^p] \leq c(p, \epsilon) e^{\epsilon(y_0 + k_y \theta_y T)},$$

which is independent of δt . On the other hand, since $\hat{y}_t \leq |\tilde{y}_t|$ and using Hölder's inequality, we get

$$(3.71) \quad \sup_{t \in [0, T]} \hat{y}_t^p \leq 3^{p-1} (y_0 + k_y \theta_y T)^p + 3^{p-1} k_y^p T^{p-1} \int_0^T \bar{y}_u^p du + 3^{p-1} \xi_y^p \sup_{t \in [0, T]} \left| \int_0^t \sqrt{\bar{y}_u} dW_u^y \right|^p.$$

Taking expectations on both sides and using the BDG inequality as in (3.63) leads to

$$(3.72) \quad \mathbb{E} \left[\sup_{t \in [0, T]} \hat{y}_t^p \right] \leq 3^{p-1} (y_0 + k_y \theta_y T)^p + \frac{1}{2} 3^{p-1} \xi_y^p C_p + 3^{p-1} \left(k_y^p + \frac{1}{2} \xi_y^p C_p \right) T^p \sup_{t \in [0, T]} \mathbb{E} [\bar{y}_t^p]$$

for some constant $C_p > 0$. However, $\sup_{t \in [0, T]} \mathbb{E} [\bar{y}_t^p] \leq \sup_{t \in [0, T]} \mathbb{E} [\hat{y}_t^p]$, and the conclusion follows from (3.70). \blacksquare

Proposition 3.8. *Under assumption (A1), the following two statements hold.*

(1) *The process g^f has uniformly bounded moments, i.e.,*

$$(3.73) \quad \mathbb{E} \left[\sup_{t \in [0, T]} (g_t^f)^p \right] < \infty \quad \forall p \geq 1.$$

(2) *The process \hat{g}^f from (3.8) has uniformly bounded moments, i.e.,*

$$(3.74) \quad \sup_{\delta t \in (0, \delta_T)} \mathbb{E} \left[\sup_{t \in [0, T]} (\hat{g}_t^f)^p \right] < \infty \quad \forall p \geq 1, \quad \forall \delta_T > 0.$$

Proof. (1) From (2.1), we know that

$$(3.75) \quad g_t^f = g_0^f + k_f \theta_f t - k_f \int_0^t g_u^f du - \rho_{sf} \xi_f \int_0^t \sigma(u, S_u) \sqrt{v_u g_u^f} du + \xi_f \int_0^t \sqrt{g_u^f} dW_u^f.$$

Since g^f is nonnegative and using the fact that $2\sqrt{|ab|} \leq |a| + |b|$, we find an upper bound

$$(3.76) \quad \begin{aligned} g_t^f &\leq g_0^f + k_f \theta_f t + \frac{1}{2} |\rho_{sf}| \xi_f \sigma_{max} \int_0^t v_u du + \frac{1}{2} (2k_f + |\rho_{sf}| \xi_f \sigma_{max}) \int_0^t g_u^f du \\ &+ \xi_f \left| \int_0^t \sqrt{g_u^f} dW_u^f \right|. \end{aligned}$$

Fix $t \in [0, T]$. Using Hölder's inequality, we get

$$(3.77) \quad \begin{aligned} \sup_{s \in [0, t]} (g_s^f)^p &\leq 4^{p-1} (g_0^f + k_f \theta_f T)^p + 2^{p-2} |\rho_{sf}|^p \xi_f^p \sigma_{max}^p T^{p-1} \int_0^T v_u^p du \\ &+ 2^{p-2} (2k_f + |\rho_{sf}| \xi_f \sigma_{max})^p T^{p-1} \int_0^t (g_u^f)^p du \\ &+ 4^{p-1} \xi_f^p \sup_{s \in [0, t]} \left| \int_0^s \sqrt{g_u^f} dW_u^f \right|^p. \end{aligned}$$

Taking expectations on both sides and using the BDG inequality as in (3.63) leads to

$$(3.78) \quad \begin{aligned} \mathbb{E} \left[\sup_{s \in [0, t]} (g_s^f)^p \right] &\leq 4^{p-1} (g_0^f + k_f \theta_f T)^p + 2^{p-2} |\rho_{sf}|^p \xi_f^p \sigma_{max}^p T^p \sup_{u \in [0, T]} \mathbb{E} [v_u^p] \\ &\quad + 2^{2p-3} \xi_f^p C_p + \left(2^{p-2} (2k_f + |\rho_{sf}| \xi_f \sigma_{max})^p T^{p-1} \right. \\ &\quad \left. + 2^{2p-3} \xi_f^p C_p T^{p-1} \right) \int_0^t \mathbb{E} \left[\sup_{s \in [0, u]} (g_s^f)^p \right] du \end{aligned}$$

for some constant $C_p > 0$. Applying Gronwall's inequality (see [15]), we get

$$(3.79) \quad \begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} (g_t^f)^p \right] &\leq \left(4^{p-1} (g_0^f + k_f \theta_f T)^p + 2^{2p-3} \xi_f^p C_p + 2^{p-2} |\rho_{sf}|^p \xi_f^p \sigma_{max}^p T^p \sup_{u \in [0, T]} \mathbb{E} [v_u^p] \right) \\ &\quad \times \exp \left\{ 2^{p-2} (2k_f + |\rho_{sf}| \xi_f \sigma_{max})^p T^p + 2^{2p-3} \xi_f^p C_p T^p \right\}. \end{aligned}$$

The conclusion follows from Proposition 3.7.

(2) Integrating the auxiliary process \tilde{g}^f defined in (3.7), we deduce that

$$(3.80) \quad \tilde{g}_t^f = g_0^f + k_f \theta_f t - k_f \int_0^t \bar{g}_u^f du - \rho_{sf} \xi_f \int_0^t \bar{\sigma}(u, \bar{S}_u) \sqrt{\bar{v}_u \bar{g}_u^f} du + \xi_f \int_0^t \sqrt{\bar{g}_u^f} dW_u^f.$$

Since $\hat{g}_t^f \leq |\tilde{g}_t^f|$, following the previous argument, we deduce that

$$(3.81) \quad \begin{aligned} \mathbb{E} \left[\sup_{s \in [0, t]} (\hat{g}_s^f)^p \right] &\leq 4^{p-1} (g_0^f + k_f \theta_f T)^p + 2^{2p-3} \xi_f^p C_p + 2^{p-2} |\rho_{sf}|^p \xi_f^p \sigma_{max}^p T^p \sup_{u \in [0, T]} \mathbb{E} [\bar{v}_u^p] \\ &\quad + \left(2^{p-2} (2k_f + |\rho_{sf}| \xi_f \sigma_{max})^p T^{p-1} + 2^{2p-3} \xi_f^p C_p T^{p-1} \right) \int_0^t \mathbb{E} [(\bar{g}_u^f)^p] du. \end{aligned}$$

Since $\sup_{u \in [0, T]} \mathbb{E} [\bar{v}_u^p] \leq \sup_{u \in [0, T]} \mathbb{E} [\hat{v}_u^p]$ and $\bar{g}_u^f \leq \sup_{s \in [0, u]} \hat{g}_s^f$, using Gronwall's inequality, we get

$$(3.82) \quad \begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} (\hat{g}_t^f)^p \right] &\leq \left(4^{p-1} (g_0^f + k_f \theta_f T)^p + 2^{2p-3} \xi_f^p C_p + 2^{p-2} |\rho_{sf}|^p \xi_f^p \sigma_{max}^p T^p \sup_{u \in [0, T]} \mathbb{E} [\hat{v}_u^p] \right) \\ &\quad \times \exp \left\{ 2^{p-2} (2k_f + |\rho_{sf}| \xi_f \sigma_{max})^p T^p + 2^{2p-3} \xi_f^p C_p T^p \right\}. \end{aligned}$$

The conclusion follows from Proposition 3.7. ■

Next, we establish the strong mean square convergence of the discretized variance and domestic short rate processes. Unlike in [37], where the focus was on the continuous-time approximation \hat{y} , we are rather interested in the behavior of \bar{y} in the limit as the time step goes to zero. Note that the convergence of \bar{g}^d extends naturally to \bar{r}^d .

Proposition 3.9. *The full truncation scheme in (3.6) for the square-root process converges strongly in the L^2 sense, i.e.,*

$$(3.83) \quad \lim_{\delta t \rightarrow 0} \sup_{t \in [0, T]} \mathbb{E} [|y_t - \bar{y}_t|^2] = 0.$$

Proof. First, fix $t \in [0, T]$. Since $|y_t - \hat{y}_t| \leq |y_t - \bar{y}_t|$, using the Cauchy–Schwarz inequality, we get

$$(3.84) \quad \sup_{s \in [0, t]} |y_s - \hat{y}_s|^2 \leq 2k_y^2 t \int_0^t (y_u - \bar{y}_u)^2 du + 2\xi_y^2 \sup_{s \in [0, t]} \left| \int_0^s (\sqrt{y_u} - \sqrt{\bar{y}_u}) dW_u^y \right|^2.$$

Taking expectations and then using Doob’s martingale inequality and Fubini’s theorem leads to

$$(3.85) \quad \begin{aligned} \mathbb{E} \left[\sup_{s \in [0, t]} |y_s - \hat{y}_s|^2 \right] &\leq 4k_y^2 T \int_0^t \mathbb{E} \left[\sup_{s \in [0, u]} |y_s - \hat{y}_s|^2 \right] du + 4k_y^2 T^2 \sup_{u \in [0, T]} \mathbb{E} [|\hat{y}_u - \bar{y}_u|^2] \\ &+ 8\xi_y^2 T \sup_{u \in [0, T]} \mathbb{E} [|y_u - \hat{y}_u|] + 8\xi_y^2 T \sup_{u \in [0, T]} \mathbb{E} [|\hat{y}_u - \bar{y}_u|]. \end{aligned}$$

Applying Gronwall’s inequality, we get

$$(3.86) \quad \begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} |y_t - \hat{y}_t|^2 \right] &\leq e^{4k_y^2 T^2} \left(4k_y^2 T^2 \sup_{t \in [0, T]} \mathbb{E} [|\hat{y}_t - \bar{y}_t|^2] + 8\xi_y^2 T \sup_{t \in [0, T]} \mathbb{E} [|\hat{y}_t - \bar{y}_t|] \right. \\ &\left. + 8\xi_y^2 T \sup_{t \in [0, T]} \mathbb{E} [|y_t - \hat{y}_t|] \right). \end{aligned}$$

On the other hand, we know that

$$(3.87) \quad \sup_{t \in [0, T]} \mathbb{E} [|y_t - \bar{y}_t|^2] \leq 2 \sup_{t \in [0, T]} \mathbb{E} [|y_t - \hat{y}_t|^2] + 2 \sup_{t \in [0, T]} \mathbb{E} [|\hat{y}_t - \bar{y}_t|^2].$$

Substituting into (3.87) with the upper bound in (3.86), we get

$$(3.88) \quad \begin{aligned} \sup_{t \in [0, T]} \mathbb{E} [|y_t - \bar{y}_t|^2] &\leq 16\xi_y^2 T e^{4k_y^2 T^2} \sup_{t \in [0, T]} \mathbb{E} [|y_t - \hat{y}_t|] + 16\xi_y^2 T e^{4k_y^2 T^2} \sup_{t \in [0, T]} \mathbb{E} [|\hat{y}_t - \bar{y}_t|] \\ &+ 2 \left(1 + 4k_y^2 T^2 e^{4k_y^2 T^2} \right) \sup_{t \in [0, T]} \mathbb{E} [|\hat{y}_t - \bar{y}_t|^2]. \end{aligned}$$

The convergence of the three terms on the right-hand side of (3.88) is a consequence of Lemma A.3 and Theorem 4.2 in [37], which concludes the proof. \blacksquare

3.4. The four-dimensional system. Even though weak convergence is important when estimating expectations of payoffs, strong convergence plays a crucial role in multilevel Monte Carlo methods and may be required for complex path-dependent derivatives. First, we prove the uniform mean square convergence of the stopped discretized spot FX rate process in (3.14).

Proposition 3.10. *Let $L_s > S_0$, $L_v > v_0$, $L_d > g_0^d$, $L_f > g_0^f > \kappa^{-1}$ be constants and define the stopping time*

$$(3.89) \quad \tau = \inf \{ t \geq 0 : S_t \geq L_s \text{ or } v_t \geq L_v \text{ or } \hat{v}_t \geq L_v \text{ or } \hat{g}_t^d \geq L_d \text{ or } \hat{g}_t^f \geq L_f \text{ or } g_t^f \leq \kappa^{-1} \}.$$

Under assumptions (A1)–(A3), the stopped process converges uniformly in L^2 , i.e.,

$$(3.90) \quad \lim_{\delta t \rightarrow 0} \mathbb{E} \left[\sup_{t \in [0, T]} |S_{t \wedge \tau} - \bar{S}_{t \wedge \tau}|^2 \right] = 0.$$

Proof. The absolute difference between the original and the discretized stopped processes can be bounded from above as follows:

$$\begin{aligned}
|S_{t \wedge \tau} - \bar{S}_{t \wedge \tau}| &\leq \left| \int_0^{t \wedge \tau} h(u) (S_u - \bar{S}_u) du \right| + \left| \int_0^{t \wedge \tau} (g_u^d - \bar{g}_u^d) S_u du \right| \\
&+ \left| \int_0^{t \wedge \tau} \bar{g}_u^d (S_u - \bar{S}_u) du \right| + \left| \int_0^{t \wedge \tau} (g_u^f - \bar{g}_u^f) S_u du \right| \\
&+ \left| \int_0^{t \wedge \tau} \bar{g}_u^f (S_u - \bar{S}_u) du \right| + \left| \int_0^{t \wedge \tau} \bar{\sigma}(u, \bar{S}_u) \sqrt{\bar{v}_u} (S_u - \bar{S}_u) dW_u^s \right| \\
&+ \left| \int_0^{t \wedge \tau} (\sigma(u, S_u) - \bar{\sigma}(u, \bar{S}_u)) \sqrt{\bar{v}_u} S_u dW_u^s \right| \\
(3.91) \quad &+ \left| \int_0^{t \wedge \tau} \sigma(u, S_u) (\sqrt{v_u} - \sqrt{\bar{v}_u}) S_u dW_u^s \right|.
\end{aligned}$$

Squaring both sides, taking the supremum over all $t \in [0, s]$, where $0 \leq s \leq T$, and then employing the Cauchy–Schwarz inequality leads to the upper bound

$$\begin{aligned}
\sup_{t \in [0, s]} |S_{t \wedge \tau} - \bar{S}_{t \wedge \tau}|^2 &\leq 8(L_d^2 + L_f^2 + 4h_{max}^2)T \int_0^{s \wedge \tau} (S_u - \bar{S}_u)^2 du + 8L_s^2 T \int_0^T (g_u^d - \bar{g}_u^d)^2 du \\
&+ 16L_s^2 T \int_0^{s \wedge \tau} (g_u^f - \hat{g}_u^f)^2 du + 16L_s^2 T \int_0^T (\hat{g}_u^f - \bar{g}_u^f)^2 du \\
&+ 8 \sup_{t \in [0, s]} \left| \int_0^{t \wedge \tau} \sigma(u, S_u) (\sqrt{v_u} - \sqrt{\bar{v}_u}) S_u dW_u^s \right|^2 \\
&+ 8 \sup_{t \in [0, s]} \left| \int_0^{t \wedge \tau} \bar{\sigma}(u, \bar{S}_u) \sqrt{\bar{v}_u} (S_u - \bar{S}_u) dW_u^s \right|^2 \\
(3.92) \quad &+ 8 \sup_{t \in [0, s]} \left| \int_0^{t \wedge \tau} (\sigma(u, S_u) - \bar{\sigma}(u, \bar{S}_u)) \sqrt{\bar{v}_u} S_u dW_u^s \right|^2.
\end{aligned}$$

Note that

$$(3.93) \quad \int_0^{s \wedge \tau} (g_u^f - \hat{g}_u^f)^2 du \leq \int_0^s (g_{u \wedge \tau}^f - \hat{g}_{u \wedge \tau}^f)^2 du \leq T \sup_{t \in [0, s]} |g_{t \wedge \tau}^f - \hat{g}_{t \wedge \tau}^f|^2.$$

Hence, taking expectations in (3.92), using Fubini's theorem, Doob's martingale inequality, and the Itô isometry, and upon noticing that a stopped martingale is also a martingale, we obtain

$$\begin{aligned}
\mathbb{E} \left[\sup_{t \in [0, s]} |S_{t \wedge \tau} - \bar{S}_{t \wedge \tau}|^2 \right] &\leq 8L_s^2 T^2 \sup_{t \in [0, T]} \mathbb{E} \left[|g_t^d - \bar{g}_t^d|^2 \right] + 32\sigma_{max}^2 L_s^2 T \sup_{t \in [0, T]} \mathbb{E} \left[|v_t - \bar{v}_t| \right] \\
&+ \left[8(L_d^2 + L_f^2 + 4h_{max}^2)T + 32\sigma_{max}^2 L_v \right] \mathbb{E} \left[\int_0^{s \wedge \tau} |S_u - \bar{S}_u|^2 du \right]
\end{aligned}$$

$$(3.94) \quad \begin{aligned} & + 32L_s^2 L_v \mathbb{E} \left[\int_0^{s \wedge \tau} |\sigma(u, S_u) - \bar{\sigma}(u, \bar{S}_u)|^2 du \right] \\ & + 16L_s^2 T^2 \sup_{t \in [0, T]} \mathbb{E} \left[|\hat{g}_t^f - \bar{g}_t^f|^2 \right] + 16L_s^2 T^2 \mathbb{E} \left[\sup_{t \in [0, s]} |g_{t \wedge \tau}^f - \hat{g}_{t \wedge \tau}^f|^2 \right]. \end{aligned}$$

First, we bound the last expectation on the right-hand side of (3.94) from above. From (3.75) and (3.80), since $|g_{t \wedge \tau}^f - \hat{g}_{t \wedge \tau}^f| \leq |g_{t \wedge \tau}^f - \bar{g}_{t \wedge \tau}^f|$, we get

$$(3.95) \quad \begin{aligned} |g_{t \wedge \tau}^f - \hat{g}_{t \wedge \tau}^f| & \leq \left| -k_f \int_0^{t \wedge \tau} (g_u^f - \hat{g}_u^f) du - k_f \int_0^{t \wedge \tau} (\hat{g}_u^f - \bar{g}_u^f) du \right. \\ & + \xi_f \int_0^{t \wedge \tau} (\sqrt{g_u^f} - \sqrt{\hat{g}_u^f}) dW_u^f + \xi_f \int_0^{t \wedge \tau} (\sqrt{\hat{g}_u^f} - \sqrt{\bar{g}_u^f}) dW_u^f \\ & - \rho_{sf} \xi_f \int_0^{t \wedge \tau} \sigma(u, S_u) \sqrt{v_u} (\sqrt{g_u^f} - \sqrt{\hat{g}_u^f}) du \\ & - \rho_{sf} \xi_f \int_0^{t \wedge \tau} \sigma(u, S_u) \sqrt{v_u} (\sqrt{\hat{g}_u^f} - \sqrt{\bar{g}_u^f}) du \\ & - \rho_{sf} \xi_f \int_0^{t \wedge \tau} \sigma(u, S_u) \sqrt{\bar{g}_u^f} (\sqrt{v_u} - \sqrt{\bar{v}_u}) du \\ & \left. - \rho_{sf} \xi_f \int_0^{t \wedge \tau} \sqrt{\bar{v}_u \bar{g}_u^f} (\sigma(u, S_u) - \bar{\sigma}(u, \bar{S}_u)) du \right|. \end{aligned}$$

Squaring both sides, taking the supremum over all $t \in [0, s']$, where $0 \leq s' \leq s$, and employing the Cauchy–Schwarz inequality, and then taking expectations and using Doob’s martingale inequality and Fubini’s theorem, we deduce that

$$(3.96) \quad \begin{aligned} \mathbb{E} \left[\sup_{t \in [0, s']} |g_{t \wedge \tau}^f - \hat{g}_{t \wedge \tau}^f|^2 \right] & \leq 8k_f^2 T \int_0^{s'} \mathbb{E} \left[(g_u^f - \hat{g}_u^f)^2 \mathbf{1}_{u < \tau} \right] du + 8k_f^2 T \int_0^T \mathbb{E} \left[(\hat{g}_u^f - \bar{g}_u^f)^2 \right] du \\ & + 32\xi_f^2 \int_0^T \mathbb{E} \left[|\hat{g}_u^f - \bar{g}_u^f| \right] du + 8\rho_{sf}^2 \xi_f^2 \sigma_{max}^2 L_v T \int_0^T \mathbb{E} \left[|\hat{g}_u^f - \bar{g}_u^f| \right] du \\ & + 32\xi_f^2 \int_0^{s'} \mathbb{E} \left[(\sqrt{g_u^f} - \sqrt{\hat{g}_u^f})^2 \mathbf{1}_{u < \tau} \right] du \\ & + 8\rho_{sf}^2 \xi_f^2 \sigma_{max}^2 L_v T \int_0^{s'} \mathbb{E} \left[(\sqrt{g_u^f} - \sqrt{\hat{g}_u^f})^2 \mathbf{1}_{u < \tau} \right] du \\ & + 8\rho_{sf}^2 \xi_f^2 \sigma_{max}^2 L_f T \int_0^T \mathbb{E} \left[|v_u - \bar{v}_u| \right] du \\ & + 8\rho_{sf}^2 \xi_f^2 L_v L_f T \mathbb{E} \left[\int_0^{s' \wedge \tau} |\sigma(u, S_u) - \bar{\sigma}(u, \bar{S}_u)|^2 du \right]. \end{aligned}$$

However,

$$(3.97) \quad (\sqrt{g_u^f} - \sqrt{\hat{g}_u^f})^2 \mathbf{1}_{u < \tau} \leq (\sqrt{g_{u \wedge \tau}^f} - \sqrt{\hat{g}_{u \wedge \tau}^f})^2 \leq \kappa (g_{u \wedge \tau}^f - \hat{g}_{u \wedge \tau}^f)^2,$$

and hence, since $s' \leq s$,

$$\begin{aligned}
\mathbb{E} \left[\sup_{t \in [0, s']} |g_{t \wedge \tau}^f - \hat{g}_{t \wedge \tau}^f|^2 \right] &\leq 8(k_f^2 T + 4\xi_f^2 \kappa + \rho_{sf}^2 \xi_f^2 \sigma_{max}^2 L_v \kappa T) \int_0^{s'} \mathbb{E} \left[\sup_{t \in [0, u]} |g_{t \wedge \tau}^f - \hat{g}_{t \wedge \tau}^f|^2 \right] du \\
&\quad + 8\rho_{sf}^2 \xi_f^2 L_v L_f T \mathbb{E} \left[\int_0^{s \wedge \tau} |\sigma(u, S_u) - \bar{\sigma}(u, \bar{S}_u)|^2 du \right] \\
&\quad + (32\xi_f^2 T + 8\rho_{sf}^2 \xi_f^2 \sigma_{max}^2 L_v T^2) \sup_{t \in [0, T]} \mathbb{E} \left[|\hat{g}_t^f - \bar{g}_t^f| \right] \\
(3.98) \quad &\quad + 8k_f^2 T^2 \sup_{t \in [0, T]} \mathbb{E} \left[(\hat{g}_t^f - \bar{g}_t^f)^2 \right] + 8\rho_{sf}^2 \xi_f^2 \sigma_{max}^2 L_f T^2 \sup_{t \in [0, T]} \mathbb{E} \left[|v_t - \bar{v}_t| \right].
\end{aligned}$$

Therefore, applying Gronwall's inequality to (3.98), since $s \leq T$, we get

$$\begin{aligned}
\mathbb{E} \left[\sup_{t \in [0, s]} |g_{t \wedge \tau}^f - \hat{g}_{t \wedge \tau}^f|^2 \right] &\leq e^{\beta_1 T} \left\{ 8\rho_{sf}^2 \xi_f^2 L_v L_f T \mathbb{E} \left[\int_0^{s \wedge \tau} |\sigma(u, S_u) - \bar{\sigma}(u, \bar{S}_u)|^2 du \right] \right. \\
&\quad + 8\rho_{sf}^2 \xi_f^2 \sigma_{max}^2 L_f T^2 \sup_{t \in [0, T]} \mathbb{E} \left[|v_t - \bar{v}_t| \right] + 8k_f^2 T^2 \sup_{t \in [0, T]} \mathbb{E} \left[(\hat{g}_t^f - \bar{g}_t^f)^2 \right] \\
(3.99) \quad &\quad \left. + (32\xi_f^2 T + 8\rho_{sf}^2 \xi_f^2 \sigma_{max}^2 L_v T^2) \sup_{t \in [0, T]} \mathbb{E} \left[|\hat{g}_t^f - \bar{g}_t^f| \right] \right\},
\end{aligned}$$

where

$$(3.100) \quad \beta_1 = 8k_f^2 T + 32\xi_f^2 \kappa + 8\rho_{sf}^2 \xi_f^2 \sigma_{max}^2 L_v \kappa T.$$

Second, since N is a multiple of N_T , assumption (A2) on the leverage function ensures that

$$\begin{aligned}
\mathbb{E} \left[\int_0^{s \wedge \tau} |\sigma(u, S_u) - \bar{\sigma}(u, \bar{S}_u)|^2 du \right] &\leq 3A^2 T (\delta t)^{2\alpha} + 3B^2 \mathbb{E} \left[\int_0^T |S_t - \bar{S}_t|^2 \mathbf{1}_{t < \tau} dt \right] \\
(3.101) \quad &\quad + 3B^2 \mathbb{E} \left[\int_0^{s \wedge \tau} \sup_{t \in [0, u]} |S_t - \bar{S}_t|^2 du \right],
\end{aligned}$$

where, for convenience of notation, we defined

$$(3.102) \quad \bar{t} = \delta t \left\lfloor \frac{t}{\delta t} \right\rfloor.$$

Furthermore, we know that

$$(3.103) \quad \int_0^{s \wedge \tau} |S_u - \bar{S}_u|^2 du \leq \int_0^{s \wedge \tau} \sup_{t \in [0, u]} |S_t - \bar{S}_t|^2 du \leq \int_0^s \sup_{t \in [0, u]} |S_{t \wedge \tau} - \bar{S}_{t \wedge \tau}|^2 du.$$

Substituting back into (3.94) with (3.99)–(3.103), we deduce that

$$\mathbb{E} \left[\sup_{t \in [0, s]} |S_{t \wedge \tau} - \bar{S}_{t \wedge \tau}|^2 \right] \leq \beta_2 \int_0^s \mathbb{E} \left[\sup_{t \in [0, u]} |S_{t \wedge \tau} - \bar{S}_{t \wedge \tau}|^2 \right] du + \beta_3 (\delta t)^{2\alpha}$$

$$\begin{aligned}
& + \beta_4 \sup_{t \in [0, T]} \mathbb{E} \left[|v_t - \bar{v}_t| \right] + \beta_5 \sup_{t \in [0, T]} \mathbb{E} \left[|g_t^d - \bar{g}_t^d|^2 \right] \\
& + \beta_6 \sup_{t \in [0, T]} \mathbb{E} \left[|\hat{g}_t^f - \bar{g}_t^f| \right] + \beta_7 \sup_{t \in [0, T]} \mathbb{E} \left[|\hat{g}_t^f - \bar{g}_t^f|^2 \right] \\
(3.104) \quad & + \beta_8 \int_0^T \mathbb{E} \left[|S_t - \bar{S}_t|^2 \mathbf{1}_{t < \tau} \right] dt,
\end{aligned}$$

where the constants β_i , for $2 \leq i \leq 8$, are defined below.

$$\begin{aligned}
\beta_2 &= 32\sigma_{max}^2 L_v + 8(L_d^2 + L_f^2 + 4h_{max}^2)T + 96B^2 L_s^2 L_v (1 + 4\rho_{sf}^2 \xi_f^2 L_f T^3 e^{\beta_1 T}), \\
\beta_3 &= 96A^2 L_s^2 L_v (1 + 4\rho_{sf}^2 \xi_f^2 L_f T^3 e^{\beta_1 T})T, \quad \beta_4 = 32\sigma_{max}^2 L_s^2 T (1 + 4\rho_{sf}^2 \xi_f^2 L_f T^3 e^{\beta_1 T}), \\
\beta_5 &= 8L_s^2 T^2, \quad \beta_6 = 128\xi_f^2 L_s^2 (4 + \rho_{sf}^2 \sigma_{max}^2 L_v T)T^3 e^{\beta_1 T}, \quad \beta_7 = 16L_s^2 T^2 (1 + 8k_f^2 T^2 e^{\beta_1 T}), \\
(3.105) \quad & \beta_8 = 96B^2 L_s^2 L_v (1 + 4\rho_{sf}^2 \xi_f^2 L_f T^3 e^{\beta_1 T}).
\end{aligned}$$

Therefore, applying Gronwall's inequality to (3.104), we obtain

$$\begin{aligned}
\mathbb{E} \left[\sup_{t \in [0, T]} |S_{t \wedge \tau} - \bar{S}_{t \wedge \tau}|^2 \right] &\leq e^{\beta_2 T} \left\{ \beta_3 (\delta t)^{2\alpha} + \beta_4 \sup_{t \in [0, T]} \mathbb{E} \left[|v_t - \bar{v}_t| \right] + \beta_5 \sup_{t \in [0, T]} \mathbb{E} \left[|g_t^d - \bar{g}_t^d|^2 \right] \right. \\
&+ \beta_6 \sup_{t \in [0, T]} \mathbb{E} \left[|\hat{g}_t^f - \bar{g}_t^f| \right] + \beta_7 \sup_{t \in [0, T]} \mathbb{E} \left[|\hat{g}_t^f - \bar{g}_t^f|^2 \right] \\
(3.106) \quad &+ \left. \beta_8 \int_0^T \mathbb{E} \left[|S_t - \bar{S}_t|^2 \mathbf{1}_{t < \tau} \right] dt \right\}.
\end{aligned}$$

The convergence of the first term on the right-hand side as $\delta t \rightarrow 0$ is trivial, whereas the convergence of the next two terms is a consequence of [Proposition 3.9](#). We now show the convergence of the L^2 difference between \hat{g}^f and \bar{g}^f .

Suppose that $t \in [t_n, t_{n+1})$. Since $|\hat{g}_t^f - \bar{g}_t^f| \leq |\bar{g}_t^f - \bar{g}_{t_n}^f|$ and $2\sqrt{|ab|} \leq |a| + |b|$, we can bound the L^2 difference from above as follows:

$$\begin{aligned}
|\hat{g}_t^f - \bar{g}_t^f|^2 &\leq \left(k_f \theta_f \delta t + \frac{1}{2} |\rho_{sf}| \xi_f \sigma_{max} \delta t \hat{v}_{t_n} + \left(k_f + \frac{1}{2} |\rho_{sf}| \xi_f \sigma_{max} \right) \delta t \hat{g}_{t_n}^f \right. \\
&+ \left. \xi_f \sqrt{\hat{g}_{t_n}^f} |W_t^f - W_{t_n}^f| \right)^2 \\
&\leq 4k_f^2 \theta_f^2 (\delta t)^2 + \rho_{sf}^2 \xi_f^2 \sigma_{max}^2 (\delta t)^2 \hat{v}_{t_n}^2 + (2k_f + |\rho_{sf}| \xi_f \sigma_{max})^2 (\delta t)^2 (\hat{g}_{t_n}^f)^2 \\
(3.107) \quad &+ 4\xi_f^2 \hat{g}_{t_n}^f (W_t^f - W_{t_n}^f)^2.
\end{aligned}$$

Hence,

$$\begin{aligned}
\sup_{t \in [0, T]} \mathbb{E} \left[|\hat{g}_t^f - \bar{g}_t^f|^2 \right] &\leq 4k_f^2 \theta_f^2 (\delta t)^2 + \rho_{sf}^2 \xi_f^2 \sigma_{max}^2 (\delta t)^2 \sup_{0 \leq n \leq N} \mathbb{E} \left[\hat{v}_{t_n}^2 \right] + 4\xi_f^2 \delta t \sup_{0 \leq n \leq N} \mathbb{E} \left[\hat{g}_{t_n}^f \right] \\
(3.108) \quad &+ (2k_f + |\rho_{sf}| \xi_f \sigma_{max})^2 (\delta t)^2 \sup_{0 \leq n \leq N} \mathbb{E} \left[(\hat{g}_{t_n}^f)^2 \right].
\end{aligned}$$

The convergence of the term on the left-hand side follows from [Propositions 3.7](#) and [3.8](#).

Finally, the integrand in the last term in [\(3.106\)](#) can be bounded from above as follows:

$$\begin{aligned}
\mathbb{E} \left[|S_t - S_{\bar{t}}|^2 \mathbf{1}_{t < \tau} \right] &= \mathbb{E} \left[\left| \int_{\bar{t}}^t [g_u^d - g_u^f + h(u)] S_u du + \int_{\bar{t}}^t \sigma(u, S_u) \sqrt{v_u} S_u dW_u^s \right|^2 \mathbf{1}_{t < \tau} \right] \\
&\leq 4\delta t \mathbb{E} \left[\int_{\bar{t}}^t [(g_u^d)^2 + (g_u^f)^2 + h(u)^2] S_u^2 \mathbf{1}_{u < \tau} du \right] \\
&\quad + 4 \mathbb{E} \left[\left| \int_{\bar{t}}^t \sigma(u, S_u) \sqrt{v_u} S_u \mathbf{1}_{u < \tau} dW_u^s \right|^2 \right] \\
&\leq 4L_s^2 (\delta t)^2 \left\{ \sup_{u \in [0, T]} \mathbb{E} [(g_u^d)^2] + \sup_{u \in [0, T]} \mathbb{E} [(g_u^f)^2] + 4h_{max}^2 \right\} \\
(3.109) \quad &\quad + 4\sigma_{max}^2 L_s^2 L_v \delta t.
\end{aligned}$$

The right-hand side is independent of the time t and tends to zero as $\delta t \rightarrow 0$ from [Propositions 3.7](#) and [3.8](#), which concludes the proof. \blacksquare

Next, we prove the uniform convergence in probability of the discretized spot FX rate process.

Proposition 3.11. *If $2k_f \theta_f > \xi_f^2$, and under assumptions $(\mathcal{A}1)$ to $(\mathcal{A}3)$, \bar{S} converges uniformly in probability, i.e.,*

$$(3.110) \quad \lim_{\delta t \rightarrow 0} \mathbb{P} \left(\sup_{t \in [0, T]} |S_t - \bar{S}_t| > \epsilon \right) = 0 \quad \forall \epsilon > 0.$$

Proof. Fix $\epsilon > 0$ and note that

$$(3.111) \quad \left\{ \sup_{t \in [0, T]} |S_t - \bar{S}_t| > \epsilon \right\} \subseteq \left\{ \sup_{t \in [0, T]} |S_{t \wedge \tau} - \bar{S}_{t \wedge \tau}| > \epsilon \right\} \cup \{\tau < T\},$$

with τ from [\(3.89\)](#). In terms of probabilities of events, the previous inclusion implies that

$$(3.112) \quad \mathbb{P} \left(\sup_{t \in [0, T]} |S_t - \bar{S}_t| > \epsilon \right) \leq \mathbb{P} \left(\sup_{t \in [0, T]} |S_{t \wedge \tau} - \bar{S}_{t \wedge \tau}| > \epsilon \right) + \mathbb{P}(\tau < T).$$

The convergence in probability of the stopped process is an immediate consequence of [Proposition 3.10](#) and Markov's inequality. Moreover, from the definition of the stopping time in [\(3.89\)](#), we get

$$\begin{aligned}
\{\tau < T\} &\subseteq \left\{ \sup_{t \in [0, T]} v_t \geq L_v \right\} \cup \left\{ \sup_{t \in [0, T]} \hat{v}_t \geq L_v \right\} \cup \left\{ \sup_{t \in [0, T]} \hat{g}_t^d \geq L_d \right\} \cup \left\{ \sup_{t \in [0, T]} \hat{g}_t^f \geq L_f \right\} \\
(3.113) \quad &\cup \left\{ \sup_{t \in [0, T]} S_t \geq L_s \right\} \cup \{\tau_\kappa < T\},
\end{aligned}$$

where we define

$$(3.114) \quad \tau_\kappa = \inf\{t \geq 0 : g_t^f \leq \kappa^{-1}\}.$$

However, assuming that $L_s > 1$, we have that

$$(3.115) \quad \left\{ \sup_{t \in [0, T]} S_t \geq L_s \right\} = \left\{ \sup_{t \in [0, T]} x_t \geq \log L_s \right\} \subseteq \left\{ \sup_{t \in [0, T]} |x_t| \geq \log L_s \right\}.$$

Therefore, using (3.113), (3.115), and Markov's inequality, we find an upper bound

$$(3.116) \quad \begin{aligned} \mathbb{P}(\tau < T) &\leq L_v^{-1} \mathbb{E} \left[\sup_{t \in [0, T]} v_t \right] + L_v^{-1} \mathbb{E} \left[\sup_{t \in [0, T]} \hat{v}_t \right] + L_d^{-1} \mathbb{E} \left[\sup_{t \in [0, T]} \hat{g}_t^d \right] + L_f^{-1} \mathbb{E} \left[\sup_{t \in [0, T]} \hat{g}_t^f \right] \\ &+ (\log L_s)^{-1} \mathbb{E} \left[\sup_{t \in [0, T]} |x_t| \right] + \mathbb{P}(\tau_\kappa < T). \end{aligned}$$

First, recall the formula for the logarithm of the spot FX rate process,

$$(3.117) \quad x_t = x_0 + \int_0^t \left(g_u^d - g_u^f + h(u) - \frac{1}{2} \sigma^2(u, S_u) v_u \right) du + \int_0^t \sigma(u, S_u) \sqrt{v_u} dW_u^s.$$

Taking the supremum over all $t \in [0, T]$ and using the fact that $a < a^2 + 1$, and then taking expectations and employing Doob's martingale inequality and the Itô isometry, we get

$$(3.118) \quad \begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} |x_t| \right] &\leq 1 + |x_0| + 2h_{max}T + T \mathbb{E} \left[\sup_{t \in [0, T]} g_t^d \right] + T \mathbb{E} \left[\sup_{t \in [0, T]} g_t^f \right] \\ &+ \frac{9}{2} \sigma_{max}^2 T \mathbb{E} \left[\sup_{t \in [0, T]} v_t \right]. \end{aligned}$$

Second, we prove the almost sure positivity of the process g^f when $2k_f\theta_f > \xi_f^2$. To this end, let $\alpha = (2k_f\theta_f - \xi_f^2)/2\xi_f^2$ and define the function $F : (0, \infty) \rightarrow \mathbb{R}$ by $F(x) = x^{-\alpha}$. Using Itô's formula, we have that

$$(3.119) \quad \begin{aligned} \mathbb{E} \left[F(g_{T \wedge \tau_\kappa}^f) \right] &= F(g_0^f) - \mathbb{E} \int_0^{T \wedge \tau_\kappa} \alpha (g_u^f)^{-(1+\alpha)} (k_f\theta_f - k_f g_u^f - \rho_{sf} \xi_f \sigma(u, S_u) \sqrt{v_u g_u^f}) du \\ &+ \frac{1}{2} \mathbb{E} \int_0^{T \wedge \tau_\kappa} \alpha(1+\alpha) \xi_f^2 (g_u^f)^{-(1+\alpha)} du - \mathbb{E} \int_0^{T \wedge \tau_\kappa} \alpha \xi_f (g_u^f)^{-(\frac{1}{2}+\alpha)} dW_u^f. \end{aligned}$$

Note that

$$(3.120) \quad \alpha k_f \theta_f - \frac{1}{2} \alpha(1+\alpha) \xi_f^2 = \frac{1}{4} \alpha [4k_f \theta_f - 2\xi_f^2 - 2\alpha \xi_f^2] = \frac{1}{2} \alpha^2 \xi_f^2$$

and also that

$$(3.121) \quad \mathbb{E} \int_0^T \alpha^2 \xi_f^2 (g_u^f)^{-(1+2\alpha)} \mathbf{1}_{u < \tau_\kappa} du \leq \alpha^2 \xi_f^2 \kappa^{1+2\alpha} T < \infty,$$

so the stochastic integral on the right-hand side of (3.119) is a true martingale. Hence,

$$(3.122) \quad \begin{aligned} \mathbb{E} \left[F(g_{T \wedge \tau_\kappa}^f) \right] &\leq F(g_0^f) - \frac{1}{2} \alpha^2 \xi_f^2 \mathbb{E} \int_0^T (g_u^f)^{-(1+\alpha)} \mathbf{1}_{u < \tau_\kappa} du + \alpha k_f \mathbb{E} \int_0^T (g_u^f)^{-\alpha} \mathbf{1}_{u < \tau_\kappa} du \\ &+ \alpha |\rho_{sf}| \xi_f \sigma_{max} \mathbb{E} \int_0^T v_u^{\frac{1}{2}} (g_u^f)^{-\left(\frac{1}{2} + \alpha\right)} \mathbf{1}_{u < \tau_\kappa} du. \end{aligned}$$

Employing Fubini's theorem and Hölder's inequality, we deduce that

$$(3.123) \quad \begin{aligned} \mathbb{E} \left[F(g_{T \wedge \tau_\kappa}^f) \right] &\leq -\frac{1}{2} \alpha^2 \xi_f^2 \int_0^T \left(\mathbb{E} \left[(g_u^f)^{-(1+\alpha)} \mathbf{1}_{u < \tau_\kappa} \right] - 2\alpha^{-1} k_f \xi_f^{-2} \mathbb{E} \left[(g_u^f)^{-(1+\alpha)} \mathbf{1}_{u < \tau_\kappa} \right]^{\frac{\alpha}{1+\alpha}} \right. \\ &\quad \left. - 2\alpha^{-1} |\rho_{sf}| \xi_f^{-1} \sigma_{max} \sup_{t \in [0, T]} \mathbb{E} \left[v_t^{1+\alpha} \right]^{\frac{1}{2(1+\alpha)}} \mathbb{E} \left[(g_u^f)^{-(1+\alpha)} \mathbf{1}_{u < \tau_\kappa} \right]^{\frac{1+2\alpha}{2(1+\alpha)}} \right) du \\ &+ F(g_0^f). \end{aligned}$$

However, the moments of the square-root process are bounded by Proposition 3.7. Furthermore, if $p, q \in (0, 1)$, $c_{p,q} \geq 0$, and $c = c_p + c_q > 0$, then for all $x \geq 0$ we have

$$(3.124) \quad \begin{aligned} x - c_p x^p - c_q x^q &= \frac{c_p}{c} (x - c x^p) + \frac{c_q}{c} (x - c x^q) \\ &= c_p c^{\frac{p}{1-p}} \left[c^{-\frac{1}{1-p}} x - \left(c^{-\frac{1}{1-p}} x \right)^p \right] + c_q c^{\frac{q}{1-q}} \left[c^{-\frac{1}{1-q}} x - \left(c^{-\frac{1}{1-q}} x \right)^q \right] \\ &\geq -c_p c^{\frac{p}{1-p}} - c_q c^{\frac{q}{1-q}}. \end{aligned}$$

Therefore, the integrand in (3.123) is bounded from below by a constant, and thus

$$(3.125) \quad \mathbb{E} \left[F(g_{T \wedge \tau_\kappa}^f) \right] \leq C$$

for some constant C independent of κ . Because g^f has continuous paths, $g_{\tau_\kappa}^f = \kappa^{-1}$, and so $F(g_{\tau_\kappa}^f) = \kappa^\alpha$. Hence, from (3.125) and the positivity of F , we deduce that

$$(3.126) \quad \mathbb{P}(\tau_\kappa < T) = \kappa^{-\alpha} \mathbb{E} \left[F(g_{\tau_\kappa}^f) \mathbf{1}_{\tau_\kappa < T} \right] \leq \kappa^{-\alpha} \mathbb{E} \left[F(g_{T \wedge \tau_\kappa}^f) \right] \leq C \kappa^{-\alpha}.$$

Taking the limit as $\delta t \rightarrow 0$ in (3.112), using the upper bounds derived in (3.116), (3.118), and (3.126), and then employing Propositions 3.7 and 3.8, the conclusion follows from the fact that we can choose L_s, L_v, L_d, L_f , and κ arbitrarily large. \blacksquare

Many models with stochastic volatility dynamics, including the Heston model, have the undesirable feature of moment instability; i.e., moments of order higher than 1 can explode in finite time [5]. This can cause problems in practice, for instance when computing the arbitrage-free price of an option with a superlinear payoff. Furthermore, establishing the existence of moments of order higher than 1 of the process and its approximation is an important ingredient in the convergence analysis (see [24]). Since the most popular FX contracts grow at most linearly in the FX rate and their risk-neutral valuation involves computing the expected

discounted payoff, it is useful to study the finiteness of moments under discounting. Define R to be the discounted exchange rate process,

$$(3.127) \quad R_t = S_0 \exp \left\{ - \int_0^t r_u^f du - \frac{1}{2} \int_0^t \sigma^2(u, S_u) v_u du + \int_0^t \sigma(u, S_u) \sqrt{v_u} dW_u^s \right\},$$

and let \bar{R} be its continuous-time approximation,

$$(3.128) \quad \bar{R}_t = S_0 \exp \left\{ - \int_0^t \bar{r}_u^f du - \frac{1}{2} \int_0^t \bar{\sigma}^2(u, \bar{S}_u) \bar{v}_u du + \int_0^t \bar{\sigma}(u, \bar{S}_u) \sqrt{\bar{v}_u} dW_u^s \right\}.$$

The next result establishes a lower bound on the explosion time of moments of order higher than 1 of the discounted process and is an important step in proving [Theorem 2.1](#) (Theorem 1 in [\[11\]](#)), which, in turn, plays a key role in the calibration of the model.

Proposition 3.12. *Let $\alpha \geq 1$. Under assumptions $(\mathcal{A}1)$ and $(\mathcal{A}3)$, if $T < T^*$, then there exists $w_1 > \alpha$ such that, for all $w \in [1, w_1)$, the following holds:*

$$(3.129) \quad \sup_{t \in [0, T]} \mathbb{E} [R_t^w] < \infty,$$

where T^* is as given below.

(1) When $k < \varphi(\alpha)\zeta$,

$$(3.130) \quad T^* = \frac{2}{\sqrt{\varphi(\alpha)^2 \zeta^2 - k^2}} \left[\frac{\pi}{2} + \arctan \left(\frac{k}{\sqrt{\varphi(\alpha)^2 \zeta^2 - k^2}} \right) \right],$$

where

$$(3.131) \quad \varphi(\alpha) = \alpha + \sqrt{(\alpha - 1)\alpha} \quad \text{and} \quad \zeta = \xi \sigma_{max}.$$

(2) When $k \geq \varphi(\alpha)\zeta$,

$$(3.132) \quad T^* = \infty.$$

Proof. First, we find it convenient to define a new stochastic process L by

$$(3.133) \quad L_t \equiv S_0 \exp \left\{ h_{max} t - \frac{1}{2} \int_0^t \sigma^2(u, S_u) v_u du + \int_0^t \sigma(u, S_u) \sqrt{v_u} dW_u^s \right\}.$$

As $R_t \leq L_t$ for all $t \in [0, T]$, it suffices to prove the finiteness of the supremum over t of

$$(3.134) \quad \mathbb{E} [L_t^w] = S_0^w \mathbb{E} \left[\exp \left\{ wh_{max} t + w \int_0^t \sigma(u, S_u) \sqrt{v_u} dW_u^s - \frac{w}{2} \int_0^t \sigma^2(u, S_u) v_u du \right\} \right].$$

Second, suppose that $T < T^*$, with T^* defined in [\(3.130\)](#)–[\(3.132\)](#). If $k < \varphi(\alpha)\zeta$, since $\varphi(\cdot)$ is increasing on $[1, \infty)$ and by a continuity argument, we can find $w_1 > \alpha$ such that, for all $w \in (\alpha, w_1)$,

$$(3.135) \quad k < \varphi(w)\zeta \quad \text{and} \quad T < \frac{2}{\sqrt{\varphi(w)^2 \zeta^2 - k^2}} \left[\frac{\pi}{2} + \arctan \left(\frac{k}{\sqrt{\varphi(w)^2 \zeta^2 - k^2}} \right) \right].$$

If $k = \varphi(\alpha)\zeta$, since $\varphi(\cdot)$ is strictly increasing on $[1, \infty)$ and

$$(3.136) \quad \lim_{w \downarrow \alpha^+} \frac{2}{\sqrt{\varphi(w)^2 \zeta^2 - k^2}} \left[\frac{\pi}{2} + \arctan \left(\frac{k}{\sqrt{\varphi(w)^2 \zeta^2 - k^2}} \right) \right] = \infty,$$

we can find $w_1 > \alpha$ such that (3.135) holds for all $w \in (\alpha, w_1)$. Finally, if $k > \varphi(\alpha)\zeta$, by a continuity argument, we can find $w_1 > \alpha$ such that, for all $w \in (\alpha, w_1)$,

$$(3.137) \quad k > \varphi(w)\zeta.$$

Third, fix $w \in (\alpha, w_1)$ and let the functions $f_w, g_w : (1, \infty) \rightarrow (1, \infty)$ be defined by

$$(3.138) \quad f_w(x) = w^2 x^2$$

and

$$(3.139) \quad g_w(x) = \frac{wx}{x-1}(wx-1) = (w + \sqrt{(w-1)w})^2 + \frac{w^2}{x-1}(x-1 - \sqrt{1-1/w})^2.$$

The first function is strictly increasing, whereas the second function attains a global minimum at $x^* = 1 + \sqrt{1-1/w}$. Furthermore, they are equal at x^* , so

$$(3.140) \quad \inf_{x>1} \max \{f_w(x), g_w(x)\} = g_w(x^*) = (w + \sqrt{(w-1)w})^2.$$

Consider the Hölder pair (p, q) satisfying $q = p/(p-1)$ such that

$$(3.141) \quad p = 1 + \sqrt{\frac{w-1}{w}} \quad \text{and} \quad q = 1 + \sqrt{\frac{w}{w-1}}.$$

This particular Hölder pair ensures an optimal lower bound T^* on the explosion time. Next, define the quantity $a = pw^2 - w$ and introduce the stochastic process

$$(3.142) \quad M_t = pw \int_0^t \sigma(u, S_u) \sqrt{v_u} dW_u^s$$

with quadratic variation

$$(3.143) \quad \langle M \rangle_t = p^2 w^2 \int_0^t \sigma^2(u, S_u) v_u du.$$

Then we can rewrite (3.134) as follows:

$$(3.144) \quad \mathbb{E} [L_t^w] = S_0^w e^{wh_{max}t} \mathbb{E} \left[\exp \left\{ \frac{1}{p} \left[M_t - \frac{1}{2} \langle M \rangle_t \right] + \frac{a}{2} \int_0^t \sigma^2(u, S_u) v_u du \right\} \right].$$

Applying Hölder's inequality with the pair (p, q) from (3.141) and taking the supremum over $[0, T]$,

$$\sup_{t \in [0, T]} \mathbb{E} [L_t^w] \leq S_0^w e^{wh_{max}T} \sup_{t \in [0, T]} \mathbb{E} \left[\exp \left\{ M_t - \frac{1}{2} \langle M \rangle_t \right\} \right]^{\frac{1}{p}}$$

$$(3.145) \quad \times \mathbb{E} \left[\exp \left\{ \frac{1}{2} qw(pw - 1) \sigma_{max}^2 \int_0^T v_u du \right\} \right]^{\frac{1}{q}}.$$

The stochastic exponential is a martingale if Novikov's condition is satisfied, i.e.,

$$(3.146) \quad \mathbb{E} \left[\exp \left\{ \frac{1}{2} \langle M \rangle_T \right\} \right] \leq \mathbb{E} \left[\exp \left\{ \frac{1}{2} p^2 w^2 \sigma_{max}^2 \int_0^T v_u du \right\} \right] < \infty.$$

The finiteness of the two expectations in (3.145) follows from (3.135)–(3.141) and Proposition 3.5. The extension to the interval $[1, w_1]$ follows from Jensen's inequality. ■

Proposition 3.13. *Let $\alpha \geq 1$. Under assumptions (A1) and (A3), if $T < T^*$, then there exists $w_2 > \alpha$ such that, for all $w \in [1, w_2]$ and $\delta_T < k^{-1}$, the following holds:*

$$(3.147) \quad \sup_{\delta t \in (0, \delta_T)} \sup_{t \in [0, T]} \mathbb{E} [(\bar{R}_t)^w] < \infty,$$

where T^* is as given below.

(1) When $2k \leq \varphi(\alpha)\zeta$,

$$(3.148) \quad T^* = \frac{1}{\varphi(\alpha)\zeta - k},$$

where

$$(3.149) \quad \varphi(\alpha) = \alpha + \sqrt{(\alpha - 1)\alpha} \quad \text{and} \quad \zeta = \xi \sigma_{max}.$$

(2) When $2k > \varphi(\alpha)\zeta$,

$$(3.150) \quad T^* = \frac{4k}{\varphi(\alpha)^2 \zeta^2}.$$

Proof. For convenience, define a new stochastic process \bar{L} by

$$(3.151) \quad \bar{L}_t \equiv S_0 \exp \left\{ h_{max} t - \frac{1}{2} \int_0^t \bar{\sigma}^2(u, \bar{S}_u) \bar{v}_u du + \int_0^t \bar{\sigma}(u, \bar{S}_u) \sqrt{\bar{v}_u} dW_u^s \right\}.$$

Since $\bar{R}_t \leq \bar{L}_t$ for all $t \in [0, T]$, it suffices to prove the finiteness of the supremum over t and δt of

$$(3.152) \quad \mathbb{E} [(\bar{L}_t)^w] = S_0^w \mathbb{E} \left[\exp \left\{ wh_{max} t + w \int_0^t \bar{\sigma}(u, \bar{S}_u) \sqrt{\bar{v}_u} dW_u^s - \frac{w}{2} \int_0^t \bar{\sigma}^2(u, \bar{S}_u) \bar{v}_u du \right\} \right].$$

Suppose that $T < T^*$, with T^* from (3.148)–(3.150). If $k \leq \frac{1}{2} \varphi(\alpha)\zeta$, by a continuity argument, we can find $w_2 > \alpha$ such that, for all $w \in (\alpha, w_2)$,

$$(3.153) \quad k < \frac{1}{2} \varphi(w)\zeta \quad \text{and} \quad T < \frac{1}{\varphi(w)\zeta - k}.$$

On the other hand, if $k > \frac{1}{2}\varphi(\alpha)\zeta$, by a continuity argument, we can find $w_2 > \alpha$ such that, for all $w \in (\alpha, w_2)$,

$$(3.154) \quad k > \frac{1}{2}\varphi(w)\zeta \quad \text{and} \quad T < \frac{4k}{\varphi(w)^2\zeta^2}.$$

Henceforth, we argue as in [Proposition 3.12](#) and use [Theorem 3.6](#) to deduce the finiteness of the supremum over t and δt of [\(3.152\)](#). \blacksquare

Higham, Mao, and Stuart [\[24\]](#) proved that for a locally Lipschitz SDE, the boundedness of the p th moments of the exact and numerical solutions, for some $p > 2$, ensures the strong mean square convergence of the Euler–Maruyama method. The existence of moment bounds for explicit Euler approximations, however, remained an open problem. Recently, Hutzenthaler, Jentzen, and Kloeden [\[27\]](#) studied SDEs with superlinearly growing coefficients and proved weak and strong divergence in L^p for all $p \geq 1$, and hence that the moment-bound assumption is not satisfied. To the best of our knowledge, the uniform boundedness of moments of order greater than 1 of discretization schemes for the Heston model and extensions thereof has not yet been established—this gap in the literature was also identified in Kloeden and Neuenkirch [\[32\]](#)—and our [Proposition 3.13](#) is the first result to address this issue.

Since the typical payoff of an FX contract grows at most linearly in the exchange rate, it suffices to know the strong convergence of the discounted process in L^1 to deduce the convergence of the time-discretization error to zero. The following theorem can be generalized to the L^α case relatively easily for all $\alpha \geq 1$, upon noticing that the critical time T^* from [\(3.130\)](#)–[\(3.132\)](#) is always greater than the one from [\(3.148\)](#)–[\(3.150\)](#).

Theorem 3.14. *Under assumptions [\(A1\)](#)–[\(A3\)](#), if $2k_f\theta_f > \xi_f^2$ and $T < T^*$, where*

$$(3.155) \quad T^* \equiv \frac{4k}{\zeta^2} \mathbb{1}_{\zeta < 2k} + \frac{1}{\zeta - k} \mathbb{1}_{\zeta \geq 2k}$$

and $\zeta = \xi\sigma_{max}$, then the discounted process converges strongly in L^1 , i.e.,

$$(3.156) \quad \lim_{\delta t \rightarrow 0} \sup_{t \in [0, T]} \mathbb{E} \left[|R_t - \bar{R}_t| \right] = 0.$$

Proof. Fix $\epsilon > 0$ and define the event $A = \{|R_t - \bar{R}_t| > \epsilon\}$; then

$$(3.157) \quad \sup_{t \in [0, T]} \mathbb{E} \left[|R_t - \bar{R}_t| \right] \leq \sup_{t \in [0, T]} \mathbb{E} \left[|R_t - \bar{R}_t| \mathbb{1}_{A^c} \right] + \sup_{t \in [0, T]} \mathbb{E} \left[|R_t - \bar{R}_t| \mathbb{1}_A \right].$$

Hence,

$$(3.158) \quad \sup_{t \in [0, T]} \mathbb{E} \left[|R_t - \bar{R}_t| \right] \leq \epsilon + \sup_{t \in [0, T]} \mathbb{E} \left[R_t \mathbb{1}_A \right] + \sup_{t \in [0, T]} \mathbb{E} \left[\bar{R}_t \mathbb{1}_A \right].$$

Choosing some $1 < w < \min\{w_1, w_2\}$ and applying Hölder’s inequality to the two expectations on the right-hand side with the pair $(p, q) = (w, \frac{w}{w-1})$ returns the following upper bound:

$$(3.159) \quad \sup_{t \in [0, T]} \mathbb{E} \left[|R_t - \bar{R}_t| \right] \leq \epsilon + \left\{ \sup_{t \in [0, T]} \mathbb{E} \left[R_t^w \right]^{\frac{1}{w}} + \sup_{t \in [0, T]} \mathbb{E} \left[(\bar{R}_t)^w \right]^{\frac{1}{w}} \right\} \sup_{t \in [0, T]} \mathbb{P} \left(|R_t - \bar{R}_t| > \epsilon \right)^{1 - \frac{1}{w}}.$$

Note that if $\zeta \geq 2k$, then

$$(3.160) \quad \frac{2}{\sqrt{\zeta^2 - k^2}} \left[\frac{\pi}{2} + \arctan \left(\frac{k}{\sqrt{\zeta^2 - k^2}} \right) \right] > \frac{\sqrt{3}}{\sqrt{\zeta^2 - k^2}} \geq \frac{1}{\sqrt{\zeta^2 - k^2}} \sqrt{\frac{\zeta + k}{\zeta - k}} = \frac{1}{\zeta - k}.$$

Moreover, if $k < \zeta < 2k$, then

$$(3.161) \quad \frac{2}{\sqrt{\zeta^2 - k^2}} \left[\frac{\pi}{2} + \arctan \left(\frac{k}{\sqrt{\zeta^2 - k^2}} \right) \right] > \frac{2}{\sqrt{\zeta^2 - k^2}} \geq \frac{4}{\sqrt{\zeta^2 - k^2}} \sqrt{\frac{k^2}{\zeta^2} \left(1 - \frac{k^2}{\zeta^2} \right)} = \frac{4k}{\zeta^2}.$$

Therefore, [Propositions 3.12](#) and [3.13](#) (with $\alpha = 1$) ensure the boundedness of moments of order w of the discounted process and its approximation. Furthermore, the convergence in probability of the discounted process is a simple consequence of [Proposition 3.11](#), by taking the domestic short rate to be zero. Finally, taking ϵ sufficiently small concludes the proof. ■

3.5. Option valuation. We now examine the convergence of Monte Carlo estimators for computing FX option prices when the dynamics of the exchange rate are governed by the Heston–2CIR++ SLV model and assumptions $(\mathcal{A}1)$ – $(\mathcal{A}3)$ are satisfied. As an aside, note that we discussed in [section 2](#) how other derivative pricing models, including popular models in equity markets, can be formulated as special cases. First, we consider European options.

Theorem 3.15. *Let $P = \mathbb{E}[e^{-\int_0^T r_t^d dt} (K - S_T)^+]$ be the arbitrage-free price of a European put option and $\bar{P} = \mathbb{E}[e^{-\int_0^T \bar{r}_t^d dt} (K - \bar{S}_T)^+]$ its approximation. If $2k_f \theta_f > \xi_f^2$, then*

$$(3.162) \quad \lim_{\delta t \rightarrow 0} |P - \bar{P}| = 0.$$

Proof. A simple string of inequalities gives the following upper bound:

$$(3.163) \quad \begin{aligned} |P - \bar{P}| &\leq \mathbb{E} \left[\left| \{e^{-\int_0^T r_t^d dt} - e^{-\int_0^T \bar{r}_t^d dt}\} (K - S_T)^+ + e^{-\int_0^T \bar{r}_t^d dt} \{(K - S_T)^+ - (K - \bar{S}_T)^+\} \right| \right] \\ &\leq K e^{h_{max} T} \mathbb{E} \left[\left| e^{-\int_0^T g_t^d dt} - e^{-\int_0^T \bar{g}_t^d dt} \right| \right] + e^{h_{max} T} \mathbb{E} \left[\left| (K - S_T)^+ - (K - \bar{S}_T)^+ \right| \right]. \end{aligned}$$

However, for any nonnegative numbers x and y , $|e^{-x} - e^{-y}| \leq |x - y|$, and so we can use Fubini's theorem to obtain an upper bound for the first expectation:

$$(3.164) \quad \begin{aligned} \sup_{t \in [0, T]} \mathbb{E} \left[\left| e^{-\int_t^T g_u^d du} - e^{-\int_t^T \bar{g}_u^d du} \right| \right] &\leq \sup_{t \in [0, T]} \int_t^T \mathbb{E} \left[|g_u^d - \bar{g}_u^d| \right] du \\ &\leq T \sup_{t \in [0, T]} \mathbb{E} \left[|g_t^d - \bar{g}_t^d| \right]. \end{aligned}$$

The right-hand side tends to zero by [Proposition 3.9](#). Define the two events $A = \{S_T < K\}$ and $\bar{A} = \{\bar{S}_T < K\}$, and denote the last expectation in [\(3.163\)](#) by \mathbb{J} . Then

$$(3.165) \quad \mathbb{J} = \mathbb{E} \left[\left| (K - S_T)^+ - (K - \bar{S}_T)^+ \right| (\mathbf{1}_{A \cap \bar{A}} + \mathbf{1}_{A \cap \bar{A}^c} + \mathbf{1}_{A^c \cap \bar{A}} + \mathbf{1}_{A^c \cap \bar{A}^c}) \right].$$

Therefore,

$$\begin{aligned}
 \mathbb{J} &\leq \mathbb{E} \left[|S_T - \bar{S}_T| \mathbf{1}_{A \cap \bar{A}} \right] + \mathbb{E} \left[(K - S_T) \mathbf{1}_{A \cap \bar{A}^c} \right] + \mathbb{E} \left[(K - \bar{S}_T) \mathbf{1}_{A^c \cap \bar{A}} \right] \\
 (3.166) \quad &\leq \mathbb{E} \left[|S_T - \bar{S}_T| \mathbf{1}_{A \cap \bar{A}} \right] + K \mathbb{P}(A \cap \bar{A}^c) + K \mathbb{P}(A^c \cap \bar{A}).
 \end{aligned}$$

Let δ be an arbitrary positive number; then we have the following inclusion of events:

$$\begin{aligned}
 A \cap \bar{A}^c &= \left(\{S_T \leq K - \delta\} \cup \{K - \delta < S_T < K\} \right) \cap \{\bar{S}_T \geq K\} \\
 &\subseteq \left(\{S_T \leq K - \delta\} \cap \{\bar{S}_T \geq K\} \right) \cup \{K - \delta < S_T < K\} \\
 (3.167) \quad &\subseteq \left\{ |S_T - \bar{S}_T| \geq \delta \right\} \cup \{K - \delta < S_T < K\}.
 \end{aligned}$$

In terms of probabilities of events, we have

$$(3.168) \quad \mathbb{P}(A \cap \bar{A}^c) \leq \mathbb{P}(|S_T - \bar{S}_T| \geq \delta) + \mathbb{P}(K - \delta < S_T < K) \quad \forall \delta > 0.$$

We can bound the second probability from above in a similar fashion,

$$(3.169) \quad A^c \cap \bar{A} \subseteq \left\{ |S_T - \bar{S}_T| \geq \delta \right\} \cup \{K \leq S_T < K + \delta\},$$

so

$$(3.170) \quad \mathbb{P}(A^c \cap \bar{A}) \leq \mathbb{P}(|S_T - \bar{S}_T| \geq \delta) + \mathbb{P}(K \leq S_T < K + \delta) \quad \forall \delta > 0.$$

For a suitable choice of δ , the last terms on the right-hand side of (3.168) and (3.170) can be made arbitrarily small, whereas the first terms tend to zero by Proposition 3.11. Therefore, the two probabilities in (3.166) converge to zero as δt goes to zero. Finally, fix $\epsilon > 0$ and let $B = \{|S_T - \bar{S}_T| > \epsilon\}$. We can bound the expectation on the right-hand side of (3.166) as follows:

$$\begin{aligned}
 \mathbb{E} \left[|S_T - \bar{S}_T| \mathbf{1}_{A \cap \bar{A}} \right] &= \mathbb{E} \left[|S_T - \bar{S}_T| \mathbf{1}_{A \cap \bar{A}} \mathbf{1}_{B^c} \right] + \mathbb{E} \left[|S_T - \bar{S}_T| \mathbf{1}_{A \cap \bar{A}} \mathbf{1}_B \right] \\
 (3.171) \quad &\leq K \mathbb{P}(|S_T - \bar{S}_T| > \epsilon) + \epsilon.
 \end{aligned}$$

Taking the limit as $\delta t \rightarrow 0$, employing Proposition 3.11, and making use of the fact that ϵ can be made arbitrarily small leads to the conclusion. \blacksquare

Theorem 3.16. *Let $C = \mathbb{E}[e^{-\int_0^T r_t^d dt} (S_T - K)^+]$ be the arbitrage-free price of a European call and $\bar{C} = \mathbb{E}[e^{-\int_0^T \bar{r}_t^d dt} (\bar{S}_T - K)^+]$ its approximation. If $2k_f \theta_f > \xi_f^2$ and $T < T^*$, with T^* from (3.155), then*

$$(3.172) \quad \lim_{\delta t \rightarrow 0} |C - \bar{C}| = 0.$$

Proof. A simple string of inequalities gives the following upper bound:

$$(3.173) \quad \begin{aligned} |C - \bar{C}| &\leq \mathbb{E} \left[\left| (R_T - Ke^{-\int_0^T r_t^d dt})^+ - (\bar{R}_T - Ke^{-\int_0^T \bar{r}_t^d dt})^+ \right| \right] \\ &\leq Ke^{h_{max}T} \mathbb{E} \left[\left| e^{-\int_0^T g_t^d dt} - e^{-\int_0^T \bar{g}_t^d dt} \right| \right] + \mathbb{E} \left[|R_T - \bar{R}_T| \right]. \end{aligned}$$

The two expectations on the right-hand side tend to zero as $\delta t \rightarrow 0$ from (3.164) and Theorem 3.14, respectively. \blacksquare

Asian options depend on the average exchange rate over a predetermined time period. Because the average is less volatile than the underlying rate, Asian options are usually less expensive than their European counterparts and are commonly used in currency and commodity markets, for instance, to reduce the foreign currency exposure of a corporation expecting payments in foreign currency. For any $0 \leq s \leq t \leq T$, define the discount factors:

$$(3.174) \quad D_{s,t} = e^{-\int_s^t r_u^d du} \quad \text{and} \quad \bar{D}_{s,t} = e^{-\int_s^t \bar{r}_u^d du}.$$

Theorem 3.17. *Consider a fixed strike Asian option with arbitrage-free price*

$$(3.175) \quad U = \mathbb{E} \left[e^{-\int_0^T r_t^d dt} [\psi(A(0,T) - K)]^+ \right],$$

$$(3.176) \quad \bar{U} = \mathbb{E} \left[e^{-\int_0^T \bar{r}_t^d dt} [\psi(\bar{A}(0,T) - K)]^+ \right].$$

If $2k_f\theta_f > \xi_f^2$ and $T < T^*$, with T^* from (3.155), then

$$(3.177) \quad \lim_{\delta t \rightarrow 0} |U - \bar{U}| = 0.$$

Here, $A(0,T)$ represents the arithmetic average and $\psi = \pm 1$ depending on the payoff (call or put). For continuous monitoring, $A(0,T) = \frac{1}{T} \int_0^T S_t dt$ and $\bar{A}(0,T) = \frac{1}{T} \int_0^T \hat{S}_t dt = \frac{1}{T} \int_0^T \bar{S}_t dt$.

Proof. The absolute difference can be bounded from above by

$$(3.178) \quad |U - \bar{U}| \leq \mathbb{E} \left[\left| [\psi(D_{0,T}A(0,T) - KD_{0,T})]^+ - [\psi(\bar{D}_{0,T}\bar{A}(0,T) - K\bar{D}_{0,T})]^+ \right| \right].$$

Therefore, we end up with the following upper bound:

$$(3.179) \quad |U - \bar{U}| \leq Ke^{h_{max}T} \mathbb{E} \left[\left| e^{-\int_0^T g_t^d dt} - e^{-\int_0^T \bar{g}_t^d dt} \right| \right] + \mathbb{E} \left[|D_{0,T}A(0,T) - \bar{D}_{0,T}\bar{A}(0,T)| \right].$$

We deduced the convergence of the first expectation in (3.164). Fubini's theorem yields

$$(3.180) \quad \begin{aligned} \mathbb{E} \left[|D_{0,T}A(0,T) - \bar{D}_{0,T}\bar{A}(0,T)| \right] &\leq \frac{1}{T} \mathbb{E} \left[\int_0^T |D_{0,T}S_t - \bar{D}_{0,T}\bar{S}_t| dt \right] \\ &\leq \sup_{t \in [0,T]} \mathbb{E} \left[|D_{t,T}R_t - \bar{D}_{t,T}\bar{R}_t| \right]. \end{aligned}$$

The triangle inequality leads to the following upper bound:

$$(3.181) \quad \begin{aligned} \sup_{t \in [0, T]} \mathbb{E} \left[\left| D_{t, T} R_t - \bar{D}_{\bar{t}, T} \bar{R}_{\bar{t}} \right| \right] &\leq \sup_{t \in [0, T]} \mathbb{E} \left[R_t |D_{t, T} - D_{\bar{t}, T}| \right] + \sup_{t \in [0, T]} \mathbb{E} \left[D_{\bar{t}, T} |R_t - R_{\bar{t}}| \right] \\ &+ \sup_{t \in [0, T]} \mathbb{E} \left[R_{\bar{t}} |D_{\bar{t}, T} - \bar{D}_{\bar{t}, T}| \right] + \sup_{t \in [0, T]} \mathbb{E} \left[\bar{D}_{\bar{t}, T} |R_{\bar{t}} - \bar{R}_{\bar{t}}| \right], \end{aligned}$$

and hence

$$(3.182) \quad \begin{aligned} e^{-h_{max} T} \sup_{t \in [0, T]} \mathbb{E} \left[\left| D_{t, T} R_t - \bar{D}_{\bar{t}, T} \bar{R}_{\bar{t}} \right| \right] &\leq \sup_{t \in [0, T]} \mathbb{E} \left[R_t |1 - D_{\bar{t}, t}| \right] + \sup_{t \in [0, T]} \mathbb{E} \left[|R_t - R_{\bar{t}}| \right] \\ &+ \sup_{t \in [0, T]} \mathbb{E} \left[R_t \left| e^{-\int_t^T g_u^d du} - e^{-\int_t^T \bar{g}_u^d du} \right| \right] \\ &+ \sup_{t \in [0, T]} \mathbb{E} \left[|R_t - \bar{R}_t| \right]. \end{aligned}$$

First, Hölder's inequality with the pair (w, γ) , where $1 < w < w_1$ and $\gamma = w/(w-1)$, yields

$$(3.183) \quad \sup_{t \in [0, T]} \mathbb{E} \left[R_t |1 - D_{\bar{t}, t}| \right] \leq \sup_{t \in [0, T]} \mathbb{E} \left[R_t^w \right]^{\frac{1}{w}} \sup_{t \in [0, T]} \mathbb{E} \left[|1 - D_{\bar{t}, t}|^\gamma \right]^{\frac{1}{\gamma}}.$$

However, using the MV and Fubini theorems and Hölder's inequality, we get

$$(3.184) \quad \sup_{t \in [0, T]} \mathbb{E} \left[|1 - D_{\bar{t}, t}|^\gamma \right]^{\frac{1}{\gamma}} \leq e^{h_{max} T} \sup_{t \in [0, T]} \mathbb{E} \left[|r_t^d|^\gamma \right]^{\frac{1}{\gamma}} \delta t,$$

and hence the convergence of the first term on the right-hand side of (3.182) is a consequence of Propositions 3.7 and 3.12 (with $\alpha = 1$). Second, employing the MV theorem and Hölder's inequality, we deduce that

$$(3.185) \quad \begin{aligned} \sup_{t \in [0, T]} \mathbb{E} \left[|R_t - R_{\bar{t}}| \right] &\leq \sup_{t \in [0, T]} \mathbb{E} \left[(R_t \vee R_{\bar{t}}) |\log(R_t) - \log(R_{\bar{t}})| \right] \\ &\leq 2 \sup_{t \in [0, T]} \mathbb{E} \left[R_t^w \right]^{\frac{1}{w}} \sup_{t \in [0, T]} \mathbb{E} \left[|\log(R_t) - \log(R_{\bar{t}})|^\gamma \right]^{\frac{1}{\gamma}}, \end{aligned}$$

and hence the convergence of the second term on the right-hand side of (3.182) is a consequence of Propositions 3.7, 3.8 and 3.12 and Theorem 1 in [16]. Third, since both g^d and \bar{g}^d are non-negative processes, for any $\gamma > 1$ we have

$$(3.186) \quad \left| e^{-\int_t^T g_u^d du} - e^{-\int_t^T \bar{g}_u^d du} \right|^\gamma \leq \left| e^{-\int_t^T g_u^d du} - e^{-\int_t^T \bar{g}_u^d du} \right| \quad \forall t \in [0, T].$$

Applying Hölder's inequality and using (3.186) yields

$$(3.187) \quad \sup_{t \in [0, T]} \mathbb{E} \left[R_t \left| e^{-\int_t^T g_u^d du} - e^{-\int_t^T \bar{g}_u^d du} \right| \right] \leq \sup_{t \in [0, T]} \mathbb{E} \left[R_t^w \right]^{\frac{1}{w}} \sup_{t \in [0, T]} \mathbb{E} \left[\left| e^{-\int_t^T g_u^d du} - e^{-\int_t^T \bar{g}_u^d du} \right|^\gamma \right]^{\frac{1}{\gamma}},$$

and hence the convergence of the third term on the right-hand side of (3.182) is a consequence of Proposition 3.12 and (3.164). Finally, the convergence of the fourth term on the right-hand side of (3.182) is due to Theorem 3.14. In case of discrete monitoring or a floating strike, we follow the same argument. \blacksquare

Barrier options continue to gain popularity in many over-the-counter markets, including the FX market. Their popularity can be explained by two key factors. First, barrier options are useful in limiting the risk exposure of an investor in the FX market. Second, they offer additional flexibility and can match an investor's view on the market for a lower price than a vanilla option.

Theorem 3.18. *Consider an up-and-out barrier call with arbitrage-free price*

$$(3.188) \quad U = \mathbb{E} \left[e^{-\int_0^T r_t^d dt} (S_T - K)^+ \mathbf{1}_{\{\sup_{t \in [0, T]} S_t \leq B\}} \right],$$

$$(3.189) \quad \bar{U} = \mathbb{E} \left[e^{-\int_0^T \bar{r}_t^d dt} (\bar{S}_T - K)^+ \mathbf{1}_{\{\sup_{t \in [0, T]} \hat{S}_t \leq B\}} \right],$$

where K is the strike price and B is the barrier. If $2k_f \theta_f > \xi_f^2$, then

$$(3.190) \quad \lim_{\delta t \rightarrow 0} |U - \bar{U}| = 0.$$

Proof. Define the events $A = \{\sup_{t \in [0, T]} S_t \leq B\}$ and $\bar{A} = \{\sup_{t \in [0, T]} \hat{S}_t \leq B\}$; then

$$(3.191) \quad \begin{aligned} |U - \bar{U}| &\leq \mathbb{E} \left[\left| (D_{0, T} - \bar{D}_{0, T}) (S_T - K)^+ \mathbf{1}_A + \bar{D}_{0, T} \left\{ (S_T - K)^+ \mathbf{1}_A - (\bar{S}_T - K)^+ \mathbf{1}_{\bar{A}} \right\} \right| \right] \\ &\leq (B - K)^+ e^{h_{max} T} \mathbb{E} \left[\left| e^{-\int_0^T g_t^d dt} - e^{-\int_0^T \bar{g}_t^d dt} \right| \right] \\ &\quad + e^{h_{max} T} \mathbb{E} \left[\left| (S_T - K)^+ \mathbf{1}_A - (\bar{S}_T - K)^+ \mathbf{1}_{\bar{A}} \right| \right]. \end{aligned}$$

The first term tends to zero by (3.164), and we can rewrite the second term as follows:

$$(3.192) \quad \begin{aligned} &\mathbb{E} \left[\left| (S_T - K)^+ (\mathbf{1}_{A \cap \bar{A}^c} + \mathbf{1}_{A \cap \bar{A}}) - (\bar{S}_T - K)^+ (\mathbf{1}_{A \cap \bar{A}} + \mathbf{1}_{A^c \cap \bar{A}}) \right| \right] \\ &\leq \mathbb{E} \left[(S_T - K)^+ \mathbf{1}_{A \cap \bar{A}^c} \right] + \mathbb{E} \left[(\bar{S}_T - K)^+ \mathbf{1}_{A^c \cap \bar{A}} \right] + \mathbb{E} \left[|S_T - \bar{S}_T| \mathbf{1}_{A \cap \bar{A}} \right] \\ &\leq (B - K)^+ \left\{ \mathbb{P}(A \cap \bar{A}^c) + \mathbb{P}(A^c \cap \bar{A}) \right\} + \mathbb{E} \left[|S_T - \bar{S}_T| \mathbf{1}_{A \cap \bar{A}} \right]. \end{aligned}$$

We can bound the last expectation from above just as in (3.171) to find

$$(3.193) \quad \mathbb{E} \left[|S_T - \bar{S}_T| \mathbf{1}_{A \cap \bar{A}} \right] \leq B \mathbb{P} \left(|S_T - \bar{S}_T| > \epsilon \right) + \epsilon \quad \forall \epsilon > 0.$$

Hence, the expectation converges to zero with the time step by Proposition 3.11. Fix $\delta > 0$. First, note that

$$\begin{aligned} A \cap \bar{A}^c &\subseteq \left\{ \sup_{t \in [0, T]} S_t \leq B \right\} \cap \left\{ \sup_{t \in [0, T]} \bar{S}_t > B \right\} \\ &\subseteq \left(\left\{ \sup_{t \in [0, T]} S_t \leq B - \delta \right\} \cap \left\{ \sup_{t \in [0, T]} \bar{S}_t > B \right\} \right) \cup \left\{ B - \delta < \sup_{t \in [0, T]} S_t \leq B \right\} \end{aligned}$$

$$(3.194) \quad \subseteq \left\{ \sup_{t \in [0, T]} |S_t - \bar{S}_t| \geq \delta \right\} \cup \left\{ B - \delta < \sup_{t \in [0, T]} S_t \leq B \right\}.$$

In terms of probabilities of events, this implies that

$$(3.195) \quad \mathbb{P}(A \cap \bar{A}^c) \leq \mathbb{P}\left(\sup_{t \in [0, T]} |S_t - \bar{S}_t| \geq \delta\right) + \mathbb{P}\left(B - \delta < \sup_{t \in [0, T]} S_t \leq B\right).$$

Second, note that

$$(3.196) \quad \begin{aligned} A^c \cap \bar{A} &= \left\{ \sup_{t \in [0, T]} S_t > B \right\} \cap \left\{ \sup_{t \in [0, T]} \hat{S}_t \leq B \right\} \\ &\subseteq \left(\left\{ \sup_{t \in [0, T]} S_t > B + 2\delta \right\} \cap \left\{ \sup_{t \in [0, T]} \hat{S}_t \leq B \right\} \right) \cup \left\{ B < \sup_{t \in [0, T]} S_t \leq B + 2\delta \right\} \\ &\subseteq \left\{ \sup_{t \in [0, T]} |S_t - \hat{S}_t| > 2\delta \right\} \cup \left\{ B < \sup_{t \in [0, T]} S_t \leq B + 2\delta \right\} \\ &\subseteq \left\{ \sup_{t \in [0, T]} |S_t - S_{\bar{t}}| > \delta \right\} \cup \left\{ \sup_{t \in [0, T]} |S_{\bar{t}} - \bar{S}_{\bar{t}}| > \delta \right\} \cup \left\{ B < \sup_{t \in [0, T]} S_t \leq B + 2\delta \right\}. \end{aligned}$$

In terms of probabilities of events, this implies that

$$(3.197) \quad \begin{aligned} \mathbb{P}(A^c \cap \bar{A}) &\leq \mathbb{P}\left(\sup_{t \in [0, T]} |S_t - S_{\bar{t}}| > \delta\right) + \mathbb{P}\left(\sup_{t \in [0, T]} |S_t - \bar{S}_t| \geq \delta\right) \\ &\quad + \mathbb{P}\left(B < \sup_{t \in [0, T]} S_t \leq B + 2\delta\right). \end{aligned}$$

Next, fix $L_s > 1 \vee S_0$ and define the stopping time

$$(3.198) \quad \tau_s = \inf \{t \geq 0 : S_t \geq L_s\}.$$

Using the MV theorem and Markov's inequality, we deduce that

$$(3.199) \quad \begin{aligned} \mathbb{P}\left(\sup_{t \in [0, T]} |S_t - S_{\bar{t}}| > \delta\right) &\leq \mathbb{P}\left(\sup_{t \in [0, T \wedge \tau_s]} |S_t - S_{\bar{t}}| > \delta\right) + \mathbb{P}(\tau_s < T) \\ &\leq \mathbb{P}\left(\sup_{t \in [0, T]} |x_t - x_{\bar{t}}| > \frac{\delta}{L_s}\right) + \mathbb{P}\left(\sup_{t \in [0, T]} |x_t| \geq \log(L_s)\right) \\ &\leq \frac{L_s}{\delta} \mathbb{E}\left[\sup_{t \in [0, T]} |x_t - x_{\bar{t}}|\right] + \frac{1}{\log(L_s)} \mathbb{E}\left[\sup_{t \in [0, T]} |x_t|\right]. \end{aligned}$$

The convergence to zero of the first term on the right-hand side of (3.199) is a consequence of Propositions 3.7 and 3.8 and Theorem 1 in [16]. Using the upper bound derived in (3.118), the conclusion follows from Proposition 3.11 since δ can be made arbitrarily small and L_s can be made arbitrarily large independently of one another. \blacksquare

Theorem 3.19. *Consider any type of barrier put option with arbitrage-free price*

$$(3.200) \quad U = \mathbb{E} \left[e^{-\int_0^T r_t^d dt} (K - S_T)^+ \mathbf{1}_A \right],$$

$$(3.201) \quad \bar{U} = \mathbb{E} \left[e^{-\int_0^T \bar{r}_t^d dt} (K - \bar{S}_T)^+ \mathbf{1}_{\bar{A}} \right],$$

where the events A and \bar{A} depend on the type of barrier:

$$(3.202) \quad \left\{ \begin{array}{ll} \text{up-and-in:} & A = \left\{ \sup_{t \in [0, T]} S_t > B \right\}, \quad \bar{A} = \left\{ \sup_{t \in [0, T]} \hat{S}_t > B \right\}, \\ \text{up-and-out:} & A = \left\{ \sup_{t \in [0, T]} S_t \leq B \right\}, \quad \bar{A} = \left\{ \sup_{t \in [0, T]} \hat{S}_t \leq B \right\}, \\ \text{down-and-in:} & A = \left\{ \inf_{t \in [0, T]} S_t < B \right\}, \quad \bar{A} = \left\{ \inf_{t \in [0, T]} \hat{S}_t < B \right\}, \\ \text{down-and-out:} & A = \left\{ \inf_{t \in [0, T]} S_t \geq B \right\}, \quad \bar{A} = \left\{ \inf_{t \in [0, T]} \hat{S}_t \geq B \right\}. \end{array} \right.$$

If $2k_f \theta_f > \xi_f^2$, then

$$(3.203) \quad \lim_{\delta t \rightarrow 0} |U - \bar{U}| = 0.$$

Proof. An upper bound for the absolute difference can be obtained as follows:

$$(3.204) \quad \begin{aligned} |U - \bar{U}| &\leq \mathbb{E} \left[\left| (D_{0,T} - \bar{D}_{0,T}) (K - S_T)^+ \mathbf{1}_A + \bar{D}_{0,T} \left\{ (K - S_T)^+ \mathbf{1}_A - (K - \bar{S}_T)^+ \mathbf{1}_{\bar{A}} \right\} \right| \right] \\ &\leq K e^{h_{max} T} \mathbb{E} \left[\left| e^{-\int_0^T g_t^d dt} - e^{-\int_0^T \bar{g}_t^d dt} \right| \right] \\ &\quad + e^{h_{max} T} \mathbb{E} \left[\left| (K - S_T)^+ \mathbf{1}_A - (K - \bar{S}_T)^+ \mathbf{1}_{\bar{A}} \right| \right]. \end{aligned}$$

The first term tends to zero by (3.164) and we can bound the second term as in (3.192):

$$(3.205) \quad \begin{aligned} \mathbb{E} \left[\left| (K - S_T)^+ \mathbf{1}_A - (K - \bar{S}_T)^+ \mathbf{1}_{\bar{A}} \right| \right] &\leq K \left\{ \mathbb{P}(A \cap \bar{A}^c) + \mathbb{P}(A^c \cap \bar{A}) \right\} \\ &\quad + \mathbb{E} \left[\left| (K - S_T)^+ - (K - \bar{S}_T)^+ \right| \right]. \end{aligned}$$

The events A and \bar{A} differ with the barrier (down-and-in, down-and-out, up-and-in, up-and-out); however, one can show in a similar way to (3.195) and (3.197) that

$$(3.206) \quad \lim_{\delta t \rightarrow 0} \mathbb{P}(A \cap \bar{A}^c) = 0 \quad \text{and} \quad \lim_{\delta t \rightarrow 0} \mathbb{P}(A^c \cap \bar{A}) = 0$$

for any type of barrier. Finally, the convergence of the last term on the right-hand side of (3.205) was derived in Theorem 3.15, which concludes the proof. \blacksquare

Theorem 3.20. *Consider a down-and-in/out or up-and-in barrier call option with arbitrage-free price*

$$(3.207) \quad U = \mathbb{E} \left[e^{-\int_0^T r_t^d dt} (S_T - K)^+ \mathbf{1}_A \right],$$

$$(3.208) \quad \bar{U} = \mathbb{E} \left[e^{-\int_0^T \bar{r}_t^d dt} (\bar{S}_T - K)^+ \mathbf{1}_{\bar{A}} \right],$$

where the events A and \bar{A} depend on the type of barrier as in (3.202). If $2k_f\theta_f > \xi_f^2$ and $T < T^*$, with T^* from (3.155), then

$$(3.209) \quad \lim_{\delta t \rightarrow 0} |U - \bar{U}| = 0.$$

Proof. An upper bound for the absolute difference can be obtained as follows:

$$(3.210) \quad \begin{aligned} |U - \bar{U}| &\leq \mathbb{E} \left[|(R_T - KD_{0,T})^+ - (\bar{R}_T - K\bar{D}_{0,T})^+| \mathbf{1}_{A \cap \bar{A}} \right] \\ &+ \mathbb{E} \left[(R_T - KD_{0,T})^+ \mathbf{1}_{A \cap \bar{A}^c} \right] + \mathbb{E} \left[(\bar{R}_T - K\bar{D}_{0,T})^+ \mathbf{1}_{A^c \cap \bar{A}} \right]. \end{aligned}$$

Therefore, we end up with

$$(3.211) \quad \begin{aligned} |U - \bar{U}| &\leq \mathbb{E} \left[|R_T - \bar{R}_T| \right] + Ke^{h_{max}T} \mathbb{E} \left[|e^{-\int_0^T g_t^d dt} - e^{-\int_0^T \bar{g}_t^d dt}| \right] \\ &+ \mathbb{E} \left[R_T \mathbf{1}_{A \cap \bar{A}^c} \right] + \mathbb{E} \left[\bar{R}_T \mathbf{1}_{A^c \cap \bar{A}} \right]. \end{aligned}$$

The convergence of the first two terms on the right-hand side is a consequence of [Theorem 3.14](#) and [\(3.164\)](#), respectively. Applying Hölder's inequality with the pair (w, γ) to the last two terms, where $1 < w < \min\{w_1, w_2\}$, we find that

$$(3.212) \quad \mathbb{E} \left[R_T \mathbf{1}_{A \cap \bar{A}^c} \right] \leq \mathbb{E} \left[(R_T)^w \right]^{\frac{1}{w}} \mathbb{P} \left(A \cap \bar{A}^c \right)^{\frac{1}{\gamma}}$$

and

$$(3.213) \quad \mathbb{E} \left[\bar{R}_T \mathbf{1}_{A^c \cap \bar{A}} \right] \leq \mathbb{E} \left[(\bar{R}_T)^w \right]^{\frac{1}{w}} \mathbb{P} \left(A^c \cap \bar{A} \right)^{\frac{1}{\gamma}}.$$

Using [Propositions 3.12](#) and [3.13](#) and [\(3.206\)](#) concludes the proof. ■

Among the most developed exotic derivatives in the FX market are the double barrier options.

Theorem 3.21. *Consider a double knock-out call option with arbitrage-free price*

$$(3.214) \quad U = \mathbb{E} \left[e^{-\int_0^T r_t^d dt} (S_T - K)^+ \mathbf{1}_{\{\inf_{t \in [0, T]} S_t \geq L, \sup_{t \in [0, T]} S_t \leq B\}} \right],$$

$$(3.215) \quad \bar{U} = \mathbb{E} \left[e^{-\int_0^T \bar{r}_t^d dt} (\bar{S}_T - K)^+ \mathbf{1}_{\{\inf_{t \in [0, T]} \hat{S}_t \geq L, \sup_{t \in [0, T]} \hat{S}_t \leq B\}} \right],$$

where K is the strike and L, B are the lower and upper barriers, respectively. If $2k_f\theta_f > \xi_f^2$, then

$$(3.216) \quad \lim_{\delta t \rightarrow 0} |U - \bar{U}| = 0.$$

Proof. Note that we have the following inclusion of events:

$$(3.217) \quad \left\{ \inf_{t \in [0, T]} S_t \geq L, \sup_{t \in [0, T]} S_t \leq B \right\} \cap \left\{ \inf_{t \in [0, T]} \hat{S}_t \geq L, \sup_{t \in [0, T]} \hat{S}_t \leq B \right\}^c \\ \subseteq \left(\left\{ \sup_{t \in [0, T]} S_t \leq B \right\} \cap \left\{ \sup_{t \in [0, T]} \hat{S}_t > B \right\} \right) \cup \left(\left\{ \inf_{t \in [0, T]} S_t \geq L \right\} \cap \left\{ \inf_{t \in [0, T]} \hat{S}_t < L \right\} \right).$$

The rest of the argument closely follows that of [Theorem 3.18](#) and is thus omitted. ■

4. Numerical results. In this section, we consider the pricing problem for exotic products, particularly for structured notes embedding barrier features. Popular FX exotic products include the power reverse dual currency note (PRDC) [9], especially long-term (e.g., 30 years), and the forward accumulator [47]. Adding stochastic rates improves the pricing of both contracts for mid- to long-term expiries. Moreover, while the plain PRDC can be seen as a strip of vanilla options, adding SLV dynamics improves the pricing of some variations of the contract, like the trigger PRDC (a PRDC with a knock-out feature).

Henceforth, we consider an autocallable barrier dual currency note (ABDC) that is indexed on the EURUSD currency pair, with USD as the domestic currency and EUR as the foreign currency, with quarterly coupon payments. While barrier dual currency notes are typically short-term investments, we examine instead a mid-term alternative with an embedded autocallable note structure. For a nominal N (in USD), a rate (strike) K , and an expiry T , the life cycle of the contract is described below.

- (1) A monthly fixing schedule is defined for the exchange rate S such that the following hold:
 - If S crosses the up barrier B_{UO} , the contract is redeemed early and a coupon C_{ER} is paid. The early redemption time is denoted by τ_{ER} and can be infinite in case of no knock-out event.
 - If S crosses the down barrier B_{DI} , a short put contract is activated at expiry. The knock-in time is denoted by τ_{KI} .
- (2) If S is above B_{DI} , a coupon C (in USD, in % of the nominal) is paid quarterly at coupon dates $\{t_i, i = 1, \dots, M\}$, for a total of M periods, such that the accumulated coupon (in USD) at expiry is

$$(4.1) \quad N \sum_{i=1}^M \frac{D_{t_i}^d}{D_T^d} C \mathbb{1}_{t_i < \tau_{\text{ER}}} \mathbb{1}_{S_{t_i} > B_{\text{DI}}} + N \frac{D_{\tau_{\text{ER}}}^d}{D_T^d} C_{\text{ER}} \mathbb{1}_{\tau_{\text{ER}} \leq T}.$$

- (3) At expiry, if S is below K , the down barrier has been activated, and early redemption has not occurred, then the nominal is converted to EUR at the rate K .

This product would suit an investor who wants to outperform the money market account by taking the risk of conversion of the nominal value to the foreign currency. The autocall feature lowers the price of the product due to early termination in case the exchange rate crosses the up-and-out barrier, whereas the down-and-in barrier feature provides protection against the conversion risk at the expense of increasing the net present value (NPV). Hence, an investor would find this product attractive under the current volatile market conditions if they expected a less volatile market in the future, which would imply a lower NPV due

to the increased chance of early redemption and conversion. Over a longer time horizon, a very stable market would be the best outcome for the investor, since they would get all the coupons without converting the nominal at maturity. If conversion occurs at expiry, the investor receives N/K in EUR in exchange for the nominal N in USD. If the investor converts this amount to USD, the loss becomes $N(S_T/K - 1)$. Hence, the profit-and-loss (PnL) of the product is

$$(4.2) \quad N \sum_{i=1}^M \frac{D_{t_i}^d}{D_T^d} \mathcal{C} \mathbb{1}_{t_i < \tau_{\text{ER}}} \mathbb{1}_{S_{t_i} > B_{\text{DI}}} + N \frac{D_{\tau_{\text{ER}}}^d}{D_T^d} \mathcal{C}_{\text{ER}} \mathbb{1}_{\tau_{\text{ER}} \leq T} - \frac{N}{K} (K - S_T)^+ \mathbb{1}_{\tau_{\text{KI}} \leq T < \tau_{\text{ER}}}.$$

Hence, this product is a yield enhancement contract with no capital guarantee. The accrued coupon is sometimes converted at expiry with the nominal at the rate K . Finally, the NPV of the contract (in % of the nominal) is

$$(4.3) \quad \text{NPV} = \mathbb{E} \left[\sum_{i=1}^M D_{t_i}^d \mathcal{C} \mathbb{1}_{t_i < \tau_{\text{ER}}} \mathbb{1}_{S_{t_i} > B_{\text{DI}}} + D_{\tau_{\text{ER}}}^d \mathcal{C}_{\text{ER}} \mathbb{1}_{\tau_{\text{ER}} \leq T} - \frac{D_T^d}{K} (K - S_T)^+ \mathbb{1}_{\tau_{\text{KI}} \leq T < \tau_{\text{ER}}} \right],$$

where the expectation is taken under the risk-neutral measure. We assume the dynamics of S under this measure as specified in (2.1).

Next, consider the contract parameters from Table 4.1 as well as the calibrated model parameters and leverage function—to the EURUSD market data from March 18, 2016—from [11]. Suppose that barriers are monitored monthly, at the fixing dates, and coupons are paid quarterly.

Table 4.1

The contract parameters of an autocallable barrier dual currency note, where the nominal N is in USD, the expiry T is in years, the strike K and the barriers B_{UO} and B_{DI} are in % of S_0 , and the coupons \mathcal{C} and \mathcal{C}_{ER} are in USD, in % of N .

N	T	S_0	K	B_{UO}	B_{DI}	\mathcal{C}	\mathcal{C}_{ER}
100000	5Y	1.1271	105%	100%	95%	2.5%	1.5%

We employ the Monte Carlo simulation scheme defined in section 3 with 5×10^7 sample paths and 376 time steps per year to price this contract, and the numerical results are displayed in Table 4.2. Note that we computed the percentage change in premium (NPV) with respect to the reference model, i.e., the Heston–2CIR++ SLV model.

We infer from Table 4.2 that the price of the contract under the 2-factor SLV model is 1.21% higher than under the 4-factor SLV model, whereas the early redemption and the knock-in probabilities are the same under the two models. This suggests that the stochastic rates have a significant impact on the price of the contract even for a 5-year expiry. We also infer from Table 4.2 that the contract is highly underpriced and the early redemption probability is overestimated under the LV model (2.7). Furthermore, we calibrated in [11] the 4-factor SLV and SV models to EURUSD market data from March 18, 2016, and observed an almost perfect fit to vanilla options with the former as opposed to a poor fit with the latter. In conclusion, we notice two things. First, pricing exotic products with barrier features under

Table 4.2

The NPV of the contract in % of N , the 95% Monte Carlo confidence interval, the early redemption and knock-in probabilities, and the percentage change in NPV, for the autocallable barrier dual currency note specified in Table 4.1 under the 4-factor hybrid SLV model (2.1), the 2-factor SLV model with deterministic rates, and the LV model (2.7).

Model	NPV	95% CI	ER prob.	KI prob.	Change
Hybrid SLV	1.7247	(1.7228, 1.7265)	95.1%	19.5%	–
SLV	1.7456	(1.7438, 1.7474)	95.1%	19.5%	+1.21%
LV	1.2079	(1.2062, 1.2097)	95.9%	19.5%	–29.96%

SLV dynamics is of paramount importance. It is a well-known fact that pure SV models underestimate the knock-out probability, whereas pure LV models overestimate it, and the true price of the contract is believed to lie in between pure SV and pure LV model prices. In our case, a small difference in the probability of no knock-out (4.1% versus 4.9%), which is leveraged by the number of coupons detached during the life cycle of the contract, can lead to a big difference in the NPV, and this explains the need to use SLV models to improve the pricing performance. Second, adding stochastic rates to an SLV model is only relevant when pricing mid- to long-term structured notes, where the effect of the stochastic rates is accentuated. The impact of stochastic rates on the contract price is expected to increase in the presence of nonzero correlations between the spot FX rate and the short rates.

We could extend the model (2.1) to multifactor short rates when pricing exotic products where the rates appear explicitly in the payoff, for example, a spread option with the payoff

$$(4.4) \quad \left[\left(\frac{S_T - S_0}{S_0} \right) - L_T - K \right]^+,$$

where L_T is the Libor rate at the fixing date T . In this case, the calibration algorithm described in section 2 can still be applied, with a higher computational cost due to the more complex simulation scheme.

Table 4.3

The Monte Carlo estimates of the NPV of the contract specified in Table 4.1 for different numbers of time steps (per year) and 5×10^7 sample paths (for a standard deviation of 9.29×10^{-4}), the difference from the previous NPV estimate, and the empirical convergence order.

Time steps	NPV	Difference	Order
12	1.9081	–	–
24	1.8110	0.0971	–
48	1.7623	0.0487	0.996
96	1.7388	0.0234	1.057
192	1.7285	0.0103	1.184

We conclude this section with an empirical convergence analysis of our Monte Carlo simulation scheme. Since the product can be decomposed into a linear combination of first-order exotics, the theoretical convergence (without a rate) of Monte Carlo estimators follows automatically from the analysis in section 3. The data in Table 4.3 suggest a first-order convergence

of the time-discretization error. As an aside, note that in the case of continuously monitored barriers, we could use Brownian bridge techniques to recover the first-order convergence.

The empirical findings of this section demonstrate the importance of SLV dynamics as well as stochastic short rate dynamics for the pricing of long-dated exotic FX products. Furthermore, we verified the convergence of our Monte Carlo simulation scheme for one such product.

5. Conclusions. The efficient pricing and hedging of vanilla and exotic FX options requires an adequate model that takes into account both the local and the stochastic features of the volatility dynamics. We have put forward a 4-factor hybrid stochastic and local volatility model that combines state-of-the-art dynamics for the exchange rate with stochastic dynamics for the domestic and foreign short rates, and proposed a simple and efficient Monte Carlo simulation scheme for pricing FX options.

The convergence of Euler schemes in a stochastic-local volatility (SLV) context has yet to be established, and even the literature on the convergence of Euler schemes under stochastic volatility is scarce. The only previous published work related to this problem that we are aware of is [23], which proves the convergence of an Euler discretization with a reflection fix in the context of Heston's model and options with bounded payoffs. In this paper, we have made some realistic assumptions on the leverage function and established, up to a critical time, the strong convergence (without a rate) of the discounted exchange rate approximation and the convergence of Monte Carlo estimators for a number of vanilla and exotic products.

The analysis carried out in this paper can be extended to other financial derivatives, including digital options, forward-start options, and double-no-touch binary options, to name just a few. Furthermore, we may consider a multifactor extension of the short rate model, in which case the convergence analysis applies with some slight modifications of the proofs.

However, several unsettled questions remain, like the exact strong convergence rate of the full truncation scheme for the CIR process and the strong convergence rate of schemes for the type of SDEs studied in this paper. On top of these being interesting and practically relevant questions in their own right, a sufficiently high order enables the use of multilevel simulation, as in [17], with substantial efficiency improvements for the estimation of expected financial payoffs.

REFERENCES

- [1] R. AHLIP AND M. RUTKOWSKI, *Pricing of foreign exchange options under the Heston stochastic volatility model and CIR interest rates*, Quant. Finance, 13 (2013), pp. 955–966.
- [2] A. ALFONSI, *On the discretization schemes for the CIR (and Bessel squared) processes*, Monte Carlo Methods Appl., 11 (2005), pp. 355–384.
- [3] A. ALFONSI, *Strong order one convergence of a drift implicit Euler scheme: Application to the CIR process*, Statist. Probab. Lett., 83 (2013), pp. 602–607.
- [4] L. ANDERSEN, *Simple and efficient simulation of the Heston stochastic volatility model*, J. Comput. Finance, 11 (2008), pp. 1–42.
- [5] L. ANDERSEN AND V. PITERBARG, *Moment explosions in stochastic volatility models*, Finance Stoch., 11 (2007), pp. 29–50.
- [6] A. BERKAOUI, M. BOSSY, AND A. DIOP, *Euler scheme for SDEs with non-Lipschitz diffusion coefficient: Strong convergence*, ESAIM Probab. Statist., 12 (2008), pp. 1–11.

- [7] D. BRIGO AND F. MERCURIO, *A deterministic-shift extension of analytically-tractable and time-homogeneous short-rate models*, *Finance Stoch.*, 5 (2001), pp. 369–387.
- [8] M. BROADIE AND O. KAYA, *Exact simulation of stochastic volatility and other affine jump diffusion processes*, *Oper. Res.*, 54 (2006), pp. 217–231.
- [9] I. J. CLARK, *Foreign Exchange Option Pricing: A Practitioner's Guide*, Wiley, New York, 2011.
- [10] J. COX, J. INGERSOLL, AND S. ROSS, *A theory of the term structure of interest rates*, *Econometrica*, 53 (1985), pp. 385–407.
- [11] A. COZMA, M. MARIAPRAGASSAM, AND C. REISINGER, *Calibration of a Hybrid Local-Stochastic Volatility Stochastic Rates Model with a Control Variate Particle Method*, preprint, <https://arxiv.org/abs/1701.06001>, 2018.
- [12] G. DEELSTRA AND F. DELBAEN, *Convergence of discretized stochastic (interest rate) processes with stochastic drift term*, *Appl. Stoch. Models Data Anal.*, 14 (1998), pp. 77–84.
- [13] G. DEELSTRA AND G. RAYEE, *Local volatility pricing models for long-dated FX derivatives*, *Appl. Math. Finance*, 20 (2013), pp. 380–402.
- [14] S. DEREICH, A. NEUENKIRCH, AND L. SZPRUCH, *An Euler-type method for the strong approximation of the Cox–Ingersoll–Ross process*, *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, 468 (2012), pp. 1105–1115.
- [15] S. S. DRAGOMIR, *Some Gronwall Type Inequalities and Applications*, Nova Science Publishers, Hauppauge, NY, 2003.
- [16] M. FISCHER AND G. NAPPO, *On the moments of the modulus of continuity of Itô processes*, *Stoch. Anal. Appl.*, 28 (2009), pp. 103–122.
- [17] M. B. GILES, D. J. HIGHAM, AND X. MAO, *Analysing multi-level Monte Carlo for options with non-globally Lipschitz payoff*, *Finance Stoch.*, 13 (2009), pp. 403–413.
- [18] P. GLASSERMAN, *Monte Carlo Methods in Financial Engineering*, *Stoch. Model. Appl. Probab.* 53, Springer, New York, 2003.
- [19] L. A. GRZELAK AND C. W. OOSTERLEE, *On the Heston model with stochastic interest rates*, *SIAM J. Financial Math.*, 2 (2011), pp. 255–286, <https://doi.org/10.1137/090756119>.
- [20] L. A. GRZELAK AND C. W. OOSTERLEE, *On cross-currency models with stochastic volatility and correlated interest rates*, *Appl. Math. Finance*, 19 (2012), pp. 1–35.
- [21] J. GUYON AND P. HENRY-LABORDÈRE, *The Smile Calibration Problem Solved*, SSRN 1885032, https://papers.ssrn.com/sol3/papers.cfm?abstract_id=1885032, 2011.
- [22] B. HAMBLY, M. MARIAPRAGASSAM, AND C. REISINGER, *A forward equation for barrier options under the Brunick & Shreve Markovian projection*, *Quant. Finance*, 16 (2016), pp. 827–838.
- [23] D. J. HIGHAM AND X. MAO, *Convergence of Monte Carlo simulations involving the mean-reverting square root process*, *J. Comput. Finance*, 8 (2005), pp. 35–62.
- [24] D. J. HIGHAM, X. MAO, AND A. M. STUART, *Strong convergence of Euler-type methods for nonlinear stochastic differential equations*, *SIAM J. Numer. Anal.*, 40 (2002), pp. 1041–1063, <https://doi.org/10.1137/S0036142901389530>.
- [25] T. R. HURD AND A. KUZNETSOV, *Explicit formulas for Laplace transforms of stochastic integrals*, *Markov Process. Rel. Fields*, 14 (2008), pp. 277–290.
- [26] A. S. HURN, K. A. LINDSAY, AND A. J. MCCLELLAND, *Estimating the parameters of stochastic volatility models using option price data*, *J. Bus. Econom. Statist.*, 33 (2014), pp. 579–594.
- [27] M. HUTZENTHALER, A. JENTZEN, AND P. E. KLOEDEN, *Strong and weak divergence in finite time of Euler's method for stochastic differential equations with non-globally Lipschitz continuous coefficients*, *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, 467 (2011), pp. 1563–1576.
- [28] M. HUTZENTHALER, A. JENTZEN, AND M. NOLL, *Strong Convergence Rates and Temporal Regularity for Cox–Ingersoll–Ross Processes and Bessel Processes with Accessible Boundaries*, preprint, <https://arxiv.org/abs/1403.6385v1>, 2014.
- [29] M. HUTZENTHALER, A. JENTZEN, AND X. WANG, *Exponential Integrability Properties of Numerical Approximation Processes for Nonlinear Stochastic Differential Equations*, preprint, <https://arxiv.org/abs/1309.7657v3>, 2016.
- [30] M. JEX, R. HENDERSON, AND D. WANG, *Pricing exotics under the smile*, *Risk*, 12 (1999), pp. 72–75.
- [31] C. KAHL AND P. JÄCKEL, *Fast strong approximation Monte Carlo schemes for stochastic volatility models*, *Quant. Finance*, 6 (2006), pp. 513–536.

- [32] P. KLOEDEN AND A. NEUENKIRCH, *Convergence of numerical methods for stochastic differential equations in mathematical finance*, in Recent Developments in Computational Finance: Foundations, Algorithms and Applications, T. Gerstner and P. Kloeden, eds., World Scientific, Hackensack, NJ, 2013, pp. 49–80.
- [33] P. E. KLOEDEN AND E. PLATEN, *Numerical Solution of Stochastic Differential Equations*, 3rd ed., Springer, New York, 1999.
- [34] A. LIPTON, *The vol smile problem*, Risk, 15 (2002), pp. 61–65.
- [35] A. LIPTON, A. GAL, AND A. LASIS, *Pricing of vanilla and first-generation exotic options in the local stochastic volatility framework: Survey and new results*, Quant. Finance, 14 (2014), pp. 1899–1922.
- [36] A. LIPTON AND W. MCGHEE, *Universal barriers*, Risk, 15 (2002), pp. 81–85.
- [37] R. LORD, R. KOEKKOEK, AND D. VAN DIJK, *A comparison of biased simulation schemes for stochastic volatility models*, Quant. Finance, 10 (2010), pp. 177–194.
- [38] A. NEUENKIRCH AND L. SZPRUCH, *First order strong approximations of scalar SDEs defined in a domain*, Numer. Math., 128 (2014), pp. 103–136.
- [39] S. NINOMIYA AND N. VICTOIR, *Weak approximation of stochastic differential equations and application to derivative pricing*, Appl. Math. Finance, 15 (2008), pp. 107–121.
- [40] Y. REN, D. MADAN, AND M. Q. QIAN, *Calibrating and pricing with embedded local volatility models*, Risk, 20 (2007), pp. 138–143.
- [41] R. SCHÖBEL AND J. ZHU, *Stochastic volatility with an Ornstein–Uhlenbeck process: An extension*, European Finance Rev., 3 (1999), pp. 23–46.
- [42] Y. TIAN, Z. ZHU, G. LEE, F. KLEBANER, AND K. HAMZA, *Calibrating and pricing with a stochastic-local volatility model*, J. Derivatives, 22 (2015), pp. 21–39.
- [43] A. VAN DER STOEP, L. A. GRZELAK, AND C. W. OOSTERLEE, *The Heston stochastic-local volatility model: Efficient Monte Carlo simulation*, Internat. J. Theoret. Appl. Finance, 17 (2014), pp. 1–30.
- [44] A. VAN HAASTRECHT, *Pricing Long-Term Options with Stochastic Volatility and Stochastic Interest Rates*, Ph.D. thesis, University of Amsterdam, Amsterdam, The Netherlands, 2010.
- [45] A. VAN HAASTRECHT, R. LORD, A. PELSSER, AND D. SCHRAGER, *Pricing long-maturity equity and FX derivatives with stochastic interest rates and stochastic volatility*, Insurance Math. Econom., 45 (2009), pp. 436–448.
- [46] A. VAN HAASTRECHT AND A. PELSSER, *Generic pricing of FX, inflation and stock options under stochastic interest rates and stochastic volatility*, Quant. Finance, 11 (2011), pp. 665–691.
- [47] U. WYSTUP, *FX Options and Structured Products*, Wiley, New York, 2007.