

# C2.6 Introduction to Schemes

Prof. Alexander F. Ritter  
University of Oxford  
HT 2020  
ritter@maths.ox.ac.uk

Feedback and corrections are welcome!

## EXERCISE SHEET 3

← Mostly topology, but useful

1) i) (Warm-up lemma)  $X$  topological space, check that  $\forall$  top-subspace  $Y \subseteq X$ :

$$Y \text{ irreducible} \implies Y \text{ connected}$$

$$Y \text{ irreducible} \implies \overline{Y} \text{ irreducible}$$

$$Y \text{ irreducible component} \implies Y \text{ closed and connected}$$

← recall: irred. component means irreducible and maximal w.r.t.  $\subseteq$

ii) Suppose  $X$  has finitely many irreducible components  $X_i$ .

Say " $X_k$  can be reached from  $X_\ell$ " if  $X_k \cap X_{i_1} \neq \emptyset, X_{i_1} \cap X_{i_2} \neq \emptyset, \dots, X_{i_n} \cap X_\ell \neq \emptyset$  some  $X_{i_r}$ .

Prove that  $X$  is connected  $\iff$  any irred. component can be reached from any other.

iii) A topological space is Noetherian if it satisfies the descending chain condition for closed sets:  $C_1 \supseteq C_2 \supseteq C_3 \supseteq \dots \implies C_N = C_{N+1} = \dots$  some  $N$ .

Prove: a Noeth. top. space has finitely many irreducible components, each containing an open dense set  $\neq \emptyset$

iv)  $R$  Noeth. ring  $\implies \text{Spec } R$  Noeth. top. space.

Check converse fails for  $k[x_1, x_2, x_3, \dots] / (x_1, x_2^2, x_3^3, \dots)$ .

← (so for a Noetherian scheme every affine open is Noeth. top. space)

v)  $X$  Noeth. top. space  $\iff$  every top. subspace of  $X$  is quasi-compact

← so for a Noeth. scheme  $X$  all subspaces are quasi-compact, not just  $X$ .

vi)  $X$  Noeth. scheme  $\implies X$  Noeth. top. space

2) i) Check  $\mathbb{A}_k^2 = \text{Spec } k[x, y]$  is a variety ( $k$  alg. closed field)

(Recall: variety = scheme which is integral, separated, finite type over  $\text{Spec } k$ .)

(Hint: you may assume as known that being "f.g.  $k$ -alg." is affine-local: notes Sec. 3.2)

ii) Show that the open subscheme  $\mathbb{A}_k^2 \setminus \{0\}$  is a variety which is not affine

iii) A variety which is affine ( $\text{Spec}(\text{ring})$ ) is an affine variety, i.e.  $\cong$  integral closed subscheme of  $\mathbb{A}_k^n$  (some  $n$ )

iv)  $(X, \mathcal{O}_X)$  variety  $\implies X$  Noetherian scheme

v) Glue two copies of  $\mathbb{A}_k^1 = \text{Spec } k[x]$  along the basic open set  $\mathbb{A}_k^1 \setminus 0 = D_x = \text{Spec } k[x, x^{-1}]$  by the isomorphism  $\text{Spec } k[s, s^{-1}] \xrightarrow{\cong} \text{Spec } k[t, t^{-1}]$  given by  $s \mapsto t$ .

Show that the glued scheme is not separated. ← (compare notes Sec. 5.3)

← (Hint: "equalizer")

vi) OPTIONAL EXERCISE  $(X, \mathcal{O}_X)$  variety,  $Z \subseteq X$  irreducible subspace ← (RMK irreducibility is not vital if allow varieties to be reducible.)

In notes Sec. 5.5 you find the definition of what it means for  $Z$  to be locally closed  $\subseteq$  scheme  $X$  and how we construct a canonical induced reduced scheme structure  $\mathcal{O}_Z$ .

• Prove  $Z$  locally closed  $\implies (Z, \mathcal{O}_Z)$  variety ← (Hint 2(iv), 1(vi), 1(v) may help)

• (harder) if you define  $\mathcal{O}_Z$  as suggested in Sec. 5.5 for  $Z \subseteq X$  irreducible subspace, prove that  $(Z, \mathcal{O}_Z)$  variety  $\implies Z \subseteq X$  locally closed

Suggestion first reduce to affine case  $Z = \text{Spec } S, X = \text{Spec } R$  by picking  $\text{Spec } R \subseteq X$  of type open/closed

Now want an open in  $Z$  s.t. generating global sections (over  $k$ ) come from sections on opens  $\subseteq X$ .

At the end, you may need to check  $\text{Spec } S \cap \text{Spec } R_f = \text{Spec } S_f \leftarrow (S_f = S \otimes_R R_f \text{ via } \varphi^\#: R \rightarrow S)$   
where  $A = \text{Spec } S \subseteq \text{Spec } R = B$   
 $\{x \in X : f(x) \neq 0 \in k(x)\}$

- 3)  $f: X \rightarrow B$  morph of schemes
- i)  $f$  is called an immersion (or locally closed immersion) if  $f: X \xrightarrow{\text{closed immersion}} U \xrightarrow{\text{open immersion}} B$
- Show that an immersion is a closed immersion  $\iff f(X) \subseteq B$  closed set  
(Hint. For  $\Leftarrow$ : glue the ideal sheaf of  $X \hookrightarrow U$  with  $\mathcal{O}_X|_{B \setminus f(X)}$ , check quasi-coherence)
- ii) Show  $\Delta_{X/B} \subseteq X \times_B X$  is closed if  $B, X$  affine  $\leftarrow$  (notation of notes Secs. 3)
- iii) Show  $\Delta_{X/B}$  immersion (Hence:  $f$  separated  $\iff \Delta_{X/B}$  closed imm.  $\iff \Delta_{X/B}$  closed set)
- iv) Call  $U, V \subseteq X$  "nice" if  $U, V, U \cap V$  affine opens and  $\mathcal{O}_X(U) \otimes_{\mathbb{Z}} \mathcal{O}_X(V) \xrightarrow{\text{surj.}} \mathcal{O}_X(U \cap V)$
- $f$  separated  $\implies (\forall$  affine open  $U, V \subseteq X$  with  $f(U), f(V) \subseteq$  affine open in  $B \implies U, V$  nice)
  - $(\exists$  open cover  $X = \cup U_i$  s.t.  $\forall x, y \in X$  with  $f(x) = f(y) \exists$  nice  $U_i, U_j \implies f$  separated with  $x \in U_i, y \in U_j$  and  $f(U_i), f(U_j) \subseteq$  affine open of  $B$ )

$\implies$  For  $B = \text{Spec } k: (\exists$  open cover  $X = \cup U_i$ , all  $U_i$  nice  $\implies f$  separated  $\implies$  (all affine opens  $U, V$  are nice)

- v) Show  $\mathbb{P}_k^n$  is separated by using (iv) ( $k$  any field). Deduce that  $\mathbb{P}_k^n$  is a variety.  
 Show any projective variety and quasi-projective variety are varieties  
 $\leftarrow$  integral closed subscheme of  $\mathbb{P}_k^n$   $\leftarrow$  irreducible open subsch. of a projective variety

4) i) Fact  $\mathbb{P}_k^n$  is complete (i.e. proper/ $k$ ). In this exercise work over alg. closed field  $k$ .

In notes, we showed  $\mathbb{A}^1$  is not complete because  $\mathbb{A}^1 \times \mathbb{A}^1 \supseteq \mathbb{V}(xy-1) \rightarrow \mathbb{A}^1$  fails the universally closed condition. Why is this not a problem for  $\mathbb{P}^1$  if consider  $\mathbb{P}^1 \times \mathbb{A}^1 \rightarrow \mathbb{A}^1$ ?

- ii)  $C \subseteq X$  closed subsc.,  $X$  complete  $\implies C$  complete  $\leftarrow$  (compare in topology: closed  $\subseteq$  compact is compact)  
 So Fact  $\implies$  also all projective varieties are complete.  $\leftarrow$  (use  $f_* \mathcal{O}_X$  on  $\text{Im}(f)$  to get scheme)
- iii)  $f: X \rightarrow Y, X$  universally closed,  $Y$  separated  $\implies \text{Im}(f) \subseteq Y$  closed & universally closed  $\leftarrow$  (compare topology: compact  $\rightarrow$  Hausdorff cts then image is closed & compact)
- iv)  $X$  complete variety  $\implies s \in \Gamma(X, \mathcal{O}_X)$  constant.  $\leftarrow$  (Hint graph)  $\leftarrow$  (Hint:  $\Gamma(X, \mathcal{O}_X) = \text{Mor}(X, \mathbb{A}^1)$  see sec 2.3 notes)
- v) Deduce that affine varieties are never complete, and that the only global sections of a projective variety  $X$  are constant morphisms  $X \rightarrow \mathbb{A}^1$ .

5) Note that any "commutative diagram" in a category  $\mathcal{C}$  can be thought of as a functor  $F: I \rightarrow \mathcal{C}$  where the objects of  $I$  are the positions  $i$  in the diagram (where you place some object  $F(i) = C_i \in \mathcal{C}$ ), the morphisms of  $I$  are the arrows of the diagram (together with all identity morphs  $i \rightarrow i$ , and composites)

"inverse limit":

The limit  $L = \varprojlim C_i \in \mathcal{C}$  (if exists) has morphs  $L \xrightarrow{\pi_i} C_i$  s.t.  $\left\{ \begin{array}{l} \text{compatible: } \forall (i \xrightarrow{\varphi} j) \in I: L \xrightarrow{\pi_i} C_i \xrightarrow{F(\varphi)} C_j \\ \text{universal property: } \forall \rho: L \rightarrow C_i \text{ s.t. } \rho \circ \pi_i = \rho \circ \pi_j \circ F(\varphi) \implies \exists! \sigma: L \rightarrow C_i \end{array} \right.$

"direct limit":

The colimit  $D = \varinjlim C_i$  is defined by reversing arrows  $\pi_i, \rho_i$  (so  $C_i \xrightarrow{\pi_i} D$ ).

EXAMPLE In sets,  $\varprojlim C_i = \{x_i \in \prod C_i : x_i \xrightarrow{F(i \rightarrow j)} x_j\}$ ,  $\varinjlim C_i = \bigsqcup C_i / \langle x_i \sim x_j \text{ if } x_i \xrightarrow{F(i \rightarrow j)} x_j \rangle$

- i) What is the functor of points interpretation of  $\varprojlim, \varinjlim$ ? (Hint: for  $\varinjlim$  consider  $I$  of  $\mathbb{P}^1$  and  $k^*$  not  $k^*$ )
- ii) Explain briefly why the product, fiber product, gluing of sheaves are limits, and the coproduct, pushout, gluing of schemes are colimits (e.g. every scheme =  $\varinjlim$  of its affine opens)

generate an equivalence e.g.  $x_i \sim x_j, x_j \sim x_k$  then declare  $x_i \sim x_k$

iii) Suppose  $f, g$  are adjoint functors  $(\text{Mor}_{\mathcal{D}}(fC, D) \rightarrow \text{Mor}_{\mathcal{C}}(C, gD))$  bijection, functorial in  $C, D$   
 Show that left adjoints commute with colimits, right adjoints commute with limits:  $\bullet g(\varinjlim C_i) = \varinjlim gC_i$   
 $\bullet f(\varprojlim C_i) = \varprojlim fC_i$