

Stability results for graphs with a critical edge

Alexander Roberts* Alex Scott*

February 3, 2017

Abstract

The classical stability theorem of Erdős and Simonovits states that, for any fixed graph with $\chi(H) = k + 1 \geq 3$, the following holds: every n -vertex graph that is H -free and has within $o(n^2)$ of the maximal possible number of edges can be made into the k -partite Turán graph by adding and deleting $o(n^2)$ edges. In this paper, we prove sharper quantitative results for graphs H with a critical edge, both for the Erdős-Simonovits Theorem (distance to the Turán graph) and for the closely related question of how close an H -free graph is to being k -partite. In many cases, these results are optimal to within a constant factor.

1 Introduction

For $n, k \geq 1$, the k -partite Turán graph $T_k(n)$ is the complete k -partite graph on n vertices with vertex classes as equal as possible (or equivalently the k -partite graph with maximum number of edges). We write $t_k(n) = e(T_k(n))$ for the number of edges in the Turán graph. A fundamental result in extremal graph theory is the Erdős-Simonovits Stability Theorem, which says that an H -free graph that is close to extremal must in fact look very much like a Turán graph.

Theorem 1.1 (Erdős-Simonovits [1]). *Let $k \geq 2$ and suppose that H is a graph with $\chi(H) = k + 1$. If G is an H -free graph with $e(G) \geq t_k(n) - o(n^2)$, then G can be formed from $T_k(n)$ by adding and deleting $o(n^2)$ edges.*

It is natural to ask how the $o(n^2)$ terms here depend on each other. Thus we will consider an H -free graph G with n vertices and $t_k(n) - f(n)$ edges, where $f(n) = o(n^2)$, and ask how close G is to the Turán graph $T_k(n)$.

In this paper, we will be interested in the case when H has a critical edge: an edge e in a graph H is said to be *critical* if $\chi(H - e) = \chi(H) - 1$. It was shown by Simonovits [14,

*Mathematical Institute, University of Oxford, Andrew Wiles Building, Radcliffe Observatory Quarter, Woodstock Road, Oxford, United Kingdom.
E-mail: {robertsa, scott}@maths.ox.ac.uk.

Theorem 2.3] that if H has a critical edge, then for sufficiently large n the Turán graph $T_k(n)$ is the unique extremal H -free graph on n -vertices. In this case, we will prove the following version of the Erdős-Simonovits Theorem.

Theorem 1.2. *Let H be a graph with a critical edge and $\chi(H) = k + 1 \geq 3$, and let $f(n) = o(n^2)$ be a positive integer. If G is an H -free graph with n vertices and $e(G) \geq t_k(n) - f(n)$, then G can be formed from $T_k(n)$ by adding and deleting $O(f(n)^{1/2}n)$ edges.*

It is easy to show that this bound is sharp up to a constant factor: if we take the Turán graph $T_k(n)$ and imbalance it by moving $\lceil f(n)^{1/2} \rceil$ vertices from one class to another then we obtain a k -partite graph G with n vertices and $t_k(n) - \Theta(f(n))$ edges. However, in order to obtain $T_k(n)$ from G we must change at least $\Omega(f(n)^{1/2}n)$ edges.

Theorem 1.2 will follow from a result on the closely related question: how many edges do we need to delete from G in order to make it k -partite? In a recent paper, Füredi [5] gave a beautiful proof of the following result of this type.¹

Theorem 1.3 (Füredi [5]). *Suppose that G is a K_{k+1} -free graph on n vertices with $t_k(n) - t$ edges. Then G can be made k -partite by deleting at most t edges.*

We will show that a much stronger bound holds. More generally, we will prove results for graphs H that contain a critical edge (note that every edge of K_{k+1} is critical. As we will see below, our bounds are sharp to within a constant factor for many graphs H (including K_{k+1}).

Theorem 1.4. *Let H be a graph with a critical edge and $\chi(H) = k + 1 \geq 3$, and let $f(n) = o(n^2)$ be a positive integer. If G is an H -free graph with n vertices and $e(G) \geq t_k(n) - f(n)$ then G can be made k -partite by deleting $O(n^{-1}f(n)^{3/2})$ edges.*

It is natural to ask what happens for small f . It follows from a result of Erdős and Simonovits [3, Theorem 1] that for sufficiently large n and $f(n) < \frac{n}{2k} - O(1)$, G is already k -partite. As we will see below, for many H the bound in Theorem 1.4 is optimal up to a constant factor; in many other cases we will be able to prove a stronger bound. In order to discuss this, we will need some definitions.

For disjoint sets A, B of vertices we write $K[A, B]$ for the edge set $\{ab : a \in A, b \in B\}$ of the complete bipartite graph $K_{A,B}$. For a graph $G = (V, E)$, recall that the *Mycielskian* [11] of G is a graph $M(G)$ with vertex set $\{v_i : v \in V, i = 1, 2\} \cup \{u\}$ and edge set $\{w_1v_i : wv \in E, i = 1, 2\} \cup \{v_2u : v \in V\}$. We define the *blown up Mycielskian graph* $M_k(a, b, c)$ as follows. Let V_1, \dots, V_k be sets of size a , let W_1, \dots, W_k be sets of size b , and let U be a set of size c (and let all these sets be disjoint). Then $M_k(a, b, c)$ has vertex set $\bigcup_{i=1}^k V_i \cup \bigcup_{i=1}^k W_i \cup U$ and edge set

$$\bigcup_{i \neq j} K[V_i, V_j] \cup \bigcup_{i \neq j} K[V_i, W_j] \cup \bigcup_i K[W_i, U].$$

¹We remark that Füredi also claims that Györi's work [6] implies the bound $O(f(n)^2n^{-2})$, but this is not correct (see Proposition 1.5).

Note that $M_k(a, b, c)$ is a blowup of the graph $M(K_k)$.




Figure 1: Mycielskian of K_3 ($M_3(1, 1, 1)$)




Figure 2: $M_3(1, 2, 2)$

If H has a critical edge but is not contained in a graph of form $M_k(a, a, a)$ then the bound in Theorem 1.4 is tight up to a constant factor.

Proposition 1.5. *Let H be a graph with a critical edge and $\chi(H) = k + 1 \geq 3$, and let $f(n) = o(n^2)$ be a positive integer. Suppose that H is not a subgraph of $M_k(a, a, a)$ for any a . Then there is an H -free graph with n vertices and $t_k(n) - O(f(n))$ edges which cannot be made k -partite by deleting $o(n^{-1}f(n)^{\frac{3}{2}})$ edges.*

If H is contained in some $M_k(a, a, a)$ then the construction used to prove Proposition 1.5 can no longer be used. However, we do have the following general lower bound that holds for all graphs H with a critical edge.

Proposition 1.6. *Let H be a graph with a critical edge and $\chi(H) = k + 1 \geq 3$, and let $f(n) = o(n^2)$ be a positive integer. Then there exists an H -free graph with n vertices and $t_k(n) - O(f(n))$ edges which cannot be made k -partite by deleting $o(n^{-2}f(n)^2)$ edges.*

Note that this is much weaker than the bound given in Proposition 1.5. However, if H is contained in $M_k(a, a, 1)$ for some a , then the bound in Proposition 1.6 is in fact tight to within a constant factor.

Theorem 1.7. *Let H be a graph with a critical edge and $\chi(H) = k + 1 \geq 3$, and suppose that H is a subgraph of $M_k(a, a, 1)$ for some a . Let $f(n) = o(n^2)$ be a positive integer. If G is an H -free graph on n vertices with $e(G) \geq t_k(n) - f(n)$ then G can be made k -partite by deleting $O(n^{-2}f(n)^2)$ edges.*

Theorems 1.4 and 1.7 give bounds that are sharp to within a constant factor when H is a subgraph of some $M_k(a, a, 1)$ and when H is not contained in any $M_k(a, a, a)$. What about graphs that are contained in some $M_k(a, a, a)$ but are not contained in any $M_k(a, a, 1)$? In this case, we do not have sharp results, but can say a little.

Theorem 1.8. *Let H be a graph with a critical edge and $\chi(H) = k + 1 \geq 3$, and suppose that H is a subgraph of $M_k(t, b, a)$. Let $f(n) = o(n^2)$ be a positive integer. If G is an H -free graph on n vertices such that $e(G) \geq t_k(n) - f(n)$ then G can be made k -partite by deleting $O(n^{\frac{1}{bk}}f(n)^{1-\frac{1}{bk}})$ edges.*

Note that the bound in Theorem 1.8 is stronger than the bound in Theorem 1.4 when $f(n) \gg n^{2-\frac{2}{5k+2}}$. This shows that the upper bound in Theorem 1.4 isn't tight for graphs that are contained in some $M_k(a, a, a)$ but are not contained in any $M_k(a, a, 1)$. In section 4, we will also give examples to show that such graphs need not satisfy the stronger bound from Theorem 1.7.

The paper is organised as follows. In Section 2 we give proofs of Theorems 1.2, 1.4, 1.7 and 1.8. In Section 3, we will prove Propositions 1.5 and 1.6 by way of constructions. In Section 4 we discuss the gap between the upper bound given by Theorem 1.8 and the lower bound given by Proposition 1.6, and then conclude the paper with some related problems and open questions.

Related results can be found in papers by Norin and Yepremyan [12, 15], and Pikhurko, Sliacan and Tyros [13]. Finally, we note that results from this paper are applied in a joint paper with Natasha Morrison [10].

2 Upper Bounds

In this section we present our proofs of Theorems 1.2, 1.4, 1.7 and 1.8. We start by recalling some important results that will be key to our argument. The first is the Erdős-Stone Theorem [4] concerning the extremal number of a complete symmetric k -partite graph.

Theorem 2.1 (Erdős-Stone [4]). *Let $k \geq 2$, $t \geq 1$, and $\epsilon > 0$. Then for n sufficiently large, if G is a graph on n vertices with*

$$e(G) \geq \left(1 - \frac{1}{k-1} + \epsilon\right) \binom{n}{2},$$

then G must contain a copy of $T_k(kt)$.

The second is a theorem proven by Simonovits regarding the extremal graph of a graph with a critical edge.

Theorem 2.2 (Simonovits [14, Theorem 2.3]). *Let H be a graph with a critical edge with $\chi(H) = k+1 \geq 3$. Then there exists some n_0 such that, for all $n \geq n_0$, we have $\text{Ex}(n; H) = \{T_k(n)\}$.*

As a stepping stone to Theorem 1.4, we first prove a weaker result. This can also be deduced from independent work of Norin and Yepremyan [12, 15, Theorem 3.1] but we include a self-contained proof here for completeness.

Lemma 2.3. *Let H be a graph with a critical edge and $\chi(H) = k+1 \geq 3$, and let $f(n) = o(n^2)$ be a positive integer. Suppose that G is an H -free graph on $[n]$ such that $e(G) \geq t_k(n) - f(n)$. Then G can be made k -partite by deleting $O(f(n))$ edges.*

The following easy lemma will be used repeatedly in the proof of Lemma 2.3.

Proposition 2.4. Fix $k \geq 2$ and $t \geq 1$. For $n \geq 1$, suppose $G \subset T_k(kn)$ is a $T_k(kt)$ -free graph. Then for sufficiently large n ,

$$e(G) \leq t_k(kn) - \frac{n^2}{2}. \quad (2.1)$$

Proof. Let $k \geq 2$ and $n, t \in \mathbb{N}$ and suppose $G \subset T_k(kn)$ is $T_k(kt)$ -free. Then $e(G) \leq \text{ex}(kn; T_k(kt))$, so we can apply Theorem 2.1 to get

$$\begin{aligned} t_k(kn) - e(G) &\geq t_k(kn) - \text{ex}(kn; T_k(kt)) \\ &\geq \frac{(kn)^2}{2} \left(1 - \frac{1}{k}\right) - \frac{(kn)^2}{2} \left(1 - \frac{1}{k-1} + o(1)\right) \\ &\geq \frac{n^2}{2}, \end{aligned}$$

for sufficiently large n . □

Proof of Lemma 2.3. Let $f(n) = o(n^2)$ be a positive integer, and let H be a graph with a critical edge and $\chi(H) = k + 1 \geq 3$. Choose t such that $H \subseteq T_k(tk) + e$, where e is any edge inside a vertex class of $T_k(tk)$. Let $\delta, \epsilon, \eta \in (0, \frac{1}{20000k^2})$ with $\epsilon \leq \frac{\eta^2}{8}$.

Suppose that G is an H -free graph on $[n]$ with $e(G) \geq t_k(n) - f(n)$. Since $f(n) = o(n^2)$, Theorem 1.1 tells us that there exists some N_0 such that when $n \geq N_0$, G is at most ϵn^2 edges away from a complete k -partite graph. By Theorem 2.2 we may assume that $\text{Ex}(n; H) = \{T_k(n)\}$ for all $n \geq N_0$.

Now suppose $n \geq 2N_0$ and let $L \subsetneq V(G)$ be the set of vertices with degree less than $(1 - \delta) \frac{n(k-1)}{k}$. Consider an arbitrary subset $B \subset L$ with $|B| < \frac{\delta n}{2}$ and let $J = G \setminus B$. We can count the number of edges in J by considering the number of edges removed from G to get that

$$e(J) \geq t_k(n) - f(n) - (1 - \delta)n \binom{k-1}{k} |B|. \quad (2.2)$$

Now $t_k(n) \geq t_k(n-1) + \frac{k-1}{k}(n-1)$, since we can form $T_k(n)$ from $T_k(n-1)$ by adding a vertex to a smallest vertex class, and so $t_k(n) \geq t_k(n-|B|) + \frac{k-1}{k}(n-|B|)|B|$. If we apply this inequality to (2.2), we see that

$$\begin{aligned} e(J) &\geq t_k(|J|) - f(n) + \binom{k-1}{k} (|B|(n-|B|) - (1-\delta)n|B|) \\ &= t_k(|J|) - f(n) + \binom{k-1}{k} (\delta n|B| - |B|^2) \\ &\geq t_k(|J|) - f(n) + \binom{k-1}{2k} \delta n|B|. \end{aligned} \quad (2.3)$$

On the other hand, J does not contain a copy of H and $|J| > n - \frac{\delta}{2}n > \frac{n}{2} \geq N_0$, so $e(J) \leq t_k(|J|)$. Comparing this upper bound with the lower bound given by (2.3), we see

that

$$|B| \leq \frac{2k}{\delta(k-1)} f(n)n^{-1}. \quad (2.4)$$

Recall that $f(n) = o(n^2)$ and so $\frac{2k}{\delta(k-1)} f(n)n^{-1} < \frac{\delta n}{4}$ for large enough n . Since B is an arbitrary subset of L with $|B| < \frac{\delta n}{2}$, we can conclude that $|L| < \frac{2k}{(k-1)\delta} f(n)n^{-1}$ for sufficiently large n .

Now fix $J = G \setminus L$. Since $|L| < \frac{2k}{(k-1)\delta} f(n)n^{-1}$, we have lost at most $\frac{2}{\delta} f(n) = O(f(n))$ edges, so it suffices to show that the graph J must be k -partite.

Let $q := |J| = (1 + o(1))n$. For sufficiently large n , the graph J has minimum degree at least $(1 - 2\delta)\frac{q(k-1)}{k}$, $e(J) = t_k(q) - O(f(n))$ and we already know that J is at most $\epsilon n^2 = \epsilon q^2(1 + o(1))$ edges away from being k -partite as a subgraph of G . We may then choose a partition V_1, \dots, V_k of $V(J)$ which contains at most ϵn^2 edges within the vertex classes. For n sufficiently large, since $f(n) = o(n^2)$, there are at most $\frac{3}{2}\epsilon n^2$ edges missing between the vertex classes. We now use this fact to derive information about the vertices in J .

Suppose some vertex v has at least ηn neighbours in each vertex class. Pick ηn neighbours of v in each vertex class to form $Q \subseteq V(J)$. Now let P be the subgraph of $J[Q]$ obtained by deleting all edges inside the classes $Q \cap V_i$. Note that if P contains a copy of $T_k(kt)$, then $J[Q \cup \{v\}]$ contains a copy of H , contradicting the fact that J is H -free. Therefore P is $T_k(tk)$ -free. An application of Proposition 2.4 then gives, for n sufficiently large,

$$e(P) \leq t_k(kn) - \frac{\eta^2}{2} n^2. \quad (2.5)$$

Since $\epsilon \leq \frac{\eta^2}{8}$, we get $e(P) \leq t_k(k\eta n) - 4\epsilon n^2$. But then at least $4\epsilon n^2$ edges between vertex classes are not present in J , which gives a contradiction. We may therefore assume that every vertex in J has at most ηn neighbours inside its own vertex class.

If $|V_i| - |V_j| \geq \frac{q}{50k}$ for some j , then $e(J) \leq t_k(q) - \frac{q^2}{10000k^2} + \epsilon q^2(1 + o(1)) < t_k(q) - \frac{\epsilon}{2} n^2$ for sufficiently large n , since $\epsilon < \frac{1}{20000k^2}$. This is impossible as $f(n) = o(n^2)$ and $e(J) = t_k(q) - O(f(n))$. So we may assume that $|V_i| \geq \frac{q}{k}(1 - \frac{1}{50})$ for each i .

Suppose, without loss of generality, that there is an edge uv inside V_1 . Consider the neighbourhoods of u and v in each of the vertex classes V_2, \dots, V_k . Note that $|\Gamma(u) \cap (V_2 \cup \dots \cup V_k)| \geq q(1 - 2\delta)\frac{(k-1)}{k} - 2\eta n$. At the same time $|V_2 \cup \dots \cup V_k| \leq q(1 - \frac{1}{k}(1 - \frac{1}{50}))$ and so $|(V_2 \cup \dots \cup V_k) \setminus \Gamma(u)| \leq q(\frac{1}{50k} + 2\delta + 2\eta) \leq \frac{q}{10k}$. The same argument applies for v and so there are most $\frac{q}{5k}$ vertices not in $\Gamma(u) \cap \Gamma(v)$ in each vertex class V_2, \dots, V_k . So $|\Gamma(u) \cap \Gamma(v) \cap V_i| \geq \frac{q}{k}(1 - \frac{1}{50}) - \frac{q}{5k} \geq \frac{q}{2k} \geq \eta n$ for each $i \geq 2$.

Pick $S_1 \subset V_1$ and $S_i \subset \Gamma(u) \cap \Gamma(v) \cap V_i$ for each $i = 2, \dots, k$ with $|S_i| = \eta n$ for each i . Let $Q = S_1 \cup \dots \cup S_k$ and P be the subgraph of $J[Q]$ obtained by deleting all edges inside the S_i . Arguing as in (2.5), we get $e(P) < t_k(k\eta n) - 4\epsilon n^2$ and so arrive at the same contradiction. Therefore there is no edge uv inside V_1 and so J must be k -partite as required. \square

We are now in a position to prove Theorem 1.4.

Proof of Theorem 1.4. Let $f(n) = o(n^2)$ be a positive integer, and let H be a graph with a critical edge and $\chi(H) = k + 1$. Choose t such that $H \subseteq T_k(tk) + e$, where e is any edge inside a vertex class of $T_k(tk)$. Let G be an H -free graph on $[n]$ with $e(G) \geq t_k(n) - f(n)$. Take a partition (V_1, \dots, V_k) of $V(G)$ which minimises the number of edges inside vertex classes. Then by Lemma 2.3, there are $O(f(n))$ edges within vertex classes and at most $O(f(n))$ edges between vertex classes are not present in G . Furthermore for each i , we must have $|V_i| = (1 + o(1))\frac{n}{k}$, otherwise G cannot contain enough edges.

Let v be a vertex in G with maximal number of neighbours inside its own vertex class. Without loss of generality, we may assume $v \in V_1$. Let $r(v) = |\Gamma(v) \cap V_1|$. By our choice of partition, v has at least $r(v)$ neighbours in every other vertex class. Pick $r(v)$ neighbours of v from each vertex class to form $Q \subseteq V(G)$ and let J be the subgraph of $G[Q]$ with all the edges within vertex classes removed (so J is k -partite). If J contains a copy of $T_k(tk)$, then by adding v to this copy, we must get a copy of H . J must then be $T_k(tk)$ -free and so we may apply Proposition 2.4 to get that if $r(v)$ is sufficiently large, then

$$e(J) \leq t_k(kr(v)) - \frac{r(v)^2}{2}. \quad (2.6)$$

As G is missing at most $O(f(n))$ edges between vertex classes, we must have $t_k(kr(v)) - e(J) = O(f(n))$ and so $r(v) = O(f(n)^{\frac{1}{2}})$.

Let $\delta \in (0, \frac{1}{4tk^2})$ and S be the set of vertices in $V(G)$ with degree less than $n(1 - \frac{1}{k})(1 - \delta)$. Arguing as in Lemma 2.3 (around (2.3)) we see that $|S| = O(f(n)n^{-1})$. We will show that each edge inside a vertex class is incident to a vertex in S .

Let E_I be the set of edges inside vertex classes. Pick some edge $e = uv \in E_I$ and suppose that neither u nor v is an element of S . Without loss of generality, assume that $u, v \in V_1$. Recall that $|V_i| = (1 + o(1))\frac{n}{k}$ for each i . Since $|\Gamma(u)| \geq n(1 - \frac{1}{k})(1 - \delta) \geq \frac{k-1}{k}n - \frac{n}{20k}$ and $|V_2 \cup \dots \cup V_k| = (1 + o(1))\frac{n(k-1)}{k}$, it follows that $|V_i \setminus \Gamma(u)| \leq \frac{n}{20k}(1 + o(1))$ for each $i \in \{2, \dots, k\}$. The same is true for v . So for sufficiently large n , $|V_i \cap \Gamma(u) \cap \Gamma(v)| \geq \frac{n}{2k}$ for each $i \in \{2, \dots, k\}$. Pick $B_1 \subset V_1$ and $B_i \subset \Gamma(u) \cap \Gamma(v) \cap V_i$ for each $i = 2, \dots, k$ with $|B_i| = \frac{n}{2k}$ for each i . If $Q = G[B_1 \cup \dots \cup B_k]$ contains a copy of $T_k(kt)$, then $G[Q \cup \{u, v\}]$ contains a copy of H , contradicting the fact that J is H -free. Therefore Q is $T_k(tk)$ -free. So by Proposition 2.4,

$$e(Q) \leq t_k\left(\frac{n}{2k}\right) - \frac{1}{8k^2}n^2.$$

This is a contradiction since G is missing $O(f(n))$ edges between vertex classes. Therefore u or v must belong to S .

We have shown that each edge in E_I is incident with S , every vertex of S is incident with at most $r(v) = O(f(n)^{\frac{1}{2}})$ edges from E_I , and $|S| = O(f(n)n^{-1})$. It follows that $|E_I| = O(f(n)^{\frac{3}{2}}n^{-1})$. \square

The proof of Theorem 1.4 was done in two parts. The first part was bounding the number of neighbours a vertex can have inside its respective vertex class by considering whether there

is a copy of $T_k(kt)$ in its neighbourhood. When we move to the regime of graphs contained in some $M_k(a, a, 1)$, we can improve the argument by considering whether there is a copy of $T_k(kt)$ present in many of the neighbourhoods of the vertex's neighbours.

Proof of Theorem 1.7. Let f be a function on the natural numbers such that $f(n) = o(n^2)$, let H be a graph with a critical edge and $\chi(H) = k + 1$ and suppose that h is such that $H \subset M_k(h, h, 1)$. Let G be an H -free graph on $[n]$ with $e(G) \geq t_k(n) - f(n)$. Let $\delta, \eta \in (0, \frac{1}{20000hk^2})$. Take a partition (V_1, \dots, V_k) of $V(G)$ which minimises the number of edges inside vertex classes and let E_I be the set of edges inside vertex classes. Furthermore, let $S = \{u \in V(G) : d(u) \leq (1 - \delta)\frac{n(k-1)}{n}\}$.

Carrying on from the end of the proof of Theorem 1.4, we know that each edge $e \in E_I$ is incident with a vertex in S and that $|S| = O(f(n)n^{-1})$. It therefore suffices to show that the maximum number of neighbours a vertex can have inside its own vertex class is $O(f(n)n^{-1})$.

Suppose without loss of generality that $v \in V_1$ has the maximum number of neighbours inside its own vertex class. Then for each i , let $A_i = V_i \cap \Gamma(v)$ and split each A_i into $B_i = S \cap A_i$ and $C_i = A_i \setminus B_i$. Let us consider the size of the C_i . Suppose that $|C_i| \geq h$ for each i and pick h -subsets $D_i \subset C_i$ for each i . Now for each $i \in [k]$, let

$$W_i = \left\{ x \in V_i \setminus (D_i \cup \{v\}) : \bigcup_{j \neq i} D_j \subset \Gamma(x) \right\}. \quad (2.7)$$

Note that for large enough n , each $u \in D_i$ is adjacent to all but at most $2\delta\frac{n}{k}$ vertices in V_j . Thus for large enough n , $|W_i| \geq \frac{n}{2k} - 2kh\delta\frac{n}{k} \geq \eta n$. So pick ηn vertices from each W_i to form a set Q of vertices and let J be the subgraph of $G[Q]$ with all edges inside vertex classes deleted. Note that if J contains a copy, $J[F]$, of $T_k(kh)$, then $G[\{v\} \cup \bigcup_{i \in [k]} D_i \cup F]$ will contain a copy of $M_k(h, h, 1)$ and so will contain a copy of H , a contradiction. So we may apply Proposition 2.4 to give that for n sufficiently large,

$$e(J) \leq t_k(k\eta n) - \frac{\eta^2}{2}n^2.$$

On the other hand, we know that there are $O(f(n))$ edges between vertex classes not present in G . Therefore, $e(J) \geq t_k(k\eta n) - O(f(n))$. We then have a contradiction since $f(n) = o(n^2)$. So there is a $j \in [k]$ such that $|C_j| < h$.

Now note since $B_j \subset L$, that $|B_j| = O(f(n)n^{-1})$ and so $|A_j| = O(f(n)n^{-1})$. Note that since we have taken the best partition with respect to edges inside vertex classes, $|A_1| \leq |A_j|$. We conclude that the maximum number of neighbours a vertex can have inside its own vertex class is $|A_1| = O(f(n)n^{-1})$. \square

Theorem 1.2 now follows as a direct corollary of Theorem 1.4.

Proof of Theorem 1.2. Let $f(n) = o(n^2)$ be a positive integer, and let H be a graph with a critical edge and $\chi(H) = k + 1$. Let G be an H -free graph on $[n]$ with $e(G) \geq t_k(n) - f(n)$. By theorem 1.4, we can delete $O(f(n)\frac{3}{2}n^{-1})$ edges from G to form a k -partite graph G' which

has $t_k(n) - O(f(n))$ edges. Suppose that V_1, \dots, V_k is a vertex colouring of G' . If the size of two colour classes differ by $2t$, then the maximum number of edges possible in the graph G' would be $t_k(n) - \Theta(t^2)$, and so two classes can differ in size by at most $O(f(n)^{\frac{1}{2}})$. It must then be the case that $||V_i| - \frac{n}{k}| = O(f(n)^{\frac{1}{2}})$ for each i and so a new graph G'' with equal class sizes can be formed by deleting the edges incident to $O(f(n)^{\frac{1}{2}})$ vertices. G'' has $t_k(n) - O(f(n)^{\frac{1}{2}}n)$ edges and has class sizes equal to that of the Turán graph. We can then attain the Turán graph by filling in the missing edges. To summarise, we deleted $O(f(n)^{\frac{1}{2}}n)$ edges to form G'' and then added $O(f(n)^{\frac{1}{2}}n)$ edges to then reach the Turán graph. \square

To end this section, we prove Theorem 1.8 using a method much like that used by Kóvari, Sós and Turán [9].

Proof of Theorem 1.8. Let $f(n) = o(n^2)$ be a positive integer, and let H be a graph with a critical edge and $\chi(H) = k + 1$. Further suppose that h, t and a are natural numbers such that H is contained in $M_k(t, h, a)$ and that H is contained in no $M_k(b, b, 1)$. Suppose that G is an H -free graph on n vertices such that $e(G) \geq t_k(n) - f(n)$. Take the best k -partition of G , W_1, \dots, W_k . By the proof of Theorem 1.4 we know that there is a set S of order $|S| = O(f(n)n^{-1})$ such that $G - S$ is a k -partite graph, that the minimum degree of the vertices in $V(G) \setminus S$ is $(1 + o(1))n^{\frac{k-1}{k}}$ and that the maximum number of edges inside a vertex class incident to a vertex in S is $O(f(n)^{\frac{1}{2}})$. We further know that there are $O(f(n))$ edges missing between any pair V_i, V_j , where $V_i = W_i \setminus S$.

Let E_I be the set of edges inside the vertex classes (so that G is $|E_I|$ edges away from being k -partite). We assume that $|E_I| = \Omega(f(n)^{1-\frac{1}{bk}}n^{\frac{1}{bk}})$, else we are done. Note that if we delete a different set of edges $F \subset E$ to obtain a k -partite subgraph of G , then since we have taken the best k -partition, it must be the case that $|F| \geq |E_I|$. So to get an upper bound on $|E_I|$ consider deleting all the edges between vertices in S , of which there are $O(f(n)^2n^{-2})$, and then deleting the edges between each vertex $s \in S$ and one of the V_i , where i may depend upon s . The best we could do (in terms of minimising edges deleted) by using this method is if for each $s \in S$ we deleted the edges between s and a V_i such that $|\Gamma(s) \cap V_i|$ is minimised. So if we let $e_I(s) = \min\{|\Gamma(s) \cap V_i| : i \in [k]\}$, we have an upper bound for the number of irregular edges in G .

$$|E_I| \leq \sum_{s \in S} e_I(s) + O(f(n)^2n^{-2}).$$

Recall that $|E_I| = \Omega(f(n)^{1-\frac{1}{bk}}n^{\frac{1}{bk}})$ and note that $f(n)^2n^{-2} = o(f(n)^{1-\frac{1}{bk}}n^{\frac{1}{bk}})$. This means that $\sum_{s \in S} e_I(s) = \Theta(|E_I|)$. Let $S' = \{s \in S : e_I(s) \geq 2h\}$. Since $|S| = O(f(n)n^{-1})$ the contribution to the sum of those vertices in $S \setminus S'$ is negligible and so

$$\sum_{s \in S'} e_I(s) = \Theta(|E_I|). \quad (2.8)$$

Following the argument of Theorem 1.7 from (2.7), we see that if we pick h -subsets D_i of each V_i , then there exists $C_i \in (V_i \setminus D_i)^{(t)}$ for each i such that $G[C_i, D_j]$ and $G[C_i, C_j]$

are homomorphic to $K_{t,h}$ and $K_{t,t}$ respectively for $i \neq j$. Therefore if there is an a -set A in S such that the vertices share h common neighbours in each V_i , then we can find a copy of $M_k(t, h, a)$ in G which contradicts our initial assumption that G is H -free. We will therefore count how many times an element of $V_1^{(h)} \times \dots \times V_k^{(h)}$ is contained within the neighbourhood of a vertex in S .

Since no element of $V_1^{(h)} \times \dots \times V_k^{(h)}$ can be contained within the neighbourhood of a distinct vertex in S , we have an upper bound given by $a|V_1^{(h)} \times \dots \times V_k^{(h)}| = O(n^{hk})$. On the other hand if we count over the vertices of S , we see that the neighbourhood of a vertex $s \in S$, contains

$$\prod_{i \in [k]} \binom{|\Gamma(s) \cap V_i|}{h} \geq \binom{e_I(s)}{h}^k \quad (2.9)$$

elements of $V_1^{(h)} \times \dots \times V_k^{(h)}$. Summing over the vertices in and comparing to the upper bound given earlier, we see that

$$\sum_{s \in S} \binom{e_I(s)}{h}^k = O(n^{hk}). \quad (2.10)$$

Note that we may easily bound the left hand side of (2.10) by summing only over S' and bounding $\binom{e_I(s)}{h}^k$ below by $(\frac{e_I(s)}{h})^{hk}$ to get that

$$\sum_{s \in S'} e_I(s)^{hk} = O(n^{hk}). \quad (2.11)$$

We can then bound the left hand side by applying Hölder's inequality to get that

$$\begin{aligned} \sum_{s \in S'} e_I(s)^{hk} &\geq |S'|^{1-hk} \left(\sum_{s \in S'} e_I(s) \right)^{hk} \\ &\geq (f(n)n^{-1})^{1-hk} \left(\sum_{s \in S'} e_I(s) \right)^{hk} \\ &= \Theta((f(n)n^{-1})^{1-hk} E_I^{hk}). \end{aligned} \quad (2.12)$$

Combining (2.11) and (2.12), we see that

$$(f(n)n^{-1})^{1-hk} E_I^{hk} = O(n^{hk}),$$

and so after rearranging, the result follows. \square

3 Lower Bounds

In this section, we prove Propositions 1.5 and 1.6.

Proof of Proposition 1.5. Fix $k \geq 2$ and let $f(n) = o(n^2)$ be a positive integer. Let $r = f(n)^{\frac{1}{2}}$ and $s = f(n)n^{-1}$. For n a large positive integer, consider the graph

$$G := M_k\left(\frac{n - s + kr}{k}, r, s\right).$$

Note that the numbers given for G may not be integer valued. This can easily be fixed but we have left it as it is for clarity and ease of reading (this will also be true of the remainder of this paper). Furthermore label subsets of the vertices of G as in Figure 3.




Figure 3: G

Now suppose that H is graph with a critical edge and $\chi(H) = k + 1$ and H is not a subgraph of $M_k(a, a, a)$ for any a . Thus H cannot be a subgraph of G .

We can obtain $T_k(n)$ from G by adding the edges between the W_i and changing the edges incident with U . In this process we add $O(f(n))$ edges and so $e(G) \geq t_k(n) - O(f(n))$. It is therefore enough to show that G is $\Omega(f(n)^{\frac{3}{2}}n^{-1})$ edges away from being k -partite.

So let Q be a k -partite subgraph on G formed by deleting edges from G . If we have deleted a fraction $\frac{1}{8k^2}$ of the edges between some pair (U, W_i) , (W_i, V_j) or (V_i, V_j) in forming Q from G , then in all cases we have deleted at least $\frac{f(n)^{\frac{3}{2}}n^{-1}}{16k^2}$ edges. Otherwise if we pick a vertex uniformly at random from each U, W_i and V_j to form a copy of $M_k(1, 1, 1)$ within G , then we expect to have deleted less than half an edge on average from this subgraph when forming Q (note that $e(M_k(1, 1, 1)) \leq 4k^2$). It must then be the case that we can pick a vertex from each U, W_i and V_j to form a copy of $M_k(1, 1, 1)$ in Q . This contradicts Q being k -partite and so there must be some pair (U, W_i) , (W_i, V_j) or (V_i, V_j) between which we have deleted a fraction $\frac{1}{8k^2}$ of the edges. In all cases we must have deleted $\Omega(f(n)^{\frac{3}{2}}n^{-1})$ edges from G and so G must be $\Omega(f(n)^{\frac{3}{2}}n^{-1})$ edges from being k -partite. \square

For critical graphs contained within some $M_k(a, a, a)$ we will consider graphs with chromatic number $k + 1$ such that every small subgraph has chromatic number at most k . We can construct an example of such a graph by adding more levels to the Mycielskian graph of a clique and blowing it up.

Definition 3.1. Let a, b, c, l and k be positive integers. Let V_1, \dots, V_k be sets of size a , let $W_1^1, \dots, W_k^1, \dots, W_1^l, \dots, W_k^l$ be sets of size b and let U be a set of size c (and let all these sets

be disjoint). Then the l -layer Mycielskian graph $M_k^{(l)}(a, b, c)$ has vertex set $V(G) \cup \bigcup_{i=1}^k V_i \cup \bigcup_{i \in [k], m \in [l]} W_i^m \cup U$ and edge set

$$\bigcup_{i \neq j} K[V_i, V_j] \cup \bigcup_{i \neq j} K[V_j, W_i^1] \cup \bigcup_{i \neq j, m \in [l-1]} K[W_i^m, W_j^{m+1}] \cup \bigcup_{i \in k} K[W_i^l, U].$$




Figure 4: $M_3^{(3)}(1, 2, 3)$

Proof of Proposition 1.6. Fix $k \geq 2$ and let $f(n) = o(n^2)$ be a positive integer. Let $s = f(n)n^{-1}$. Suppose that H is a graph with a critical edge on N vertices with $\chi(H) = k + 1$. For n a large positive integer, consider the graph G ,

$$G = M_k^{(N)}\left(\frac{n - (Nk + 1)s}{k}, s, s\right).$$

Furthermore label subsets of the vertices of G as in Figure 3 (we have drawn an example with $k = 4$).




Figure 5: G

Note that if we delete U or $\bigcup_i V_i$, then we are left with a k -partite and a bipartite graph respectively. It must then be the case that any subgraph $J \subset G$ with chromatic number $k + 1$ must contain vertices in both U and $\bigcup_i V_i$ and so must contain at least $N + 2$ vertices. It follows that G is an H -free graph since $|H| = N$.

We can obtain $T_k(n)$ from G by adding the edges between the W_i^j and changing the edges incident with U . In this process we add $O(f(n))$ edges and so $e(G) \geq t_k(n) - O(f(n))$.

Therefore if we can show that G is $\Omega(f(n)^2 n^{-2})$ edges away from being k -partite, then we will be done.

So let Q be a k -partite subgraph on G formed by deleting edges from G . If we have deleted a fraction $\frac{1}{8N^2 k^2}$ of the edges between some pair $(U, W_i^1), (W_i^l, W_j^{l+1}), (W_i^N, V_j)$ or (V_i, V_j) in forming Q from G , then in all cases we have deleted at least $\frac{f(n)^2 n^{-2}}{16N^2 k^2}$ edges. Otherwise if we pick a vertex uniformly at random from each U, W_i^l and V_j to form a copy of $M_k^{(N)}(1, 1, 1)$ within G , then we expect to have deleted less than half an edge on average from this subgraph when forming Q (Note that $e(M_k^{(N)}(1, 1, 1)) \leq 4N^2 k^2$). It must then be the case that we can pick a vertex from each $U, W_i^{(l)}$ and V_j to form a copy of $M_k^{(N)}(1, 1, 1)$ in Q . This contradicts Q being k -partite and so there must be some pair $(U, W_i^l), (W_i^l, W_i^{l+1})$ or (W_i, V_j) between which we have deleted a fraction $\frac{1}{8N^2 k^2}$ of the edges. In all cases we must have deleted $\Omega(f(n)^{\frac{3}{2}} n^{-1})$ edges from G and so G must be $\Omega(f(n)^2 n^{-2})$ edges from being k -partite. \square

4 Conclusion

We have given tight bounds for graphs with a critical edge that are not contained in any $M_k(a, a, a)$ and also graphs that are contained in some $M_k(a, a, 1)$; the remaining cases appear more difficult to handle. Theorem 1.8 shows that, for graphs contain in some $M_k(t, b, a)$, we can improve on the $O(f(n)^{\frac{3}{2}} n^{-1})$ upper bound of Theorem 1.4. The arguments used in the proof of Theorem 1.8 are rather crude, and it seems likely that stronger results should hold, at least when $f(n)$ is quite large. For instance, what can we say if $f(n) = n^{2-o(1)}$?

Let us also note that, for some graphs, we can improve on the $\Omega(f(n)^2 n^{-2})$ lower bound of Proposition 1.6. Let $r = f(n)^{\frac{1}{2}}$ and $s = f(n)n^{-1}$. Consider the graph given in Figure 3

$$G = M_k\left(\frac{n - s + kr}{k}, r, s\right),$$

and label the subsets $U, W_1, \dots, W_k, V_1, \dots, V_k$ as in Figure 3. Suppose we wanted to avoid a copy of $M_k(1, 1, 2)$, then it would be sufficient to changed the edges between U and $W_1 \cup \dots \cup W_k$ so that for each pair of vertices $u_1 \neq u_2 \in U$, there is a $j \in [k]$ so that u_1 and u_2 have no common neighbour in W_j .

Let $q = \lceil (\frac{f(n)}{n} + 1)^{\frac{1}{k}} \rceil$ and for each $i \in [k]$, let $W_i^{(0)}, \dots, W_i^{(q-1)}$ be disjoint subsets of W_i all of size $\lfloor \frac{f(n)^{\frac{1}{2}}}{q} \rfloor$. Now enumerate $U = \{u_1, \dots, u_r\}$ (where $r = \lceil f(n)n^{-1} \rceil$). Note that each $j \in [r]$ can be expressed as a q -ary number with k digits

$$j = \sum_{i \in [k]} a_i q^{i-1}, \tag{4.1}$$

where $a_i \in \{0, \dots, q-1\}$ for each i . Now create a new graph G' from G , where each vertex u_j in U has neighbourhood $\bigcup_{i \in [k]} W_i^{(a_i)}$ where the a_i are as in (4.1). The graph G' does not

contain $M_k(c, b, a)$ for any a, b and c with $a \geq 2$, and it is $\Theta(f(n)^{\frac{3}{2}-\frac{1}{k}}n^{-1+\frac{1}{k}})$ edges away from being k -partite. This improves a little upon the bound given by Proposition 1.6.

Finally, in this paper we have discussed graphs H with a critical edge. It would be interesting to get sharp results for all graphs both for the Erdős-Simonovits problem of distance from the Turán graph, and for the problem of the distance from being k -partite.

References

- [1] P. Erdős. Some recent results on extremal problems in graph theory. *Proc. Theory of Graphs Internl. Symp. Rome (1966)* 118-123.
- [2] P. Erdős and M. Simonovits. A limit theorem in graph theory. *Studia Sci. Math. Hungar* 1 1966 51-57.
- [3] P. Erdős and M. Simonovits. On a valence problem in extremal graph theory. *Discrete Mathematics* 5 (1973) 323-334.
- [4] P. Erdős and A. H. Stone. On the structure of linear graphs. *Bull. Amer. Math. Soc* 52 (1946), 1087-1091.
- [5] Z. Füredi. A proof of the stability of extremal graphs, Simonovits' stability from Szemerédi's regularity. *Journal of Combinatorial Theory, Series B* 115 (2015) 66-71
- [6] E. Győri. On the number of edge disjoint cliques in graphs of given size. *Combinatorica* 11 (3) (1991) 231-243
- [7] G. Hajós. Über eine Konstruktion nicht n -färbbarer Graphen. *Wiss. Z. Martin-Luther-Univ. Halle-Wittenberg Math.-Natur. Reihe* 10: 116-117
- [8] D. Hanson and B. Toft. k -saturated graphs of chromatic number at least k . *Ars Combin.* 31 (1991), 159-164
- [9] T. Kóvari, V. Sós and P. Turán. On a problem of K. Zarankiewicz. *Colloquium Mathematicae* 3.1 (1954): 50-57.
- [10] N. Morrison, A. Roberts and A. Scott. Maximising the number of cycles in graphs with forbidden subgraphs.
- [11] J. Mycielski. Sur le coloriage des graphes. *Colloq. Math.* 3 (1955): 161-162
- [12] S. Norin and L. Yepremyan. Turán numbers of extensions. *arXiv:1510.04689*
- [13] O. Pikhurko, J. Sliacan and K. Tyros. Strong forms of stability from flag algebra calculations.
- [14] M. Simonovits. Extremal graph problems with symmetrical extremal graphs. Additional chromatic conditions. *Discrete Mathematics* 7.3 (1974): 349-376.

[15] Liana Yepremyan, personal communication.