

Benjamini-Schramm local limits of sparse random planar graphs

Mihyun Kang


Joint work with Michael Misethan




Oxford Discrete Mathematics and Probability Seminar, 18 May 2021

Guiding questions/themes

(1) What does a random graph **locally** look like?



(2) How does a **global structure** of a random graph **affect** its **local structure**?



(3) What about a local structure of a random graph if a **global constraint** (e.g. planarity) is imposed to a random graph?

Part I.

Erdős-Rényi random graph

- $G = G(n, m) \in_R \mathcal{G}(n, m)$

a graph chosen uniformly at random from the class $\mathcal{G}(n, m)$ of all vertex-labelled simple graphs on vertex set $[n] := \{1, \dots, n\}$ with $m = m(n)$ edges

- all asymptotics are taken as $n \rightarrow \infty$

- **whp** = with high probability

- = with probability tending to one as $n \rightarrow \infty$


Phase transition in Erdős-Rényi random graph

$G = G(n, m) \in_R \mathcal{G}(n, m)$ and $2m/n \xrightarrow{n \rightarrow \infty} c \in [0, \infty)$

Theorem

[ERDŐS-RÉNYI 1959–60]

- If $c < 1$ ('subcritical'), whp
 - each component is of order $O(\log n)$
 - each component is either a tree or unicyclic component
- If $c > 1$ ('supercritical'), whp
 - unique largest ('giant') component of order $\Theta(n)$
 - largest comp contains \geq two cycles ('complex') and is not planar



Largest component in Erdős-Rényi random graph

$G = G(n, m) \in_R \mathcal{G}(n, m)$ and $2m/n \xrightarrow{n \rightarrow \infty} c \in [0, \infty)$


L largest component in G

ρ positive solution of $1 - \rho = \exp(-c \rho)$

Theorem

whp

$$|L| = (1 + o(1)) \rho n$$



Part II.


Local structure of Erdős-Rényi random graph

Part II.

Local structure of Erdős-Rényi random graph

● $r \in_R V(G(n, m))$

a vertex chosen uniformly at random from $V(G(n, m))$




$$d(r) \sim \text{Po}(c)$$

Part II.

Local structure of Erdős-Rényi random graph

● $r \in_R V(G(n, m))$

a vertex chosen uniformly at random from $V(G(n, m))$



$$d(r) \sim \text{Po}(c)$$

$$d(u) \sim \text{Po}(c)$$

$$d(v) \sim \text{Po}(c)$$

Benjamini–Schramm local weak limit

[BENJAMINI–SCHRAMM 2001; ALDOUS–STEELE 2004]


- a rooted graph is a pair (H, r) of a graph H and a vertex $r \in V(H)$
- two rooted graphs (H_1, r_1) and (H_2, r_2) are isomorphic,

$$(H_1, r_1) \cong (H_2, r_2)$$

if \exists isomorphism ϕ from H_1 onto H_2 with $\phi(r_1) = r_2$

- given a rooted graph (H, r) and $\ell \in \mathbb{N} := \{1, 2, \dots\}$, let

$$B_\ell(H, r) := H \left[\{v \in V(H) : d_H(v, r) \leq \ell\} \right]$$



Benjamini–Schramm local weak limit — cont'd

Definition


[BENJAMINI–SCHRAMM 2001; ALDOUS–STEELE 2004]

Given two **random** rooted graphs (G_1, r_1) and (G_2, r_2) with $G_1 = G_1(n)$, the **local weak limit** of (G_1, r_1) is (G_2, r_2) , denoted by

$$(G_1, r_1) \xrightarrow{d} (G_2, r_2)$$

if for each fixed rooted graph (H, r_H) and $\ell \in \mathbb{N}$

$$\mathbb{P} \left[B_\ell(G_1, r_1) \cong (H, r_H) \right] \xrightarrow{n \rightarrow \infty} \mathbb{P} \left[B_\ell(G_2, r_2) \cong (H, r_H) \right]$$



Erdős-Rényi random graph vs Galton-Watson tree


• $G = G(n, m) \in_R \mathcal{G}(n, m)$

• $r \in_R V(G)$


If $2m/n \xrightarrow{n \rightarrow \infty} c \in [0, \infty)$, then

$$(G, r) \xrightarrow{d} \text{GWT}(c)$$

where $\text{GWT}(c)$ is the Galton-Watson tree with offspring distribution $P_O(c)$



\xrightarrow{d}



i.e.

$$\mathbb{P}[B_\ell(G, r) \simeq (H, r_H)]$$

$\xrightarrow{n \rightarrow \infty}$

$$\mathbb{P}[B_\ell(\text{GWT}(c)) \simeq (H, r_H)]$$

Local weak limit of a random tree

• $T = T(n) \in_R \mathcal{T}(n)$

a tree chosen uniformly at random from the class of all trees
(i.e. acyclic connected graphs) on vertex set $[n]$


• $r \in_R V(T)$

Theorem

[GRIMMETT 1980/1981]

$$(T, r) \xrightarrow{d} T_\infty$$

Skeleton tree T_∞




a rooted tree obtained by replacing each vertex of an infinite path
by an independent Galton-Watson tree GWT (1)


Part III.

Local weak limit of a random planar graph

GWT(c) Galton–Watson tree



T_∞ Skeleton tree



Local weak limit of a random planar graph

- $P = P(n, m) \in_R \mathcal{P}(n, m)$ a uniform random planar graph
- $r \in_R V(P)$ a vertex chosen uniformly at random from $V(P)$


Local weak limit of a random planar graph

- $P = P(n, m) \in_R \mathcal{P}(n, m)$ a uniform random planar graph
- $r \in_R V(P)$ a vertex chosen uniformly at random from $V(P)$

Theorem

[K.-MISSETHAN 2021+]

- If $2m/n \xrightarrow{n \rightarrow \infty} c \in [0, 1]$, then $(P, r) \xrightarrow{d} \text{GWT}(c)$




Local weak limit of a random planar graph

- $P = P(n, m) \in_R \mathcal{P}(n, m)$ a uniform random planar graph
- $r \in_R V(P)$ a vertex chosen uniformly at random from $V(P)$

Theorem

[K.-MISSETHAN 2021+]

- If $2m/n \xrightarrow{n \rightarrow \infty} c \in [0, 1]$, then $(P, r) \xrightarrow{d} \text{GWT}(c)$
- If $2m/n \xrightarrow{n \rightarrow \infty} 2$, then $(P, r) \xrightarrow{d} T_\infty$



Local weak limit of a random planar graph


• $P = P(n, m) \in_R \mathcal{P}(n, m)$ and $2m/n \xrightarrow{n \rightarrow \infty} c \in (1, 2)$

• $r \in_R V(P)$

Theorem

[K.-MISSETHAN 2021+]

$$(P, r) \xrightarrow{d} (2 - c) \text{GWT}(1) + (c - 1) T_\infty$$



Local weak limit of a random planar graph


• $P = P(n, m) \in_R \mathcal{P}(n, m)$ and $2m/n \xrightarrow{n \rightarrow \infty} c \in (1, 2)$

• $r \in_R V(P)$

Theorem

[K.-MISSETHAN 2021+]

$$(P, r) \xrightarrow{d} (2 - c) \text{GWT}(1) + (c - 1) T_\infty$$



i.e. for each rooted graph (H, r_H) and $\ell \in \mathbb{N}$, we have

$$\mathbb{P}[B_\ell(P, r) \cong (H, r_H)] \xrightarrow{n \rightarrow \infty} (2 - c) \mathbb{P}[B_\ell(\text{GWT}(1)) \cong (H, r_H)] + (c - 1) \mathbb{P}[B_\ell(T_\infty) \cong (H, r_H)]$$

Part IV.

Main proof ideas

Phase transition in a random planar graph

$P = P(n, m) \in_R \mathcal{P}(n, m)$ and $2m/n \rightarrow c \in (1, 6]$


L largest component of P

Theorem

[K.-ŁUCZAK 2012]

● If $c \in (1, 2)$, whp $|L| = (1 + o(1)) (c - 1) n$



● If $c \in [2, 6]$, whp $|L| = (1 + o(1)) n$



Global structure of a random planar graph

$P = P(n, m) \in_R \mathcal{P}(n, m)$ and $2m/n \rightarrow c \in (1, 2)$

L largest component of P





Global structure of a random planar graph

$P = P(n, m) \in_R \mathcal{P}(n, m)$ and $2m/n \rightarrow c \in (1, 2)$

L largest component of P

$S = P \setminus L$ 'small' part of P

C 2-core = max subgraph of L with min deg \geq two



Global structure of a random planar graph

$P = P(n, m) \in_R \mathcal{P}(n, m)$ and $2m/n \rightarrow c \in (1, 2)$

L largest component of P



$S = P \setminus L$ 'small' part of P

C 2-core = max subgraph of L with min deg \geq two

Theorem


[K.-ŁUCZAK 2012; K.-MOSSHAMMER-SPRÜSSEL 2020]

- S 'behaves similarly' like a critical ER random graph $G(\bar{n}, \bar{m})$ with $\bar{n} = (2 - c)n$ and $2\bar{m}/\bar{n} \rightarrow 1$
- $L = C +$ each vertex in $V(C)$ replaced by a rooted tree



Local weak limit of a random forest

- $F = F(n, t) \in_R \mathcal{F}(n, t)$ a forest with t tree components
- $r_F \in_R V(F)$ a vertex chosen uniformly at random from $V(F)$



Local weak limit of a random forest


- $F = F(n, t) \in_R \mathcal{F}(n, t)$ a forest with t tree components
- $r_F \in_R V(F)$ a vertex chosen uniformly at random from $V(F)$
- r_T the root of the tree component T in F that contains r_F

Lemma

[K.-MISSETHAN 2021+]

If $t = t(n) = o(n)$, then whp $d_F(r_F, r_T) = \omega(1)$ and

$$(F, r_F) \xrightarrow{d} T_\infty$$



Finer view of local weak limits

$P = P(n, m) \in_R \mathcal{P}(n, m)$ and $2m/n \rightarrow c \in (1, 2)$


L largest component of P and $|L| \sim (c-1)n$

$r_L \in_R V(L)$

Theorem

[K.-MISSETHAN 2021+]

$$(L, r_L) \xrightarrow{d} T_\infty$$



Finer view of local weak limits

$P = P(n, m) \in_R \mathcal{P}(n, m)$ and $2m/n \rightarrow c \in (1, 2)$

L largest component of P and $|L| \sim (c-1)n$

$S = P \setminus L \sim$ **critical ER** random graph and $|S| \sim (2-c)n$


$r_S \in_R V(S)$, $r_L \in_R V(L)$

Theorem

[K.-MISSETHAN 2021+]

$(S, r_S) \xrightarrow{d} \text{GWT}(1)$

$(L, r_L) \xrightarrow{d} T_\infty$



Finer view of local weak limits

$P = P(n, m) \in_R \mathcal{P}(n, m)$ and $2m/n \rightarrow c \in (1, 2)$

L largest component of P and $|L| \sim (c-1)n$

$S = P \setminus L \sim$ **critical ER** random graph and $|S| \sim (2-c)n$

$r_S \in_R V(S)$, $r_L \in_R V(L)$, and $r_P \in_R V(P)$


Theorem

[K.-MISSETHAN 2021+]

$(S, r_S) \xrightarrow{d} \text{GWT}(1)$

$(L, r_L) \xrightarrow{d} T_\infty$

$(P, r_P) \xrightarrow{d} (2-c)\text{GWT}(1) + (c-1)T_\infty$




Local limit of a random planar graph with root in 2-core

$P = P(n, m) \in_R \mathcal{P}(n, m)$ and $2m/n \rightarrow c \in (1, 2)$

C 2-core = maximal subgraph of largest comp of P with min deg ≥ 2

$r_C \in_R V(C)$

$(P, r_C) \xrightarrow{d} T_\infty^{(2)}$



$T_\infty^{(k)}$ a rooted tree obtained by replacing each vertex of k infinite paths rooted at a common vertex by an independent GWT (1)


Local limit of a random planar graph with root in 2-core

$P = P(n, m) \in_R \mathcal{P}(n, m)$ and $2m/n \rightarrow c \in (1, 2)$

C 2-core = maximal subgraph of largest comp of P with min deg ≥ 2

$r_C \in_R V(C)$

$$(P, r_C) \xrightarrow{d} T_\infty^{(2)}$$



$T_\infty^{(k)}$ a rooted tree obtained by replacing each vertex of k infinite paths rooted at a common vertex by an independent GWT (1)


cf. local weak limit of a random planar graph with $r \in_R V(P)$

$$\begin{aligned} (P, r) &\xrightarrow{d} (2-c) \text{GWT}(1) + (c-1) T_\infty \\ &= (2-c) T_\infty^{(0)} + (c-1) T_\infty^{(1)} \end{aligned}$$


Summary and open problems

(1) Phase transitions and critical phenomena

Uniform random graph $G(n, m)$




Random planar graph $P(n, m)$




Summary and open problems

(1) Phase transitions and critical phenomena

Uniform random graph $G(n, m)$



Random planar graph $P(n, m)$




- $S = G(n, m) \setminus L$ 'behaves similarly' like a **subcritical** ER random graph $G(\bar{n}, \bar{m})$ with $\bar{n} = (1 - \rho)n$ and $2\bar{m}/\bar{n} < 1$
- $S = P(n, m) \setminus L$ 'behaves similarly' like a **critical** ER random graph $G(\bar{n}, \bar{m})$ with $\bar{n} = (2 - c)n$ and $2\bar{m}/\bar{n} \rightarrow 1$

Summary and open problems — cont'd

(2) Local weak limit of a random planar graph

If $P = P(n, m) \in_R \mathcal{P}(n, m)$, $2m/n \rightarrow c \in (1, 2)$, and $r \in_R V(P)$, then

$$(P, r) \xrightarrow{d} (2-c) \text{GWT}(1) + (c-1) T_\infty$$




Summary and open problems — cont'd

Let $P = P(n, m) \in_R \mathcal{P}(n, m)$ and $r \in_R V(P)$

(3) In 2nd critical regime when $2m/n \rightarrow 2$

$$|2\text{-core}| = o(n) \quad \text{and} \quad (P, r) \xrightarrow{d} T_\infty = T_\infty^{(1)}$$




Summary and open problems — cont'd

Let $P = P(n, m) \in_R \mathcal{P}(n, m)$ and $r \in_R V(P)$

(3) In 2nd critical regime when $2m/n \rightarrow 2$ and $m \leq n + o(n(\log n)^{-2/3})$

$$|2\text{-core}| = o(n) \quad \text{and} \quad (P, r) \xrightarrow{d} T_\infty = T_\infty^{(1)}$$




Summary and open problems — cont'd

Let $P = P(n, m) \in_{\mathcal{R}} \mathcal{P}(n, m)$ and $r \in_{\mathcal{R}} V(P)$

(3) In 2nd critical regime when $2m/n \rightarrow 2$ and $m \leq n + o(n(\log n)^{-2/3})$

$$|2\text{-core}| = o(n) \quad \text{and} \quad (P, r) \xrightarrow{d} T_{\infty} = T_{\infty}^{(1)}$$



(4) Conjecture: $\exists 2 < t_c < 4.42$ and $0 < a, b < 1$ s.t. $2m/n \rightarrow \beta \in (2, t_c)$,

• $|2\text{-core}| = (a + b + o(1))n$, $|\text{kernel}| = (b + o(1))n$, and

• $(P, r) \xrightarrow{d} (1 - a - b)T_{\infty}^{(1)} + aT_{\infty}^{(2)} + bT_{\infty}^{(\geq 3)}$