

The Ramsey number of books

David Conlon*

Abstract

We show that in every two-colouring of the edges of the complete graph K_N there is a monochromatic K_k which can be extended in at least $(1 + o_k(1))2^{-k}N$ ways to a monochromatic K_{k+1} . This result is asymptotically best possible, as may be seen by considering a random colouring. Equivalently, defining the book $B_n^{(k)}$ to be the graph consisting of n copies of K_{k+1} all sharing a common K_k , we show that the Ramsey number $r(B_n^{(k)}) = 2^k n + o_k(n)$. In this form, our result answers a question of Erdős, Faudree, Rousseau and Schelp and establishes an asymptotic version of a conjecture of Thomason.

1 Introduction

The Ramsey number $r(H)$ of a graph H is the smallest natural number N such that every two-colouring of the edges of the complete graph K_N contains a monochromatic copy of H . The problem of determining Ramsey numbers is notoriously hard. For instance, when H is a complete graph, work of Erdős and Szekeres [8, 12] in the 1930s and 40s showed that $\sqrt{2}^t \leq r(K_t) \leq 4^t$, but the only improvements to these bounds since that time [3, 24] have been to lower order terms.

We investigate the Ramsey numbers of books, a study which bears close relation to the problem of determining $r(K_t)$. The book $B_n^{(k)}$ is the graph consisting of n copies of K_{k+1} , all sharing a common K_k . Embracing the metaphor, we refer to the common K_k as the spine of the book and the n points completing each copy of K_{k+1} as the pages or leaves.

The Ramsey problem for these books was first studied by Erdős, Faudree, Rousseau and Schelp [10] and then by Thomason [25]. Both papers contain bounds of the form

$$2^k n + o_k(n) \leq r(B_n^{(k)}) \leq 4^k n,$$

where the lower bound follows from considering the random graph $G(n, 1/2)$ and the upper bound from a standard neighbourhood chasing argument. In their paper, Erdős et al. asked whether one of these bounds might be asymptotically correct and Thomason conjectured that the lower bound is. In fact, he made a very precise conjecture about the value of $r(B_n^{(k)})$, namely, that

$$r(B_n^{(k)}) \leq 2^k(n + k - 2) + 2.$$

For $k = 2$, this conjecture is known to hold [23] and is tight for infinitely many values of n . The main contribution of this paper is a proof of an approximate version of Thomason's conjecture, thus answering the question of Erdős et al. (see also [21]).

*Mathematical Institute, Oxford OX2 6GG, United Kingdom. Email: david.conlon@maths.ox.ac.uk. Research supported by a Royal Society University Research Fellowship and ERC Starting Grant 676632.

Theorem 1 *For every natural number k ,*

$$r(B_n^{(k)}) = 2^k n + o_k(n).$$

To see something of why this is interesting, suppose that we have a red/blue-colouring of K_N with no monochromatic copy of K_t . Then this colouring contains no monochromatic book $B_n^{(k)}$ with n equal to the off-diagonal Ramsey number $r(K_{t-k}, K_t)$. Indeed, suppose that the book is red. If the set induced by the pages contains a blue K_t , we have a contradiction, so it must contain a red K_{t-k} . But together with the red spine K_k , this forms a red K_t . But, by Theorem 1, if t , and therefore n , is sufficiently large in terms of k , we have

$$r(K_t) \leq r(B_n^{(k)}) \leq 2^{k+1} r(K_{t-k}, K_t) \leq 2^{k+1} \binom{2t-k}{t-k},$$

where the last inequality follows from a classical estimate of Erdős and Szekeres [12]. In particular, if the theorem applied for t linear in k , this would give an exponential improvement on the upper bound for diagonal Ramsey numbers.

Another motivation for Theorem 1 is its relation to a well-known, but false, conjecture of Erdős [9] (see also [2]) asserting that every two-colouring of the edges of K_N contains at least

$$(1 + o_k(1)) 2^{1 - \binom{k}{2}} \binom{N}{k}$$

monochromatic copies of K_k . That is, he conjectured that a random colouring should asymptotically minimise the number of monochromatic copies of K_k . While true for $k = 3$ by a result of Goodman [16], this conjecture was disproved for $k \geq 4$ by Thomason [26]. However, Theorem 1 is equivalent to a local version of Erdős' conjecture, saying that there is some monochromatic copy of K_{k-1} which is contained in asymptotically as many monochromatic K_k as in a random colouring. In some ways, this interpretation is more appealing than the original formulation in terms of books. It also connects our result with the study of Ramsey multiplicity, which has drawn considerable attention in recent years (see, for instance, [4, 7, 13, 19, 20]).

2 Preliminaries

In this section, we collect several results that we will need for the proof of Theorem 1.

2.1 Regularity and counting lemmas

One of the main ingredients in our proof is a simple corollary of Szemerédi's regularity lemma. To state this fundamental result, we first recall some standard definitions. A bipartite graph between two sets U and V is said to be ϵ -regular if, for all sets $U' \subseteq U$, $V' \subseteq V$ with $|U'| \geq \epsilon|U|$ and $|V'| \geq \epsilon|V|$, $|d(U', V') - d(U, V)| \leq \epsilon$. A partition $V(G) = \cup_{i=1}^m V_i$ of the vertex set of a graph G is said to be equitable if $||V_i| - |V_j|| \leq 1$ for all i and j . The regularity lemma is now as follows.

Lemma 1 *For every $0 < \epsilon < 1$ and natural number m_0 , there exists a natural number M such that every graph G with at least m_0 vertices has an equitable partition $V(G) = \cup_{i=1}^m V_i$ with $m_0 \leq m \leq M$ parts such that all but ϵm^2 pairs (V_i, V_j) with $1 \leq i \neq j \leq m$ are ϵ -regular.*

We will also need the following lemma from [5]. We say that a subset U of the vertex set of a graph G is ϵ -regular if the pair (U, U) is ϵ -regular.

Lemma 2 *For every $0 < \epsilon < 1$, there exists a constant δ such that every graph contains an ϵ -regular vertex subset U with $|U| \geq \delta|V(G)|$.*

The key lemma we will need is the following. We note a superficial similarity to a lemma used in the proof of the induced removal lemma [1], though that lemma requires the stronger condition that every pair (W_i, W_j) be regular.

Lemma 3 *For every $0 < \eta < 1$ and natural number m_0 , there exists a natural number M such that every graph G with at least m_0 vertices has an equitable partition $V(G) = \cup_{i=1}^m V_i$ with $m_0 \leq m \leq M$ parts and subsets $W_i \subseteq V_i$ such that W_i is η -regular for all i and, for all but ηm^2 pairs (i, j) with $1 \leq i \neq j \leq m$, (V_i, V_j) , (W_i, V_j) and (W_i, W_j) are η -regular with $|d(W_i, V_j) - d(V_i, V_j)| \leq \eta$ and $|d(W_i, W_j) - d(V_i, V_j)| \leq \eta$.*

Proof: Apply the regularity lemma to G with $\epsilon = \eta \cdot \delta(\eta)$, with δ as in Lemma 2. This yields an equitable partition $V(G) = \cup_{i=1}^m V_i$ where all but $\epsilon m^2 \leq \eta m^2$ pairs (V_i, V_j) with $1 \leq i \neq j \leq m$ are ϵ -regular. Within each piece V_i , now apply Lemma 2 to find a set W_i of order at least $\delta(\eta)$ which is η -regular. Note that if (V_i, V_j) is ϵ -regular, then, since $|W_i| \geq \delta|V_i|$ and $\epsilon = \eta \cdot \delta(\eta)$, the pairs (W_i, V_j) and (W_i, W_j) are η -regular with $|d(W_i, V_j) - d(V_i, V_j)| \leq \epsilon \leq \eta$ and $|d(W_i, W_j) - d(V_i, V_j)| \leq \eta$. \square

In order to apply Lemma 3, we need a standard counting lemma (see, for example, [22, Theorem 18]).

Lemma 4 *For any $\delta > 0$, there is $\eta > 0$ such that if U_1, \dots, U_k are (not necessarily distinct) vertex sets with (U_i, U_j) η -regular of density d_{ij} for all $1 \leq i < j \leq k$, then there are*

$$\prod_{i < j} d_{ij} \prod_{i=1}^k |U_i| \pm \delta \prod_{i=1}^k |U_i|$$

copies of K_k with vertex i in U_i for each $1 \leq i \leq k$.

In practice, we will always use this lemma in the following form.

Lemma 5 *For any $\delta > 0$, there is $\eta > 0$ such that if $U_1, \dots, U_k, U_{k+1}, \dots, U_{k+\ell}$ are (not necessarily distinct) vertex sets with (U_i, U_j) η -regular of density d_{ij} for all $1 \leq i < j \leq k + \ell$ and $d_{ij} \geq \delta$ for all $1 \leq i < j \leq k$, then there is a copy of K_k with vertex i in U_i for each $1 \leq i \leq k$ which is contained in at least*

$$\sum_{s=1}^{\ell} \left(\prod_{i=1}^k d_{i, k+s} - \delta \right) |U_{k+s}|$$

copies of K_{k+1} with vertex $k+1$ in $\cup_{s=1}^{\ell} U_{k+s}$.

Proof: By Lemma 4 applied with $\delta' = \delta \binom{k}{2} + 1/2$ instead of δ , there exists η such that the number of copies of K_k with vertex i in U_i for each $1 \leq i \leq k$ is at most

$$\prod_{1 \leq i < j \leq k} d_{ij} \prod_{i=1}^k |U_i| + \delta' \prod_{i=1}^k |U_i|.$$

Moreover, the number of copies of K_{k+1} with vertex i in U_i for each $1 \leq i \leq k$ and vertex $k+1$ in $\cup_{s=1}^{\ell} U_{k+s}$ is at least

$$\sum_{s=1}^{\ell} \prod_{i=1}^k d_{i,k+s} |U_{k+s}| \prod_{1 \leq i < j \leq k} d_{ij} \prod_{i=1}^k |U_i| - \delta' \sum_{s=1}^{\ell} |U_{k+s}| \prod_{i=1}^k |U_i|.$$

Therefore, there must be some K_k which is in at least

$$\begin{aligned} \frac{\sum_{s=1}^{\ell} \prod_{i=1}^k d_{i,k+s} |U_{k+s}| \prod_{1 \leq i < j \leq k} d_{ij} - \delta' \sum_{s=1}^{\ell} |U_{k+s}|}{\prod_{1 \leq i < j \leq k} d_{ij} + \delta'} &\geq \frac{\sum_{s=1}^{\ell} \left(\prod_{i=1}^k d_{i,k+s} - \delta^{-\binom{k}{2}} \delta' \right) |U_{k+s}|}{1 + \delta^{-\binom{k}{2}} \delta'} \\ &\geq \sum_{s=1}^{\ell} \left(\prod_{i=1}^k d_{i,k+s} - \delta \right) |U_{k+s}| \end{aligned}$$

copies of K_{k+1} , as required. \square

2.2 A coloured extremal result

We will need a coloured version of the celebrated Erdős–Stone theorem [11]. Recall that a blow-up of a graph H is a graph where each vertex of H is replaced by a vertex set and the bipartite graph between two such vertex sets is complete whenever the corresponding vertices are joined by an edge.

Lemma 6 *For any natural numbers k and t and any $\delta > 0$, there exists a natural number n_0 such that if the edges of the complete graph on $n \geq n_0$ vertices are coloured in red and blue, then, provided the blue density is at least $1 - \frac{1}{k-1} + \delta$, there is a blue blow-up of K_k with t vertices in each part, where each part is itself a monochromatic clique.*

Proof: Since the blue density is at least $1 - \frac{1}{k-1} + \delta$, the Erdős–Stone theorem implies that for $n \geq n_0$ there is a blue blow-up of K_k with at least $r(K_t)$ vertices in each part. Applying Ramsey’s theorem within each part then gives the required monochromatic cliques. \square

In practice, we will use a slight variant of this lemma, where the underlying graph is not necessarily complete.

Lemma 7 *For any natural numbers k and t and any $\delta > 0$, there exists a natural number n_1 and $\epsilon > 0$ such that if the edges of a graph on $n \geq n_1$ vertices with $(1 - \epsilon) \binom{n}{2}$ edges are coloured in red and blue, then, provided the blue density is at least $1 - \frac{1}{k-1} + \delta$, there is a blue blow-up of K_k with t vertices in each part, where each part is itself a monochromatic clique.*

Proof: Suppose first that the $\epsilon \binom{n}{2}$ missing edges are coloured blue, so that the underlying graph is complete. Then, by Lemma 6, every subset of the graph of order n_0 contains the required blow-up of K_k with monochromatic parts of order t . But then, for n sufficiently large in terms of n_0 , there must be at least

$$\binom{n}{n_0} / \binom{n-kt}{n_0-kt} = \binom{n}{kt} / \binom{n_0}{kt} \geq \frac{n^{kt}}{2n_0^{kt}}$$

such blow-ups of K_k . However, at most $\epsilon n^2 \cdot n^{kt-2} = \epsilon n^{kt}$ such copies contain an edge from the missing set. Therefore, for $\epsilon < 1/2n_0^{kt}$, we must have the required blue blow-up of K_k with at least t vertices in each part, where each part is a monochromatic clique. \square

2.3 Some technical lemmas

The proof requires a small degree of optimisation, almost all of which is contained in the following two lemmas.

Lemma 8 *For each $i = 1, \dots, k$, let x_i be a real number between 0 and t . Then either*

$$\sum_{i=1}^k (t - x_i)^k \geq (t/2)^k k \quad \text{or} \quad \prod_{i=1}^k x_i \geq (t/2)^k.$$

Proof: We claim that the minimum value of $\sum_i (t - x_i)^k$ subject to the constraint $\prod_i x_i = (t/2)^k$ occurs when all the x_i are equal to $t/2$. Since decreasing the product can only increase the sum, this will suffice to prove the result. As the claim is easily checked for $k = 2$ and 3 , we can assume without loss of generality that $k \geq 4$. But then we may assume that none of the x_i are less than t/k , since $(t - t/k)^k > (t/2)^k k$ for $k \geq 4$.

Now make the substitution $x_i = e^{y_i}$. The problem then becomes to minimise $\sum_{i=1}^k (t - e^{y_i})^k$ subject to the constraint $\sum_{i=1}^k y_i = k \log(t/2)$. But the function $(t - e^y)^k$ is easily seen to be convex for $t/k \leq e^y \leq t$. Therefore, the minimum occurs when all of the e^{y_i} and, hence, all of the x_i are equal. \square

Lemma 9 *Suppose that $k \leq \ell$ and, for each $i = 1, \dots, \ell$, let x_i be a real number between 0 and 1. Then*

$$\sum_{1 \leq i_1 < \dots < i_k \leq \ell} \prod_{j=1}^k x_{i_j} \geq \binom{\sum_i x_i}{k}.$$

Proof: Suppose that $\sum_i x_i = c$ and we wish to minimise the left-hand side of the required inequality under this constraint. Consider the function $f(x_1, \dots, x_{k-1})$ obtained by replacing x_ℓ with $c - \sum_{i=1}^{k-1} x_i$ in

$$\sum_{1 \leq i_1 < \dots < i_k \leq \ell} \prod_{j=1}^k x_{i_j}.$$

For each of the variables x_i with $1 \leq i \leq k-1$, this function has the form

$$f = -Ax_i^2 + Bx_i + C,$$

where A is a function of $x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_{k-1}$ which is non-negative on the allowed domain. Hence, if we fix all variables other than x_i , f is minimised when x_i is either as large or as small as is allowed by the constraints $0 \leq x_i \leq 1$ and $x_i \leq c - \sum_{j \neq i, \ell} x_j \leq x_i + 1$. Using symmetry, we can now check that the minimum is obtained when $\lfloor c \rfloor$ of the x_i are 1, one of the x_i is $\{c\}$ and the remainder are 0. Therefore,

$$\sum_{1 \leq i_1 < \dots < i_k \leq \ell} \prod_{j=1}^k x_{i_j} \geq \binom{\lfloor c \rfloor}{k} + \{c\} \binom{\lfloor c \rfloor}{k-1} \geq \binom{c}{k}.$$

To establish the final inequality, suppose that X is a random subset of a $(\lfloor c \rfloor + 1)$ -element set, where the first element is chosen with probability $\{c\}$ and all other elements with probability 1. The expected number of subsets of size k in this random set is then

$$\binom{\lfloor c \rfloor}{k} + \{c\} \binom{\lfloor c \rfloor}{k-1}.$$

But it is also equal to

$$(1 - \{c\}) \binom{\lfloor c \rfloor}{k} + \{c\} \binom{\lfloor c \rfloor + 1}{k},$$

which by convexity of $\binom{x}{k}$ is at least $\binom{c}{k}$. \square

3 Proof of Theorem 1

Suppose that we have a red/blue-colouring of the edges of the complete graph on $N = (2^k + \epsilon)n$ vertices. Assume that η is taken sufficiently small and m_0 sufficiently large in terms of k and ϵ and apply Lemma 3 with η and m_0 to the red subgraph to obtain an equitable partition $\cup_{i=1}^m V_i$ of the vertex set $[N]$ with $m \geq m_0$ and subsets $W_i \subseteq V_i$ such that W_i is η -regular for all i and, for all but ηm^2 pairs (i, j) with $1 \leq i \neq j \leq m$, (V_i, V_j) , (W_i, V_j) and (W_i, W_j) are η -regular with $|d(W_i, V_j) - d(V_i, V_j)| \leq \eta$ and $|d(W_i, W_j) - d(V_i, V_j)| \leq \eta$, where $d(U, V)$ measures the red density between vertex sets U and V . Because the colours are complementary, the same conclusion holds for the blue subgraph. For convenience of notation, we will assume below that all V_i have precisely the same order N/m .

We now form a coloured reduced graph with vertex set v_1, \dots, v_m . We leave an edge uncoloured if (W_i, V_j) , (V_i, V_j) and (W_i, W_j) are not all η -regular with $|d(W_i, V_j) - d(V_i, V_j)| \leq \eta$ and $|d(W_i, W_j) - d(V_i, V_j)| \leq \eta$. Otherwise, we fix a constant δ (which will be taken sufficiently small in terms of k and ϵ) and colour the edge $v_i v_j$ red if the red density between V_i and V_j is at least $1 - \delta$ and blue if the blue density is at least δ , breaking ties arbitrarily. Note that there are at most ηm^2 ordered pairs (i, j) whose corresponding edge is uncoloured. Therefore, by deleting at most $\sqrt{\eta}m$ vertices, we may assume that each vertex is adjacent to at most $\sqrt{\eta}m$ uncoloured edges. In what follows, when referring to the reduced graph, we will assume that these vertices have been removed. To each remaining v_i , we assign a colour c_i , either red or blue, depending on which colour has the higher density inside W_i , again breaking ties arbitrarily. By the pigeonhole principle, at least $s := \lceil (1 - \sqrt{\eta})m/2 \rceil$ of the c_i are the same. Without loss of generality, we will assume that this colour is red and c_1, \dots, c_s are red.

Suppose now that there is a red vertex v_a in the reduced graph which has degree at least $\ell := 2^{-k}m$ in red, with neighbours $v_{b_1}, \dots, v_{b_\ell}$. Since the density of red edges in W_a is at least $1/2$, we may apply Lemma 5 with $U_1 = \dots = U_k = W_a$ and $U_{k+s} = V_{b_s}$ for $s = 1, \dots, \ell$ to conclude that, for η sufficiently small in terms of δ , there is a red K_k which is contained in at least

$$\sum_{s=1}^{\ell} (d(W_a, V_{b_s})^k - \delta) |V_{b_s}| \geq ((1 - \delta - \eta)^k - \delta) \ell \frac{N}{m} = ((1 - \delta - \eta)^k - \delta) 2^{-k} N$$

red K_{k+1} . Provided η and δ are sufficiently small in terms of k and ϵ , this is at least n , so we obtain the required book $B_n^{(k)}$. We may therefore assume that we are in the other case, where every red vertex in the reduced graph has blue degree at least $m - \ell - 2\sqrt{\eta}m \geq (1 - 2^{-k} - 2\sqrt{\eta})m$.

The degree of each red vertex is therefore at least $(1 - 2^{-k} - 2\sqrt{\eta})m$ in blue. If we restrict to a set S consisting of s of the red vertices, the blue degree of each vertex inside this set is at least $s - (2^{-k} + 2\sqrt{\eta})m \geq (1 - 2^{-(k-1)} - 16\sqrt{\eta})s$. Since $1 - 2^{-(k-1)} - 16\sqrt{\eta} > 1 - (k-1)^{-1} + \beta$ for some β depending only on k and the number of uncoloured edges is at most $\eta m^2 \leq 8\eta s^2$, Lemma 7 implies that for m sufficiently large and η sufficiently small in terms of k and t , where t is a constant to be fixed below, the reduced graph contains a blue blow-up of K_k with at least t vertices in each part, where each part is itself a monochromatic clique.

We now claim that none of these monochromatic cliques can be blue. Indeed, suppose otherwise and C is a blue clique of order t . If any of the vertices in C , say v_a , is such that $\sum_s d(W_a, V_s) \geq \frac{1}{2}m$, where the sum is taken over all s such that (W_a, V_s) is η -regular, then we have

$$\sum_s d(W_a, V_s)^k \geq m \left(\frac{\sum_s d(W_a, V_s)}{m} \right)^k \geq 2^{-k}m.$$

Again, since the density of red edges in W_a is at least $1/2$, we may apply Lemma 5 with $U_1 = \dots = U_k = W_a$ and U_{k+s} equal in turn to each of the V_s for which (W_a, V_s) is η -regular to conclude that, for η sufficiently small in terms of δ , there is a red K_k which is contained in at least

$$\sum_s (d(W_a, V_s)^k - \delta)|V_s| \geq (2^{-k} - \delta)N$$

red K_{k+1} . Provided η and δ are sufficiently small in terms of k and ϵ , this is at least n , so we again obtain the required book $B_n^{(k)}$.

Therefore, writing $\bar{d}(U, V)$ for the blue density between sets U and V , we must have $\sum_s \bar{d}(W_a, V_s) \geq (\frac{1}{2} - 2\sqrt{\eta})m$ for all $v_a \in C$, where the sum is now over all s such that v_s is in the reduced graph. Writing $\bar{d}_C(V_s) = \sum_{v_a \in C} \bar{d}(W_a, V_s)$, we see, by applying Lemma 9 and summing over all s such that v_s is in the reduced graph, that

$$\sum_s \sum_{(a_1, \dots, a_k) \in \binom{C}{k}} \prod_{i=1}^k \bar{d}(W_{a_i}, V_s) \geq \sum_s \binom{\bar{d}_C(V_s)}{k} \geq m \binom{\sum_s \bar{d}_C(V_s)/m}{k}.$$

Therefore, since $\sum_b \bar{d}_C(V_b) \geq \frac{1}{2}(1 - 4\sqrt{\eta})m|C|$, we have, for $t \geq (1 + \xi)k/(\xi - 4\sqrt{\eta})$, that

$$\sum_s \sum_{(a_1, \dots, a_k) \in \binom{C}{k}} \prod_{i=1}^k \bar{d}(W_{a_i}, V_s) \geq 2^{-k}(1 - k\xi)m \binom{|C|}{k}.$$

Hence, there exists a choice of a_1, \dots, a_k such that

$$\sum_s \prod_{i=1}^k \bar{d}(W_{a_i}, V_s) \geq 2^{-k}(1 - k\xi)m.$$

Since, in the reduced graph, each v_{a_i} has at most $\sqrt{\eta}m$ neighbours v_s such that (W_{a_i}, V_s) is not η -regular, if we now sum only over those s such that (W_{a_i}, V_s) is η -regular for all i , we have that

$$\sum_s \prod_{i=1}^k \bar{d}(W_{a_i}, V_s) \geq 2^{-k}(1 - k\xi)m - k\sqrt{\eta}m.$$

We now apply Lemma 5 with $U_i = W_{a_i}$ for each $1 \leq i \leq k$ and U_{k+s} equal in turn to each V_s with (W_{a_i}, V_s) η -regular for all $1 \leq i \leq k$ to conclude that, for η sufficiently small in terms of δ , there is a blue K_k which is contained in at least

$$\sum_s \left(\prod_{i=1}^k \bar{d}(W_{a_i}, V_s) - \delta \right) |V_s| \geq (2^{-k}(1 - k\xi) - k\sqrt{\eta} - \delta)N$$

blue K_{k+1} . Provided η , δ and ξ are sufficiently small (and t is sufficiently large) in terms of k and ϵ , this is again at least n .

This completes the proof of the claim. We may therefore assume that all of the cliques are red and focus on the subgraph of the reduced graph consisting of the k red cliques C_1, \dots, C_k , each of order t , where every edge between C_i and C_j with $i \neq j$ is blue.

Now, for each vertex v in the reduced graph, let $e_i(v)$ be the weighted blue degree of v in each C_i . That is, $e_i(v) = \sum_{w \in C_i} \bar{d}(v, w)$. By Lemma 8, either $\sum_v \sum_i (t - e_i(v))^k \geq (t/2)^k k m'$ or $\sum_v \prod_i e_i(v) \geq (t/2)^k m'$, where $m' = (1 - \sqrt{\eta})m$. In the second case, we see that there must exist a choice of vertices v_{c_1}, \dots, v_{c_k} with $v_{c_i} \in C_i$ such that

$$\sum_s \prod_{i=1}^k \bar{d}(W_{c_i}, V_s) \geq \frac{\sum_s \sum_{c_1, \dots, c_k} \prod_i \bar{d}(W_{c_i}, V_s)}{t^k} = \frac{\sum_s \prod_i (\sum_{c_i \in C_i} \bar{d}(W_{c_i}, V_s))}{t^k} = \frac{\sum_v \prod_i e_i(v)}{t^k} \geq 2^{-k} m'.$$

Since there are at most $k\sqrt{\eta}m$ vertices v_s such that (W_{c_i}, V_s) is not η -regular for all $1 \leq i \leq k$, we may apply Lemma 5 with $U_i = W_{c_i}$ for $i = 1, \dots, k$ and U_{k+s} equal in turn to each V_s such that (W_{c_i}, V_s) is η -regular for each $i \leq i \leq k$ to conclude that, for η sufficiently small in terms of δ , there is a blue K_k which is contained in at least

$$\sum_s \left(\prod_{i=1}^k \bar{d}(W_{c_i}, V_s) - \delta \right) |V_s| \geq (2^{-k}(1 - \sqrt{\eta}) - k\sqrt{\eta} - \delta)N$$

blue K_{k+1} , again giving the required book for η and δ sufficiently small in terms of k and ϵ .

In the first case, there exists a C_j such that $\sum_v (t - e_j(v))^k \geq (t/2)^k m'$. There must therefore exist (not necessarily distinct) vertices $d_1, \dots, d_k \in C_j$ such that

$$\sum_s \prod_{i=1}^k d(W_{d_i}, V_s) \geq \frac{\sum_s \sum_{d_1, \dots, d_k} \prod_i d(W_{d_i}, V_s)}{t^k} = \frac{\sum_s (\sum_{d \in C_j} d(W_d, V_s))^k}{t^k} = \frac{\sum_v (t - e_j(v))^k}{t^k} \geq 2^{-k} m'.$$

If we again remove the at most $k\sqrt{\eta}m$ vertices v_s such that (W_{d_i}, V_s) is not η -regular for all $1 \leq i \leq k$, we may apply Lemma 5 with $U_i = W_{d_i}$ for $i = 1, \dots, k$ and U_{k+s} equal in turn to each V_s such that (W_{d_i}, V_s) is η -regular for each $1 \leq i \leq k$ to conclude that, for η sufficiently small in terms of δ , there is a red K_k which is contained in at least

$$\sum_s \left(\prod_{i=1}^k d(W_{d_i}, V_s) - \delta \right) |V_s| \geq (2^{-k}(1 - \sqrt{\eta}) - k\sqrt{\eta} - \delta)N$$

red K_{k+1} , giving the required book in this final case provided η and δ are again small enough in terms of k and ϵ . This completes the proof.

4 Concluding remarks

One obvious question is whether a multicolour analogue of Theorem 1 might hold. This is certainly not the case when the number of colours is large. To see this, we use the fact that there exist q -colourings of the complete graph on vertex set $\{1, 2, \dots, 2^{qk/4}\}$ with no monochromatic K_k . Fix such a colouring χ . We consider the $(q+1)$ -coloured complete graph whose vertex set is split into $t = 2^{qk/4}$ vertex sets

V_1, \dots, V_t , each of order n , where every edge between V_i and V_j receives the colour $\chi(i, j)$ and edges internal to any V_i all receive a $(q + 1)$ st colour. This colouring contains no monochromatic $B_n^{(k)}$, so the $(q + 1)$ -colour Ramsey number $r(B_n^{(k)}; q + 1) \geq 2^{qk/4}n$, far greater than the $(q + 1)^k n$ bound one might hope for. More generally, we have $r(B_n^{(k)}; q + 1) \geq (r(k; q) - 1)n$, so, if true, the problem of showing that $r(B_n^{(k)}; 3) \leq 3^k n + o_k(n)$ is at least as hard as showing that $r(k) \leq 3^{k+o(k)}$.

It is also tempting to generalise Theorem 1 to hypergraphs. To this end, we define $B_n^{(k,s)}$ to be the s -uniform hypergraph consisting of n copies of $K_{k+1}^{(s)}$, all sharing a common $K_k^{(s)}$. The natural conjecture would then be that

$$r(B_n^{(k,s)}) = 2^{-\binom{k}{s-1}}n + o_{k,s}(n).$$

However, this is false for $s \geq 4$. To see this, suppose that $s \geq 3$, k is a multiple of s and there is a 2-colouring χ of the s -uniform hypergraph on vertex set $\{1, 2, \dots, r - 1\}$ with no monochromatic $K_{k/s}^{(s)}$. Consider the complete s -uniform hypergraph whose vertex set is split into $r - 1$ vertex sets V_1, \dots, V_{r-1} , each of order n . To colour this hypergraph, suppose that $\{v_1, \dots, v_s\}$ is an edge and $v_j \in V_{i_j}$ for all $1 \leq j \leq s$. If the i_j are all distinct, we colour the edge by $\chi(i_1, \dots, i_s)$ and if the i_j are all the same, we colour the edge red. Otherwise, we colour the edge blue. Since the colouring χ contains no monochromatic $K_{k/s}^{(s)}$, at least s elements of the spine are contained in the same set V_i . But this implies that the book must be red and, therefore, entirely contained within V_i , which is not large enough to contain it, a contradiction. Since we may take r to be $r(K_{k/s}^{(s)})$, this implies that

$$r(B_n^{(k,s)}) \geq (r(K_{k/s}^{(s)}) - 1)n.$$

The value of $r(K_k^{(s)})$ is known to be at least an $(s - 2)$ -fold exponential in k (see, for example, [6]), so this disproves the conjecture for $s \geq 4$. The $s = 3$ case remains unresolved, though a negative answer would again follow from improved lower bounds for $r(K_k^{(3)})$.

As a final remark, we note that there is a strong analogy between Theorem 1 and Green's popular progression theorem [17] (see also [18]). This says that for every $\epsilon > 0$ there exists n_0 such that if $n \geq n_0$ and A is a subset of $\{1, 2, \dots, n\}$ of size αn , then there is $d \neq 0$ such that A contains at least $(\alpha^3 - \epsilon)n$ arithmetic progressions of length 3 with common difference d . That is, there are asymptotically at least as many arithmetic progressions of length 3 in A with common difference d as there would be in a random subset of $\{1, 2, \dots, n\}$ of the same size. A surprising recent result of Fox, Pham and Zhao [14, 15] says that n_0 grows as a tower-type function of ϵ , showing that an application of the (arithmetic) regularity lemma in Green's proof is in some sense necessary. It would be very interesting if a similar phenomenon held for our result, though this seems unlikely to this author.

Acknowledgements. This paper was partially written while I was visiting the California Institute of Technology as a Moore Distinguished Scholar and I am extremely grateful for their kind support.

References

- [1] N. Alon, E. Fischer, M. Krivelevich and M. Szegedy, Efficient testing of large graphs, *Combinatorica* **20** (2000), 451–476.
- [2] S. A. Burr and V. Rosta, On the Ramsey multiplicity of graphs – problems and recent results, *J. Graph Theory* **4** (1980), 347–361.

- [3] D. Conlon, A new upper bound for diagonal Ramsey numbers, *Ann. of Math.* **170** (2009), 941–960.
- [4] D. Conlon, On the Ramsey multiplicity of complete graphs, *Combinatorica* **32** (2012), 171–186.
- [5] D. Conlon and J. Fox, Bounds for graph regularity and removal lemmas, *Geom. Funct. Anal.* **22** (2012), 1191–1256.
- [6] D. Conlon, J. Fox and B. Sudakov, Recent developments in graph Ramsey theory, in *Surveys in Combinatorics 2015*, 49–118, London Math. Soc. Lecture Note Ser., 424, Cambridge Univ. Press, Cambridge, 2015.
- [7] J. Cummings, D. Král', F. Pfender, K. Sperfeld, A. Treglown and M. Young, Monochromatic triangles in three-coloured graphs, *J. Combin. Theory Ser. B* **103** (2013), 489–503.
- [8] P. Erdős, Some remarks on the theory of graphs, *Bull. Amer. Math. Soc.* **53** (1947), 292–294.
- [9] P. Erdős, On the number of complete subgraphs contained in certain graphs, *Magyar Tud. Akad. Mat. Kutató Int. Közl.* **7** (1962), 459–464.
- [10] P. Erdős, R. J. Faudree, C. C. Rousseau and R. H. Schelp, The size Ramsey number, *Period. Math. Hungar.* **9** (1978), 145–161.
- [11] P. Erdős and A. H. Stone, On the structure of linear graphs, *Bull. Amer. Math. Soc.* **52** (1946), 1087–1091.
- [12] P. Erdős and G. Szekeres, A combinatorial problem in geometry, *Compos. Math.* **2** (1935), 463–470.
- [13] J. Fox, There exist graphs with super-exponential Ramsey multiplicity constant, *J. Graph Theory* **57** (2008), 89–98.
- [14] J. Fox and H. T. Pham, Popular progression differences in vector spaces, preprint available at [arXiv:1708.08482](https://arxiv.org/abs/1708.08482) [math.CO].
- [15] J. Fox, H. T. Pham and Y. Zhao, Tower-type bounds for Roth's theorem with popular differences, preprint.
- [16] A. W. Goodman, On sets of acquaintances and strangers at any party, *Amer. Math. Monthly* **66** (1959), 778–783.
- [17] B. Green, A Szemerédi-type regularity lemma in abelian groups, with applications, *Geom. Funct. Anal.* **15** (2005), 340–376.
- [18] B. Green and T. Tao, An arithmetic regularity lemma, an associated counting lemma, and applications, in *An Irregular Mind (Szemerédi is 70)*, 261–334, Springer, 2010.
- [19] H. Hatami, J. Hladký, D. Král', S. Norine and A. Razborov, Non-three-colorable common graphs exist, *Combin. Probab. Comput.* **21** (2012), 734–742.
- [20] C. Jagger, P. Šťovíček and A. Thomason, Multiplicities of subgraphs, *Combinatorica* **16** (1996), 123–141.

- [21] V. Nikiforov, C. C. Rousseau and R. H. Schelp, Book Ramsey numbers and quasi-randomness, *Combin. Probab. Comput.* **14** (2005), 851–860.
- [22] V. Rödl and M. Schacht, Regularity lemmas for graphs, in *Fete of combinatorics and computer science*, 287–325, Bolyai Soc. Math. Stud., 20, János Bolyai Math. Soc., Budapest, 2010.
- [23] C. C. Rousseau and J. Sheehan, On Ramsey numbers of books, *J. Graph Theory* **2** (1978), 77–87.
- [24] J. H. Spencer, Ramsey’s theorem — a new lower bound, *J. Combin. Theory Ser. A* **18** (1975), 108–115.
- [25] A. Thomason, On finite Ramsey numbers, *Eur. J. Combin.* **3** (1982), 263–273.
- [26] A. Thomason, A disproof of a conjecture of Erdős in Ramsey theory, *J. London Math. Soc.* **39** (1989), 246–255.