

# Proof of the Kalai-Meshulam conjecture

Maria Chudnovsky<sup>1</sup>

Princeton University, Princeton, NJ 08544

Alex Scott<sup>2</sup>

Mathematical Institute, University of Oxford, Oxford OX2 6GG, UK

Paul Seymour<sup>3</sup>

Princeton University, Princeton, NJ 08544

Sophie Spirkl<sup>4</sup>

Princeton University, Princeton, NJ 08544

June 4, 2018; revised April 29, 2019

<sup>1</sup>Supported by NSF Grant DMS-1763817. This material is based upon work supported in part by the U.S. Army Research Laboratory and the U.S. Army Research Office under grant number W911NF-16-1-0404.

<sup>2</sup>Supported by a Leverhulme Trust Research Fellowship.

<sup>3</sup>Supported by ONR grant N00014-14-1-0084 and NSF grant DMS-1265563.

<sup>4</sup>Current address: Rutgers University, New Brunswick, NJ 08901

### **Abstract**

Let  $G$  be a graph, and let  $f_G$  be the sum of  $(-1)^{|A|}$ , over all stable sets  $A$ . If  $G$  is a cycle with length divisible by three, then  $f_G = \pm 2$ . Motivated by topological considerations, G. Kalai and R. Meshulam [8] made the conjecture that, if no induced cycle of a graph  $G$  has length divisible by three, then  $|f_G| \leq 1$ . We prove this conjecture.

# 1 Introduction

In the late 1990's, G. Kalai and R. Meshulam [8] made an intriguing sequence of conjectures about the connections between induced cycle lengths, chromatic number, and the number of stable sets of different parities in a graph.

A graph is *ternary* if no induced cycle has length a multiple of three; thus, ternary graphs have no triangles. (All graphs in this paper are finite and have no loops or parallel edges.) First, Kalai and Meshulam conjectured:

**1.1** *There exists  $c$  such that every ternary graph is  $c$ -colourable.*

This was proved by Bonamy, Charbit and Thomassé [1], for some large constant  $c$  (although it may be that all ternary graphs are 3-colourable, and this remains open). A much stronger result was later proved by two of us [9]: that for all integers  $p, q \geq 0$ , every graph with bounded clique number and with no induced cycle of length  $p$  modulo  $q$  has bounded chromatic number.

Second, Kalai and Meshulam conjectured:

**1.2** *For every ternary graph, the number of stable sets with even cardinality and the number with odd cardinality differ by at most one.*

This has remained open, and we prove it in this paper.

Two further conjectures of Kalai and Meshulam were proved in [9]. The stronger of these conjectures stated that for all  $k$  there exists  $c$ , such that, if for every induced subgraph of  $G$  the number of even stable sets and the number of odd ones differ by at most  $k$ , then  $G$  is  $c$ -colourable. This follows from a generalization of the strengthening of 1.1 mentioned above.

A final Kalai-Meshulam conjecture concerns Betti numbers and ternary graphs. The *independence complex*  $I(G)$  of a graph  $G$  is the simplicial complex whose faces are the stable sets of vertices of  $G$ . Let  $b_i$  denote the  $i$ th Betti number of  $I(G)$  and let  $b(G)$  denote the sum of the Betti numbers.

**1.3 Conjecture:** *A graph  $G$  is ternary if and only if  $|b(H)| \leq 1$  for every induced subgraph  $H$ .*

Let  $f_G(\emptyset)$  denote the number of even stable sets in  $G$  minus the number of odd ones. If  $|b(H)| \leq 1$  for every induced subgraph  $H$ , then  $G$  has no induced cycle of length divisible by 3, since  $b(H) = 2$  for every cycle  $H$  of length divisible by three. For the converse, suppose  $G$  has no such induced cycle. Then by 1.2,  $|f_G(\emptyset)| \leq 1$ , but we need to prove that  $b(G) \leq 1$ . Now  $f_G(\emptyset)$  is the Euler characteristic of  $I(G)$ , and in particular there is a connection between  $f_G(\emptyset)$  and  $b(G)$ . It is a basic theorem from homology theory that the Euler characteristic of  $I(G)$  is the alternating sum of the Betti numbers of  $I(G)$  (see [6]). It follows that  $|f_G(\emptyset)| \leq b(G)$ ; but this inequality is in the wrong direction for us, and the conjecture remains open.

We mention a few other related results:

- Chen and Saito [3] proved that every non-null graph with no cycle of length divisible by three (not just induced cycles) has a vertex of degree at most two (and so all such graphs are 3-colourable).
- G. Gauthier [5] found an explicit construction for all graphs with no cycle of length divisible by three.

- D. Král’ asked (unpublished): is it true that in every ternary graph with an edge, there is an edge  $e$  such that the graph obtained by deleting  $e$  is also ternary? This would have implied that all ternary graphs are 3-colourable, but has very recently been disproved; a counterexample was found by M. Wrochna. (Take the disjoint union of a 5-cycle and a 10-cycle, and join each vertex of the 5-cycle to two opposite vertices of the 10-cycle, in order.)
- The difference between the numbers of odd and even stable sets has also appeared in statistical physics. Let us define the polynomial

$$I_G(z) = \sum_I z^{|I|},$$

where the sum is over stable sets  $I$  in  $G$ . This polynomial is known in combinatorics as the *independent set polynomial* and statistical physics as the *partition function of the hard-core lattice gas* (see, for instance, [10]). We see that  $I_G(-1)$  is the number of even stable sets minus the number of odd stable sets. The question of when  $|I_G(-1)| \leq 1$  has been the focus of considerable study, particularly on the square lattice (see [2, 4, 7]).

If  $G$  is a graph, and  $X, Y$  are disjoint subsets of  $V(G)$ , let  $f_G(X, Y)$  be the sum of  $(-1)^{|A|}$ , summed over all stable sets  $A$  in  $G$  that include  $X$  and are disjoint from  $Y$ . Our main theorem states:

**1.4** *If  $G$  is ternary then  $|f_G(\emptyset, \emptyset)| \leq 1$ .*

The proof of 1.4 is by induction on  $|V(G)|$ , and it follows easily that if  $G$  is a minimum counterexample then  $f_G(\emptyset, \emptyset) = \pm 2$ . It is very helpful to know the value of  $f_G(\emptyset, \emptyset)$ , and so the proof breaks into two cases, depending whether this value is 2 or  $-2$ . The proof for the second is obtained from the first proof by negating  $f_G$  throughout, and we would like to say “we may assume that  $f_G(\emptyset, \emptyset) = 2$  without loss of generality”; but this gives us a difficulty, because negating  $f_G$  does not give a function that equals  $f_H$  for some graph  $H$ . We overcome this as follows.

Let  $G$  be a graph, and with  $f_G$  as before, let us say the functions  $f_G$  and  $-f_G$  are *counters* on  $G$ . We will prove that if  $G$  is ternary and  $g$  is a counter on  $G$ , then  $|g(\emptyset, \emptyset)| \leq 1$ . Now we are free to replace  $g$  by its negative if that is convenient.

We will frequently need to talk about  $g(X, Y)$  when  $Y = \emptyset$ ; so often that it is worthwhile to make a special convention for it. We define  $g(X) = g(X, \emptyset)$  (and the same for  $f_G$ ).

If  $g$  is a counter on  $G$ , we say  $g$  is a *good counter* if for all disjoint  $X, Y \subseteq V(G)$  with  $X \cup Y \neq \emptyset$ :

- $|g(X, Y)| \leq 1$ ; and
- $|g(X \cup \{u\}, Y) - g(X \cup \{v\}, Y)| \leq 1$  for all  $u, v \in V(G) \setminus (X \cup Y)$ .

In section 3, we show that:

**1.5** *If  $g$  is a good counter on a graph  $G$ , then  $|g(\{u\}) - g(\{v\})| \leq 1$  for all  $u, v \in V(G)$ .*

Then in section 4, we show that:

**1.6** *If  $g$  is a good counter on a ternary graph  $G$ , then  $|g(\emptyset)| \leq 1$ .*

**Proof of 1.4, assuming 1.5 and 1.6.** We prove by induction on  $|V(G)|$  that for every ternary graph  $G$ , if  $g$  is a counter on  $G$ , then  $|g(\{u\}) - g(\{v\})| \leq 1$  for all  $u, v \in V(G)$ , and  $|g(\emptyset)| \leq 1$ . Thus we may assume that these two statements hold for every proper induced subgraph of  $G$ . Now  $g$  is a counter on  $G$ , and so  $g = \pm f_G$ . If the result holds for  $-g$  then it holds for  $g$ ; so we may assume that  $g = f_G$ , by replacing  $g$  by  $-g$  if necessary.

(1) *If  $X, Y \subseteq V(G)$  are disjoint, with  $X \cup Y \neq \emptyset$ , then  $|f_G(X, Y)| \leq 1$ .*

We may assume that  $X$  is a stable set. Let  $H$  be the graph obtained from  $G$  by deleting  $X \cup Y$  and deleting all vertices with a neighbour in  $X$ . Thus, if  $A$  is a stable set of  $G$  including  $X$  and disjoint from  $Y$ , then  $A \setminus X$  is a stable set of  $H$ ; and conversely, if  $B$  is a stable set of  $H$ , then  $X \cup B$  is a stable set of  $G$  including  $X$  and disjoint from  $Y$ . In particular,  $f_H(\emptyset) = (-1)^{|X|} f_G(X, Y)$ ; but from the inductive hypothesis,  $|f_H(\emptyset)| \leq 1$ , and so  $|f_G(X, Y)| \leq 1$ . This proves (1).

(2) *If  $X, Y \subseteq V(G)$  are disjoint, with  $X \cup Y \neq \emptyset$ , and  $u, v \in V(G) \setminus (X \cup Y)$ , then*

$$|f_G(X \cup \{u\}, Y) - f_G(X \cup \{v\}, Y)| \leq 1.$$

We may assume that  $X$  is stable. Suppose first that  $u$  has a neighbour in  $X$ . Then  $f_G(X \cup \{u\}, Y) = 0$  (because  $X \cup \{u\}$  is not a subset of any stable set). Also  $|f_G(X \cup \{v\}, Y)| \leq 1$ , by (1), and the claim follows. So we may assume that  $u$  and similarly  $v$  has no neighbour in  $X$ ; and so  $u, v \in V(H)$ , if we define  $H$  as before. Thus  $f_G(X \cup \{u\}, Y) = (-1)^{|X|} f_H(\{u\})$ , and  $f_G(X \cup \{v\}, Y) = (-1)^{|X|} f_H(\{v\})$ ; and from the inductive hypothesis,  $|f_H(\{u\}) - f_H(\{v\})| \leq 1$ . It follows that  $|f_G(X \cup \{u\}, Y) - f_G(X \cup \{v\}, Y)| \leq 1$ . This proves (2).

From (1) and (2),  $g$  is a good counter on  $G$ . From 1.6 and 1.5, it follows that  $|g(\{u\}) - g(\{v\})| \leq 1$  for all  $u, v \in V(G)$ , and  $|g(\emptyset)| \leq 1$ . This completes the inductive proof; and 1.4 follows.  $\blacksquare$

## 2 Some lemmas

Here are a few useful lemmas. First, we observe:

**2.1** *Let  $g$  be a counter on  $G$ , let  $X, Y \subseteq V(G)$  be disjoint, and let  $Y' \subseteq Y$ . Then*

$$g(X, Y) = \sum_{Z \subseteq Y \setminus Y'} (-1)^{|Z|} g(X \cup Z, Y').$$

**Proof.** We may assume that  $g = f_G$ , by replacing  $g$  by  $-g$  if necessary. If  $A$  is a stable set of  $G$  including  $X$  and disjoint from  $Y'$ , define  $n_A$  to be

$$\sum_{Z \subseteq A \cap Y} (-1)^{|A| - |Z|}.$$

Thus  $n_A = 0$  unless  $A \cap Y = \emptyset$ , in which case  $n_A = (-1)^{|A|}$ . But  $\sum_{Z \subseteq Y \setminus Y'} (-1)^{|Z|} f_G(X \cup Z, Y')$  is the sum of  $n_A$ , over all stable sets  $A$  of  $G$  including  $X$  and disjoint from  $Y'$ . It follows that  $\sum_{Z \subseteq Y \setminus Y'} (-1)^{|Z|} f_G(X \cup Z, Y')$  is the sum of  $(-1)^{|A|}$  over all stable sets of  $G$  that include  $X$  and are disjoint from  $Y$ . But this sum equals  $f_G(X, Y)$ . This proves 2.1.  $\blacksquare$

In evaluating an expression given by 2.1, it often happens that for some number  $\ell$ ,  $g(X \cup Z) = \ell$  for “most” subsets  $Z \subseteq Y$ , and if so the following is helpful:

**2.2** *Let  $g$  be a counter on  $G$ , let  $X, Y \subseteq V(G)$  be disjoint, with  $Y \neq \emptyset$ , and let  $\ell$  be some number. Then*

$$g(X, Y) = \sum_{Z \subseteq Y} (-1)^{|Z|} (g(X \cup Z) - \ell).$$

**Proof.** By 2.1,

$$g(X, Y) = \sum_{Z \subseteq Y} (-1)^{|Z|} g(X \cup Z),$$

and  $\sum_{Z \subseteq Y} (-1)^{|Z|} (-\ell) = 0$  since  $Y \neq \emptyset$ . This proves 2.2. ■

**2.3** *Let  $g$  be a good counter on  $G$ , let  $X, Y \subseteq V(G)$  be disjoint, and let  $v \in V(G) \setminus (X \cup Y)$ . Then  $|g(X, Y) - g(X \cup \{v\}, Y)| \leq 1$  and  $|g(X, Y) - g(X, Y \cup \{v\})| \leq 1$ .*

**Proof.** We may assume that  $g = f_G$ . Every stable set including  $X$  and disjoint from  $Y$  either includes  $X \cup \{v\}$  or is disjoint from  $Y \cup \{v\}$ , and not both. Consequently

$$g(X, Y) = g(X \cup \{v\}, Y) + g(X, Y \cup \{v\}).$$

But  $|g(Y \cup \{v\})| \leq 1$  since  $g$  is a good counter, and therefore  $|g(X, Y) - g(X \cup \{v\}, Y)| \leq 1$ ; and the second claim follows similarly. ■

For  $X \subseteq V(G)$ , let  $N[X]$  denote the set of vertices in  $G$  that either belong to  $X$  or have a neighbour in  $X$ . We observe that

**2.4** *Let  $g$  be a counter on  $G$ . If  $X, Y \subseteq V(G)$  are disjoint with  $g(X, Y) \neq 0$ , and  $v \in V(G) \setminus (N[X] \cup Y)$ , then  $v$  has a neighbour in  $V(G) \setminus (N[X] \cup Y)$ .*

**Proof.** We may assume that  $g = f_G$ , by replacing  $g$  by  $-g$  if necessary. The stable sets of  $G$  that include  $X$  and are disjoint from  $Y$  are obtained from the stable sets of  $G \setminus (N[X] \cup Y)$  ( $= H$  say) by adding the set  $X$  to each such stable set; and so  $f_H(\emptyset) \neq 0$ . But  $f_K(\emptyset) = 0$  for every graph  $K$  with a vertex of degree zero, and so  $H$  has no vertex of degree zero. The result follows. ■

**2.5** *Let  $g$  be a good counter on  $G$ , let  $X, Y \subseteq V(G)$  be disjoint, and let  $u, v \in V(G) \setminus (X \cup Y)$ . If  $g(X, Y) = g(X \cup \{u, v\}, Y) \neq 0$ , then  $g(X, Y) = g(X \cup \{v\}, Y)$ .*

**Proof.** We proceed by induction on  $|V(G) \setminus (X \cup Y)|$ . By replacing  $g$  by  $-g$  if necessary we may assume that  $g(X, Y) > 0$ . For all disjoint  $A, B \subseteq V(G) \setminus (X \cup Y)$ , let  $h(A, B) = g(X \cup A, Y \cup B)$  (and  $h(A)$  means  $h(A, \emptyset)$ ). Since  $g$  is a good counter it follows that  $|h(\{u, v\})| \leq 1$ , and so  $h(\{u, v\}) = h(\emptyset) = 1$ . We suppose for a contradiction that  $h(\{v\}) \neq 1$ . Hence  $u \neq v$ , and  $X \cup \{u, v\}$  is stable. By 2.3, it follows that  $h(\{v\}) = 0$ . Since  $|h(\emptyset, \{u, v\})| \leq 1$ , 2.1 implies that

$$h(\emptyset) - h(\{u\}) - h(\{v\}) + h(\{u, v\}) \leq 1.$$

Consequently  $h(\{u\}) \geq 1$ , and so  $h(\{u\}) = 1$ . From 2.4,  $v$  has a neighbour  $w$ .

Now  $h(\emptyset, \{v\}) = h(\emptyset) - h(\{v\}) = 1$ , and  $h(\{u\}, \{v\}) = h(\{u\}) - h(\{u, v\}) = 0$ , and so from the inductive hypothesis,  $h(\{u, w\}, \{v\}) \neq 1$ . Consequently  $h(\{u, w\}) - h(\{u, v, w\}) \neq 1$ , and since  $h(\{u, v, w\}) = 0$ , it follows that  $h(\{u, w\}) \neq 1$ . By 2.3,  $h(\{u, w\}) = 0$ . Thus  $h(\{u\}, \{w\}) = 1$  by 2.1, since  $h(\{u\}) = 1$ . Since  $h(\{v\}, \{w\}) = 0$  and  $h(\{u, v\}, \{w\}) = 1$  by 2.1 (the first since  $h(\{v, w\}) = 0$  and  $h(\{v\}) = 0$ , and the second since  $h(\{u, v, w\}) = 0$  and  $h(\{u, v\}) = 1$ ), it follows from the inductive hypothesis that  $h(\emptyset, \{w\}) \neq 1$ , and so  $h(\emptyset, \{w\}) = 0$  by 2.3. Hence  $h(\emptyset) - h(\{w\}) = 0$  by 2.1, and so  $h(\{w\}) = 1$ . But then  $h(\{w\}, \{u\}) = 1$ , because  $h(\{u, w\}) = 0$ ; and  $h(\{v\}, \{u\}) = -1$ , since  $h(\{v\}) = 0$  and  $h(\{u, v\}) = 1$ . This contradicts that  $g$  is good, and so proves 2.5.  $\blacksquare$

The next result has been independently discovered several times.

**2.6** *Let  $G$  be a nonnull graph and let  $A_1, A_2, A_3$  be the classes of a 3-colouring of  $G$ . Suppose that for  $i = 1, 2, 3$ , every vertex in  $A_i$  has a neighbour in  $A_{i+1}$ , where  $A_4$  means  $A_1$ . Then  $G$  is not ternary.*

**Proof.** Throughout we read subscripts modulo 3. For  $i = 1, 2, 3$ , direct each edge of  $G$  between  $A_i$  and  $A_{i+1}$  from  $A_i$  to  $A_{i+1}$ . Since each vertex has positive outdegree, the digraph we form has a directed cycle, and hence an induced directed cycle. But such a cycle is an induced cycle of  $G$ , and has length a multiple of three.  $\blacksquare$

**2.7** *Let  $\mathcal{H}$  be a set of subsets of some set  $V$ , all of the same cardinality  $k$ ; and suppose that for every subset  $X \subseteq V$  with  $|X| = k + 1$ , if  $X$  includes a member of  $\mathcal{H}$  then it includes at least two such members. Then there is a partition  $P_1, \dots, P_n$  of  $V$  with  $P_1, \dots, P_n$  all nonempty, such that for all distinct  $u, v \in V$ , either there exists  $i \in \{1, \dots, n\}$  with  $u, v \in P_i$ , or there exists  $B \in \mathcal{H}$  with  $u, v \in B$ , and not both.*

**Proof.** Say two vertices  $u, v \in V$  are *equivalent* if either  $u = v$ , or:

- there is no member of  $\mathcal{H}$  containing both  $u, v$ ; and
- for each  $C \subseteq V \setminus \{u, v\}$ ,  $C \cup \{u\} \in \mathcal{H}$  if and only if  $C \cup \{v\} \in \mathcal{H}$ .

We claim that this is an equivalence relation. To see this, we may assume that  $u, v, w \in V(G)$  are distinct, and  $v$  is equivalent to both  $u$  and  $w$ ; and we must show that  $u, w$  are equivalent. If there exists  $B \in \mathcal{H}$  containing  $u, w$ , then  $v \notin B$  (since  $u, v$  are equivalent) and so  $(B \setminus \{u\}) \cup \{v\} \in \mathcal{H}$  (since  $(B \setminus \{u\}) \cup \{u\} \in \mathcal{H}$  and  $u, v$  are equivalent), and so this is a member of  $\mathcal{H}$  containing  $v, w$ , a contradiction. Thus there is no such  $B$ . Let  $C \subseteq V \setminus \{u, w\}$ , with  $C \cup \{u\} \in \mathcal{H}$ . Consequently  $v \notin C$ , and  $C \cup \{v\} \in \mathcal{H}$  (because  $u, v$  are equivalent), and consequently  $C \cup \{w\} \in \mathcal{H}$  (since  $v, w$  are equivalent). Similarly  $C \cup \{u\} \in \mathcal{H}$  if and only if  $C \cup \{w\} \in \mathcal{H}$ . This proves that equivalence is indeed an equivalence relation.

We claim that for all distinct  $u, v \in V$ , if they do not belong to the same equivalence class then some member of  $\mathcal{H}$  contains both  $u, v$ . To see this, since  $u, v$  are not equivalent, if no member of  $\mathcal{H}$  contains both  $u$  and  $v$ , then we may assume (exchanging  $u, v$  if necessary) that there exists  $C \subseteq V \setminus \{u, v\}$  such that  $C \cup \{u\} \in \mathcal{H}$  and  $C \cup \{v\} \notin \mathcal{H}$ . Thus  $|C| = k - 1$ , and since  $C \cup \{u, v\}$  includes a member of  $\mathcal{H}$ , by hypothesis it includes at least two members. But since no member of  $\mathcal{H}$  contains both  $u, v$ , and  $C \cup \{v\} \notin \mathcal{H}$ , this is impossible. This proves 2.7.  $\blacksquare$

### 3 The value on distinct vertices

In this section we prove 1.5. Thus, throughout this section, let  $g$  be a good counter on a graph  $G$ . For  $i = -1, 0, 1$  let  $A_i$  be the set of vertices  $v$  of  $G$  such that  $g(\{v\}) = i$ . Thus  $A_{-1}, A_0, A_1$  are disjoint and have union  $V(G)$ . We need to show that one of  $A_{-1}, A_1$  is empty, and so we assume for a contradiction that they are both nonempty. We will prove a series of statements about  $G, g$ . We begin with:

**3.1** *The following hold:*

- $g(\emptyset) = 0$ ;
- $G$  is connected;
- $A_1, A_{-1}$  are both stable sets;
- there is not both an edge between  $A_1, A_0$  and an edge between  $A_{-1}, A_0$ .

**Proof.** Since there exists  $v \in A_1$ , and hence with  $g(\{v\}) = 1$ , we deduce from 2.3 that  $g(\emptyset) \geq 0$ , and similarly  $g(\emptyset) \leq 0$ . This proves the first statement.

For the second statement, we may assume (replacing  $g$  by  $-g$  if necessary) that  $g = f_G$ . By assumption, there exist  $u_i \in V(G)$  with  $g(\{u_i\}) = i$ , for  $i \in \{1, -1\}$ . Suppose that  $G$  is not connected, and let  $G_1$  be a component of  $G$  containing  $u_1$ , and let  $G_2$  be obtained from  $G$  by deleting  $G_1$ . Write  $g_i$  for  $f_{G_i}$  ( $i = 1, 2$ ). Thus for disjoint  $X, Y \subseteq V(G)$ ,

$$g(X, Y) = g_1(X \cap V(G_1), Y \cap V(G_1))g_2(X \cap V(G_2), Y \cap V(G_2)),$$

and in particular,  $g_1(X) = g(X, V(G_2))$  for  $X \subseteq V(G_1)$ , and  $g_2(X) = g(X, V(G_1))$  for  $X \subseteq V(G_2)$ . Since  $0 = g(\emptyset) = g_1(\emptyset)g_2(\emptyset)$ , one of  $g_1(\emptyset), g_2(\emptyset)$  is zero.

Since  $g(\{u_1\}) = g_1(\{u_1\})g_2(\emptyset)$ , it follows that  $g_2(\emptyset) \neq 0$ , and so  $g_1(\emptyset) = 0$ . In particular,  $G_1$  is the unique component  $C$  of  $G$  such that  $f_C(\emptyset) = 0$ , and so  $u_{-1} \in V(G_1)$ . Thus  $g(\{u_{-1}\}) = g_1(\{u_{-1}\})g_2(\emptyset)$ , and so one of  $g_1(\{u_1\}), g_1(\{u_{-1}\})$  equals 1 and the other equals  $-1$ , contradicting that  $g$  is good. This proves the second statement.

For the third, suppose that  $u, v \in A_1$  are adjacent. By 2.1,

$$g(\emptyset, \{u, v\}) = g(\emptyset) - g(\{u\}) - g(\{v\}) + g(\{u, v\});$$

but the last term is zero since  $u, v$  are adjacent, and since  $u, v \in A_1$  and  $g(\emptyset) = 0$ , we deduce that  $g(\emptyset, \{u, v\}) = -2$ , contradicting that  $g$  is good.

For the fourth statement, suppose that  $u_1 \in A_1$  is adjacent to  $v_1 \in A_0$ , and  $u_{-1} \in A_{-1}$  is adjacent to  $v_{-1} \in A_0$ . Suppose first that  $g(\{v_1, u_{-1}\}) = 0$ . Then by two applications of 2.1,  $g(\{u_{-1}\}, \{v_1\}) = g(\{u_{-1}\}) - g(\{u_{-1}, v_1\}) = -1$ , and  $g(\{u_1\}, \{v_1\}) = g(\{u_1\}) - g(\{u_1, v_1\}) = 1$  (since  $u_1, v_1$  are adjacent), contradicting that  $g$  is good. This proves that  $g(\{v_1, u_{-1}\}) \neq 0$ , and so  $g(\{v_1, u_{-1}\}) = -1$  by 2.3. Similarly  $g(\{v_{-1}, u_1\}) = 1$  (and in particular,  $v_1 \neq v_{-1}$ ). But by 2.1,

$$g(\{v_1\}, \{u_1, u_{-1}\}) = g(\{v_1\}) - g(\{v_1, u_1\}) - g(\{v_1, u_{-1}\}) + g(\{v_1, u_1, u_{-1}\});$$

and since  $g(\{v_1\}) = 0$  and  $g(\{v_1, u_1\}) = g(\{v_1, u_1, u_{-1}\}) = 0$  (since  $u_1, v_1$  are adjacent) it follows that  $g(\{v_1\}, \{u_1, u_{-1}\}) = 1$ . Similarly  $g(\{v_{-1}\}, \{u_1, u_{-1}\}) = -1$ , contradicting that  $g$  is good. This proves 3.1. ■



In the same notation, because of the third statement of 3.1, we may assume (replacing  $g$  by  $-g$  if necessary) that there are no edges between  $A_{-1}$  and  $A_0$ . Let  $B_1$  be the set of vertices  $v \in A_0$  such that  $g(\{u, v\}) = 1$  for each  $u \in A_1$  and  $g(\{u, v\}) = 0$  for each  $u \in A_{-1}$ ; and let  $B_{-1}$  be the set of vertices  $v \in A_0$  such that  $g(\{u, v\}) = 0$  for each  $u \in A_1$  and  $g(\{u, v\}) = -1$  for each  $u \in A_{-1}$ .

**3.2** *Every vertex in  $A_0$  belongs to one of  $B_1, B_{-1}$ .*

**Proof.** Let  $v \in A_0$ , and for  $i \in \{1, -1\}$  let  $u_i \in A_i$ . Not both  $g(\{v, u_1\}) = 1$  and  $g(\{v, u_{-1}\}) = -1$ , since  $g$  is good. Suppose that neither of these holds. Then  $g(\{v, u_1\}) = 0$  and  $g(\{v, u_{-1}\}) = 0$ , by 2.3. Then by two applications of 2.1,  $g(\{u_1\}, \{v\}) = g(\{u_1\}) - g(\{u_1, v\}) = 1$ , and  $g(\{u_{-1}\}, \{v\}) = g(\{u_{-1}\}) - g(\{u_{-1}, v\}) = -1$ , contradicting that  $g$  is good. It follows that either  $g(\{v, u_1\}) = 1$  and  $g(\{v, u_{-1}\}) = 0$ , or  $g(\{v, u_1\}) = 0$  and  $g(\{v, u_{-1}\}) = -1$ . Since this holds for all  $u_1, u_{-1}$ , it follows that  $v \in B_1 \cup B_{-1}$ . This proves 3.2.  $\blacksquare$

**3.3**  *$A_0$  is empty.*

**Proof.** Suppose that  $A_0 \neq \emptyset$ . Since  $G$  is connected by 3.1, and by assumption there are no edges between  $A_{-1}$  and  $A_0$ , it follows that there is an edge between  $A_0$  and  $A_1$ , say between  $b \in A_0$  and  $a_1 \in A_1$ . Consequently  $g(\{a_1, b\}) = 0$ , and so  $b \notin B_1$  from the definition of  $B_1$ ; and so  $b \in B_{-1}$  by 3.2. Choose  $a_{-1} \in A_{-1}$ . By three applications of 2.1,  $g(\emptyset, \{a_1\}) = g(\emptyset) - g(\{a_1\}) = -1$ ,  $g(\{b\}, \{a_1\}) = g(\{b\}) - g(\{b, a_1\}) = 0$ , and  $g(\{b, a_{-1}\}, \{a_1\}) = g(\{b, a_{-1}\}) - g(\{b, a_{-1}, a_1\}) = -1$ , contrary to 2.5. Thus  $A_0 = \emptyset$ . This proves 3.3.  $\blacksquare$

Now we prove 1.5, which we restate:

**3.4** *If  $g$  is a good counter on a graph  $G$ , then  $|g(\{u\}) - g(\{v\})| \leq 1$  for all  $u, v \in V(G)$ .*

**Proof.** As all through this section, we assume that  $G, g$  is a counterexample. In the previous notation, 3.3 and 3.1 imply that  $G$  is bipartite, and  $(A_1, A_{-1})$  is a bipartition. We recall that  $g(\emptyset) = 0$ .

(1) *Every vertex of  $G$  has degree at least two.*

Since  $G$  is connected by 3.1, all vertices have degree at least one; suppose that  $v \in A_1$  has only one neighbour  $u \in A_{-1}$  say. Since  $G$  is connected and  $|V(G)| \geq 3$ ,  $u$  has another neighbour  $v' \in A_1$ . Now  $g(\{v'\}) = 1$ , and since  $v \in V(G) \setminus N[\{v'\}]$ , 2.4 implies that  $v$  has a neighbour in  $V(G) \setminus N[\{v'\}]$ , a contradiction. This proves (1).

(2) *There is a subset  $X \subseteq A_1$  with  $g(X) = 0$ .*

Choose  $v \in A_1$ , and let  $X = A_1 \setminus \{v\}$ . Since  $v \in V(G) \setminus N[X]$ , and  $v$  has no neighbour in  $V(G) \setminus N[X]$  (by (1)), 2.4 implies that  $g(X) = 0$ . This proves (2).

For  $i \in \{1, -1\}$  let  $k_i > 0$  be minimum such that some subset  $B$  of  $A_i$  with cardinality  $k_i$  satisfies  $g(B) \neq i$ . Thus  $k_i \geq 2$ ; and by 2.3,  $g(B) = 0$  or  $i$  for each subset  $B \subseteq A_i$  with  $|B| = k_i$ .

(3) For  $i \in \{1, -1\}$ ,  $k_i$  is odd.

Choose  $B \subseteq A_i$  with cardinality  $k_i$  such that  $g(B) \neq i$ , and hence  $g(B) = 0$ . Since  $g$  is good,  $|g(\emptyset, B)| \leq 1$ ; and so by 2.2,

$$\left| \sum_{Z \subseteq B} (-1)^{|Z|} (g(Z) - i) \right| \leq 1.$$

But  $g(Z) = i$  for all  $Z \subseteq B$  with  $Z \neq B, \emptyset$ , and  $g(Z) = 0$  if  $Z = B, \emptyset$ ; and consequently  $|-i - i(-1)^{k_i}| \leq 1$ , and so  $k_i$  is odd. This proves (3).

Let  $\mathcal{H}_i$  be the set of all subsets  $B$  of  $A_i$  such that  $|B| = k_i$  and  $g(B) = 0$ .

(4) For every subset  $X$  of  $A_i$  with cardinality  $k_i + 1$ , if  $X$  includes a member of  $\mathcal{H}_i$  then it includes at least two such members.

Let  $X = \{v_0, \dots, v_{k_i}\}$ , and suppose that  $\{v_1, \dots, v_{k_i}\}$  is the only member of  $\mathcal{H}_i$  included in  $X$ . Then  $g(X) \neq i$ , by 2.5, and  $g(X) \neq -i$  by 2.3; so  $g(X) = 0$ . Let  $Y = \{v_2, \dots, v_{k_i}\}$ . By 2.2 and (3):

$$\begin{aligned} g(\emptyset, Y) &= \sum_{Z \subseteq Y} (-1)^{|Z|} (g(Z) - i) = -i, \\ g(\{v_0\}, Y) &= \sum_{Z \subseteq Y} (-1)^{|Z|} (g(Z \cup \{v_0\}) - i) = 0, \\ g(\{v_0, v_1\}, Y) &= \sum_{Z \subseteq Y} (-1)^{|Z|} (g(Z \cup \{v_0, v_1\}) - i) = -(-1)^{|Y|} i = -i, \end{aligned}$$

contrary to 2.5. This proves (4).

(5) There exist  $B_i \in \mathcal{H}_i$  for  $i \in \{1, -1\}$ , such that there are two edges of  $G$  between  $B_1$  and  $B_{-1}$  with no end in common.

By (4) and 2.7, there is a partition  $P_1, \dots, P_m$  of  $A_1$  such that every two vertices in  $A_1$  either belong to the same  $P_i$  or to some member of  $\mathcal{H}_1$ , and not both; and let  $Q_1, \dots, Q_n \subseteq A_{-1}$  be defined analogously. Say  $P_i, Q_j$  are adjacent if there is an edge in  $G$  between a vertex in  $P_i$  and a vertex in  $Q_j$ . Since  $m, n \geq 2$  and each  $P_i$  is adjacent to some  $Q_j$  and vice versa, there are distinct  $P_1, P_2$  (say) and distinct  $Q_1, Q_2$  such that  $P_1$  is adjacent to  $Q_1$  and  $P_2$  to  $Q_2$ . Choose  $p_i \in P_i$  and  $q_i \in Q_i$  ( $i = 1, 2$ ) such that  $p_1 q_1$  and  $p_2 q_2$  are edges of  $G$ . Since  $p_1, p_2$  do not belong to the same one of  $P_1, \dots, P_m$ , there exists  $B_1 \in \mathcal{H}_1$  containing  $p_1, p_2$ ; and similarly there exists  $B_{-1} \in \mathcal{H}_{-1}$  containing  $q_1, q_2$ . This proves (5).

For  $i \in \{1, -1\}$  choose  $B_i$  as in (5).

(6) For  $i \in \{1, -1\}$ , let  $X_i \subseteq B_i$  with  $\emptyset \neq X_i \neq B_i$ . Then  $g(X_1 \cup X_{-1}) = 0$ .

Suppose not, and for  $i \in \{1, -1\}$  choose  $X_i \subseteq B_i$  with  $\emptyset \neq X_i \neq B_i$ , with  $X_1 \cup X_{-1}$  minimal

such that  $g(X_1 \cup X_{-1}) \neq 0$ . We may assume that  $g(X_1 \cup X_{-1}) = 1$ , by replacing  $g$  by  $-g$  if necessary. By 2.1 and the minimality of  $X_1 \cup X_{-1}$ ,

$$g(X_1, X_{-1}) = g(X_1) + (-1)^{|X_{-1}|} g(X_1 \cup X_{-1}) = 1 + (-1)^{|X_{-1}|},$$

and so  $|X_{-1}|$  is odd; and similarly  $|X_1|$  is even. Choose  $u \in X_1$  and  $v \in X_{-1}$ . Then by three applications of 2.1,

$$\begin{aligned} g(X_1 \setminus \{u\}, X_{-1} \setminus \{v\}) &= g(X_1 \setminus \{u\}) = 1, \\ g((X_1 \cup \{v\}) \setminus \{u\}, X_{-1} \setminus \{v\}) &= 0, \\ g(X_1 \cup \{v\}, X_{-1} \setminus \{v\}) &= (-1)^{|X_{-1} \setminus \{v\}|} g(X_1 \cup X_{-1}) = 1, \end{aligned}$$

contrary to 2.5. This proves (6).

Choose  $C_1 \subseteq B_1$  maximal such that either  $C_1 = \emptyset$  or  $g(C_1 \cup B_{-1}) \neq 0$ , and choose  $C_{-1} \subseteq B_{-1}$  maximal such that either  $C_{-1} = \emptyset$  or  $g(C_{-1} \cup B_1) \neq 0$ . It follows that  $|C_i| \leq k_i - 2$  for  $i \in \{1, -1\}$ , since there is a 2-edge matching between  $B_1, B_{-1}$ . For  $i \in \{1, -1\}$  let  $D_i = B_i \setminus C_i$ , and let  $C = C_1 \cup C_{-1}$  and  $D = D_1 \cup D_{-1}$ .

(7) *If  $C_1 \neq \emptyset$  then  $g(C_1 \cup B_{-1}) = 1$ ; and if  $C_{-1} \neq \emptyset$  then  $g(C_{-1} \cup B_1) = -1$ .*

Since  $g(C_1, B_{-1}) \neq 2$  (because  $g$  is good), and  $g(C_1 \cup Z) = 0$  for all  $Z \subseteq B_{-1}$  with  $Z \neq \emptyset, B_{-1}$  by (6), 2.1 implies that  $g(C_1) + (-1)^{|k_1-1|} g(C_1 \cup B_{-1}) \leq 1$ . But  $g(C_1) = 1$  (since  $C_1 \neq \emptyset$ ), and  $k_1$  is odd, and so  $g(C_1 \cup B_{-1}) = 1$ . Similarly if  $C_{-1} \neq \emptyset$  then  $g(C_{-1} \cup B_1) = -1$ . This proves (7).

(8) *One of  $C_1, C_{-1}$  is empty.*

Suppose they are both nonempty. By 2.1,

$$g(C, D) = \sum_{Z \subseteq D} (-1)^{|Z|} g(C \cup Z).$$

But for  $Z \subseteq D$ ,  $g(C \cup Z) \neq 0$  only if  $Z$  includes one of  $D_1, D_{-1}$  by (6), and only if one of  $Z \cap B_1, Z \cap B_{-1}$  is empty (from the definition of  $C_1, C_{-1}$ ); that is, only if  $Z$  is one of  $D_1, D_{-1}$ . These two sets are distinct, since they are nonempty. Consequently

$$g(C, D) = (-1)^{|D_1|} g(B_1 \cup C_{-1}) + (-1)^{|D_{-1}|} g(B_{-1} \cup C_1)$$

and so by (7),  $g(C, D) = (-1)^{|D_1|+1} + (-1)^{|D_{-1}|}$ . Since  $|g(C, D)| \leq 1$  (because  $g$  is good) it follows that  $|D_1|, |D_{-1}|$  have the same parity.

Choose  $u \in D_1$  and  $v \in D_{-1}$ . Then by 2.1,

$$g(C \cup \{u\}, D \setminus \{u, v\}) = \sum_{Z \subseteq D \setminus \{u, v\}} (-1)^{|Z|} g(C \cup \{u\} \cup Z).$$

But for  $Z \subseteq D \setminus \{u, v\}$ ,  $g(C \cup \{u\} \cup Z) \neq 0$  only if  $Z = D_1 \setminus \{u\}$  (by (6) and the definition of  $C_1, C_{-1}$ ) and so

$$g(C \cup \{u\}, D \setminus \{u, v\}) = (-1)^{|D_1 \setminus \{u\}|} g(B_1 \cup C_{-1}) = (-1)^{|D_1|}.$$

Similarly

$$g(C \cup \{v\}, D \setminus \{u, v\}) = (-1)^{|D_2 \setminus \{v\}|} g(B_{-1} \cup C_1) = (-1)^{|D_{-1}|+1}.$$

Since  $|D_1|, |D_{-1}|$  have the same parity, one of  $g(C \cup \{u\}, D \setminus \{u, v\}), g(C \cup \{v\}, D \setminus \{u, v\})$  equals 1 and the other equals  $-1$ , contradicting that  $g$  is good. This proves (8).

From (8) we may assume that  $C_{-1} = \emptyset$  (replacing  $g$  by  $-g$  if necessary).

(9)  $|D_1|$  is odd.

To prove this, we may assume that  $C_1 \neq \emptyset$ , since  $|B_1|$  is odd. By 2.1,

$$g(C_1, B_{-1} \cup D_1) = \sum_{Z \subseteq B_{-1} \cup D_1} (-1)^{|Z|} g(C_1 \cup Z).$$

But, by (6), for  $Z \subseteq B_{-1} \cup D_1$ ,  $g(C_1 \cup Z)$  is nonzero only if  $Z \subseteq D_1$  or  $Z = B_{-1}$ ; and then it has value 1 if  $Z \subseteq D_1$  and  $Z \neq D_1$ ; 0 if  $Z = D_1$ ; and 1 if  $Z = B_{-1}$ . Thus  $g(C_1, B_{-1} \cup D_1) = (-1)^{|D_1|+1} + (-1)^{|B_{-1}|}$  and since  $|B_{-1}|$  is odd by (5), and  $|g(C_1, B_{-1} \cup D_1)| \leq 1$  since  $g$  is good, it follows that  $|D_1|$  is odd. This proves (9).

Now  $|C_1| \leq |B_1| - 2$  as we saw. Choose  $u \in D_1$  and  $v \in B_{-1}$ , and let  $W = (D_1 \cup B_{-1}) \setminus \{u, v\}$ . By 2.1,

$$g(C_1 \cup \{u\}, W) = \sum_{Z \subseteq W} (-1)^{|Z|} g(C_1 \cup \{u\} \cup Z).$$

But for  $Z \subseteq W$ ,  $g(C_1 \cup \{u\} \cup Z)$  is nonzero only if  $Z \subseteq D_1$ , and in that case it has value 1 if  $Z \neq D_1 \setminus \{u\}$ , and 0 if  $Z = D_1 \setminus \{u\}$ . Since  $|D_1| \geq 2$ , it follows that

$$g(C_1 \cup \{u\}, W) = (-1)^{|D_1|} = -1$$

since  $|D_1|$  is odd by (9). On the other hand, by 2.1,

$$g(C_1 \cup \{v\}, W) = \sum_{Z \subseteq W} (-1)^{|Z|} g(C_1 \cup \{v\} \cup Z).$$

We claim that  $g(C_1 \cup \{v\}, W) = 1$ . To see this there are two cases, depending whether  $C_1 \neq \emptyset$  or not. First, suppose that  $C_1 \neq \emptyset$ . Then for  $Z \subseteq W$ ,  $g(C_1 \cup \{v\} \cup Z)$  is nonzero only if  $Z = B_{-1} \setminus \{v\}$ , by (6) and the maximality of  $C_1$ ; so

$$g(C_1 \cup \{v\}, W) = (-1)^{|B_{-1}|-1} g(C_1 \cup B_{-1}) = 1,$$

by (7) and (3), contradicting that  $g$  is good. Now suppose that  $C_1 = \emptyset$ . Then, again by (6), for  $Z \subseteq W$ ,  $g(C_1 \cup \{v\} \cup Z)$  is nonzero only if  $Z \subsetneq B_{-1} \setminus \{v\}$ , and in that case it has value  $-1$ . Consequently

$$g(C_1 \cup \{v\}, W) = (-1)^{|B_{-1} \setminus \{v\}|} = 1,$$

again contradicting that  $g$  is good. This proves 3.4. ■

## 4 The value on the null set

In this section we prove 1.6, thereby completing the inductive proof of 1.4. We need to show that if  $g$  is a good counter on a ternary graph  $G$ , then  $|g(\emptyset)| \leq 1$ . The proof is divided into several steps. We may assume the statement is false, for a contradiction; and by replacing  $g$  by  $-g$  if necessary, we may assume that  $g(\emptyset) \geq 2$ . Throughout this section,  $G$  is a counterexample to 1.6, and  $g$  is a good counter on  $G$ , with  $g(\emptyset) \geq 2$ .

**4.1** *The following hold:*

- $g(\emptyset) = 2$ ;
- $g(\{v\}) = 1$  for every vertex  $v \in V(G)$ ; and
- $G$  is connected.

**Proof.** Let  $v \in V(G)$ ; since  $g$  is good, it follows that  $|g(\{v\})| \leq 1$ , and so 2.3 implies that  $g(\{v\}) = 1$  and  $g(\emptyset) = 2$ . This proves the first two statements.

Suppose that  $G$  is not connected, let  $G_1$  be a component of  $G$  and let  $G_2$  be obtained from  $G$  by deleting  $V(G_1)$ . Since  $f_{G_1}(\emptyset) = \pm g(\emptyset, V(G_2))$ , and  $g$  is good, it follows that  $|f_{G_1}(\emptyset)| \leq 1$ , and similarly  $|f_{G_2}(\emptyset)| \leq 1$ . But

$$g(\emptyset) = \pm f_G(\emptyset) = \pm f_{G_1}(\emptyset) f_{G_2}(\emptyset),$$

a contradiction. This proves the third statement, and so proves 4.1. ▀

In particular, if  $u, v \in V(G)$  are distinct, then since  $g(\{u\}) = 1$  by the second statement of 4.1, it follows that  $g(\{u, v\}) \in \{0, 1\}$  by 2.3. Let  $H$  be the graph with vertex set  $V(G)$  in which distinct  $u, v$  are adjacent if  $g(\{u, v\}) = 1$ .

**4.2** *Every component of  $H$  is a complete graph, and  $H$  has at least two and at most four components.*

**Proof.** Suppose the first statement is false. Then there are three distinct vertices  $u, v, w \in V(H)$  such that  $uv, vw \in E(H)$  and  $uw \notin E(H)$ . From 2.3,  $g(\{u, w\}) = 0$ . Now

$$\begin{aligned} g(\emptyset, \{w\}) &= g(\emptyset) - g(\{w\}) = 1, \\ g(\{v\}, \{w\}) &= g(\{v\}) - g(\{v, w\}) = 0 \\ g(\{u, v\}, \{w\}) &= g(\{u, v\}) - g(\{u, v, w\}); \end{aligned}$$

and by 2.5,  $g(\{u, v\}, \{w\}) \neq 1$ . Consequently  $g(\{u, v, w\}) = 1$ . But then  $g(\{w\}) = 1$ ,  $g(\{u, w\}) = 0$  and  $g(\{u, v, w\}) = 1$ , contrary to 2.5. This proves that every component of  $H$  is a complete graph.

Since each edge of  $H$  joins two vertices that are nonadjacent in  $G$ , it follows that  $H$  has at least two components. Suppose it has at least five. Since  $G$  is connected, there is a vertex of  $H$  that has neighbours (in  $G$ ) in at least two components of  $H$ . Thus we can choose  $v_1, \dots, v_5 \in V(G)$ , all in different components of  $H$ , where  $v_1$  is adjacent (in  $G$ ) to  $v_2, v_3$ . Let  $a, b, c \in \{v_1, \dots, v_5\}$  be distinct. Since  $|g(\emptyset, \{a, b, c\})| \leq 1$ , and  $g(\{a, b\}) = 0$  (because  $g(\{a, b\}) \neq 1$  since  $a, b$  belong to different components of  $H$ , and  $g(\{a, b\}) \neq -1$  by 2.3), and the same for  $\{a, c\}$  and  $\{b, c\}$ , it follows from 2.1 that  $|2 - 3 + 0 - g(\{a, b, c\})| \leq 1$ , and so  $g(\{a, b, c\}) \neq 1$ . Hence  $g(\{a, b, c\}) \in \{0, -1\}$  for every triple  $a, b, c$  of distinct members of  $\{v_1, \dots, v_5\}$ .

Note that since  $v_1v_2, v_1v_3 \in E(G)$ , it follows that  $g(\{v_1, v_2, v_i\}) = 0$  for every  $i \in \{3, 4, 5\}$  and  $g(\{v_1, v_3, v_j\}) = 0$  for every  $j \in \{2, 4, 5\}$ . Let  $\mathcal{T}$  be the set of all subsets  $T \subseteq \{v_1, \dots, v_5\}$  with  $|T| = 3$  and  $g(T) = -1$ . Thus  $g(T) = 0$  for all triples  $T \notin \mathcal{T}$ . Since  $|g(\emptyset, \{v_1, v_2, v_3, v_4\})| \leq 1$ , it follows from 2.1 that  $\{v_2, v_3, v_4\} \in \mathcal{T}$ , and similarly  $\{v_2, v_3, v_5\} \in \mathcal{T}$ .

Suppose that  $\{v_1, v_4, v_5\} \notin \mathcal{T}$ . Now 2.1 implies that

$$g(\emptyset, \{v_1, v_2, v_4, v_5\}) = 2 - 4 + 0 - g(\{v_2, v_4, v_5\}),$$

and so  $\{v_2, v_4, v_5\} \in \mathcal{T}$ , and similarly  $\{v_3, v_4, v_5\} \in \mathcal{T}$ . But then

$$g(\{v_5\}, \{v_2, v_3, v_4\}) = -2 - g(v_2, v_3, v_4, v_5) \leq -1$$

and  $g(\{v_1\}, \{v_2, v_3, v_4\}) = 1$ , contradicting that  $g$  is good. Thus  $\{v_1, v_4, v_5\} \in \mathcal{T}$ .

If also  $\{v_2, v_4, v_5\} \in \mathcal{T}$  then  $g(\{v_4, v_5\}, \{v_1, v_2\}) = 2$ , contradicting that  $g$  is good; so  $\{v_2, v_4, v_5\} \notin \mathcal{T}$ , and similarly  $\{v_3, v_4, v_5\} \notin \mathcal{T}$ . Since  $g(\{v_2, v_3\}, \{v_4, v_5\}) \leq 1$ , it follows that  $g(\{v_2, v_3, v_4, v_5\}) = -1$ . But then  $g(\{v_4\}, \{v_2\}) = 1$ ,  $g(\{v_4, v_5\}, \{v_2\}) = 0$  and  $g(\{v_3, v_4, v_5\}, \{v_2\}) = 1$ , contrary to 2.5. This proves 4.2. ▀

**4.3** Let  $C_1, C_2$  be distinct components of  $H$ , and let  $X \subseteq C_1 \cup C_2$ . Suppose that

- $X \cap C_1, X \cap C_2 \neq \emptyset$ ;
- $g(X) \neq 0$ ; and
- for all  $X' \subseteq X$ , if  $g(X') \neq 0$  then either  $X' = X$  or  $X' \subseteq C_1$  or  $X' \subseteq C_2$ .

If  $|X \cap C_1| > 1$  then there is a subset  $B \subseteq X \cap C_1$  with  $g(B) = 0$ .

**Proof.** Let  $X_i = X \cap C_i$  for  $i = 1, 2$ ; and suppose there is no  $B \subseteq X_1$  with  $g(B) = 0$ . From 2.3 it follows that  $g(B) = 1$  for all nonempty subsets  $B$  of  $X_1$ , and in particular,  $g(X_1) = 1$ . Let  $g(X) = i = \pm 1$ . Because of the third bullet of the hypothesis, 2.1 implies that

$$g(X_1, X_2) = \sum_{Z \subseteq X_2} (-1)^{|Z|} g(X_1 \cup Z) = g(X_1) + (-1)^{|X_2|} i;$$

and since  $g(X_1, X_2) \leq 1$ , it follows that  $(-1)^{|X_2|} i = -1$ , that is,  $|X_2|$  is odd if  $i = 1$ , and even if  $i = -1$ . Choose  $u \in X_1$  and  $v \in X_2$ ; then by 2.1,  $g(X_1 \setminus \{u\}, X_2 \setminus \{v\}) = 1$  (since  $|X_1| > 1$ ),  $g(X_1 \cup \{v\} \setminus \{u\}, X_2 \setminus \{v\}) = 0$ , and by 2.1,

$$g(X_1 \cup \{v\}, X_2 \setminus \{v\}) = \sum_{Z \subseteq X_2 \setminus \{v\}} (-1)^{|Z|} g(X_1 \cup Z \setminus \{v\}) = (-1)^{|X_2|-1} g(X) = 1,$$

contrary to 2.5. This proves 4.3. ▀

Let  $C$  be a component of  $H$ , and let  $D \subseteq C$ . We say that  $B \subseteq D$  is a *base* of  $D$  if  $g(B) \neq 1$  and there is no  $B' \subseteq D$  with  $|B'| < |B|$  and with  $g(B') \neq 1$ .

**4.4** Let  $C$  be a component of  $H$ , and let  $D \subseteq C$ .

- If there is a vertex  $v$  of  $G$  such that all its neighbours belong to  $D$ , then  $D$  has a base.
- If  $B$  is a base of  $D$  then  $g(B) = 0$ , and  $|B|$  is even and at least four.
- If  $D$  has a base, of cardinality  $k$  say, then every subset of  $D$  of cardinality  $k + 1$  includes two bases of  $D$ , and so every vertex of  $D$  belongs to a base of  $D$ .
- If  $D$  has a base, of cardinality  $k$ , then there is a partition of  $D$  into nonempty sets  $D_1, \dots, D_n$ , such that for all distinct  $u, v \in D$ , there is a base of  $D$  containing both  $u, v$  if and only if  $u, v$  do not belong to the same set  $D_i$ ; and consequently  $n \geq k$ .

**Proof.** For the first statement, suppose that all neighbours of  $v$  belong to  $D$ . If  $V(G) = C \cup \{v\}$ , then  $v$  is adjacent to all other vertices (since no vertex has degree zero, by 2.4), contradicting that  $g(\emptyset) = 2$ . Thus we may choose  $u \notin C \cup \{v\}$ . By 2.4,  $g(\{u\}, D) = 0$ , but  $g(\{u\}) = 1$ , and so by 2.1, there exists a nonempty subset  $Z \subseteq D$  such that  $g(Z \cup \{u\}) \neq 0$ . Since  $C$  is the vertex set of a component of  $H$ , it follows that  $|Z| \geq 2$ . From 4.3, there exists  $B \subseteq Z$  with  $g(B) = 0$ . This proves the first statement.

For the second, let  $B$  be a base of  $D$ . Then  $g(B) \neq 1$  by hypothesis, and in particular  $|B| \geq 3$ , since  $B \subseteq C$ . For every  $B' \subseteq B$  with  $B' \neq \emptyset, B$ , we have  $g(B') = 1$ , and since there is such a choice of  $B'$  with  $|B'| = |B| - 1$ , 2.3 implies that  $g(B) \neq -1$ ; and hence  $g(B) = 0$  since  $g$  is good. But by 2.2,

$$g(\emptyset, B) = \sum_{Z \subseteq B} (-1)^{|Z|} (g(Z) - 1) = (g(\emptyset) - 1) + (-1)^{|B|} (g(B) - 1) = 1 - (-1)^{|B|},$$

and so  $|B|$  is even. This proves the second statement.

For the third, let  $B$  be a base of  $D$ , with  $|B| = k$  say; it suffices to prove that for all  $v \in D \setminus B$ ,  $B \cup \{v\}$  includes at least two bases of  $D$ . Let  $X = B \cup \{v\}$ , and choose  $u \in B$ . Thus  $g(X \setminus \{u, v\}) = 1$  and  $g(X \setminus \{v\}) = 0$ , so by 2.5,  $g(X) \neq 1$ . We may assume that  $g(X \setminus \{u\}) = 1$ , and so by 2.3,  $g(X) = 0$ . By 2.1,  $g(\emptyset, X \setminus \{u, v\}) = 1$  and  $g(\{u, v\}, X \setminus \{u, v\}) = 1$ , so by 2.5,  $g(\{v\}, X \setminus \{u, v\}) = 1$ . Hence by 2.1, since  $|X| \geq 3$ , there exists  $Z \subseteq X \setminus \{u, v\}$  with  $g(Z \cup \{v\}) \neq 1$ . Then  $|Z| \leq |B|$ , and since  $B$  is a base for  $D$ , it follows that  $Z$  is minimal with  $g(Z) \neq 1$ , and hence  $Z$  is another base for  $D$ . This proves the third statement.

The fourth statement follows from 2.7. This proves 4.4. ■

We call a partition  $D_1, \dots, D_n$  as in the fourth statement of 4.4 the *induced partition* of  $D$ , and the sets  $D_1, \dots, D_n$  are called its *classes*. (If the partition exists then it is unique, as is easily seen.)

**4.5** Let  $C_1, C_2$  be distinct components of  $H$ , and for  $i = 1, 2$ , let  $D_i \subseteq C_i$ , including a base for  $D_i$ . Then for one of  $i = 1, 2$ , there is a class of the induced partition of  $D_i$  that meets all edges between  $D_1$  and  $D_2$ .

**Proof.** Let the induced partition of  $D_1$  have classes  $P_1, \dots, P_m$ , and let the induced partition of  $D_2$  have classes  $Q_1, \dots, Q_n$ . We may assume that there is no  $i \in \{1, \dots, m\}$  such that all edges between

$D_1, D_2$  have an end in  $P_i$ , and there is no  $j \in \{1, \dots, n\}$  similarly. By König's theorem, there exist distinct  $i_1, i_2 \in \{1, \dots, m\}$  and distinct  $j_1, j_2 \in \{1, \dots, n\}$  such that there is an edge between  $P_{i_1}$  and  $Q_{j_1}$ , and an edge between  $P_{i_2}$  and  $Q_{j_2}$ . Hence there is a base  $B_1$  for  $D_1$  and a base  $B_2$  for  $D_2$ , such that there are two edges of  $G$  between  $B_1, B_2$  with no end in common.

(1) Suppose that there exists  $M_1 \subseteq B_1$  with  $g(B_2 \cup M_1) \neq 0$ , and choose  $M_1$  maximal with this property. Then  $|M_1| \leq |B_1| - 2$ , and  $g(B_2 \cup M_1) = -1$ , and  $|M_1|$  is odd.

Since there are two edges of  $G$  between  $B_1, B_2$  with no end in common, and both have an end in  $B_2$ , it follows that neither has an end in  $M_1$ , and so  $|M_1| \leq |B_1| - 2$ . Let  $A_1 = B_1 \setminus M_1$ . By 2.1,

$$g(M_1, B_2) = \sum_{Z \subseteq B_2} (-1)^{|Z|} g(M_1 \cup Z).$$

But for  $Z \subseteq B_2$ ,  $g(M_1 \cup Z) \neq 0$  only if  $Z = \emptyset$  or  $Z = B_2$ , by 4.3. Consequently  $g(M_1, B_2) = g(M_1) + (-1)^{|B_2|} g(M_1 \cup B_2)$ . But  $g(M_1) = 1$  and  $|B_2|$  is even, so  $g(M_1 \cup B_2) = -1$  since  $g$  is good. Now by 2.1,

$$g(M_1, A_1 \cup B_2) = \sum_{Z \subseteq A_1 \cup B_2} (-1)^{|Z|} g(M_1 \cup Z).$$

But for  $Z \subseteq A_1 \cup B_2$ ,  $g(M_1 \cup Z) \neq 0$  only if  $Z \subseteq A_1$  or  $Z = B_2$ ; and so

$$g(M_1, A_1 \cup B_2) = (-1)^{|A_1|} (g(B_1) - 1) + (-1)^{|B_2|} g(M_1 \cup B_2).$$

Since  $|B_2|$  is even,  $g(B_1) = 0$  and  $g(M_1 \cup B_2) = -1$ , it follows that  $g(M_1, A_1 \cup B_2) = (-1)^{|A_1|+1} - 1$ , and so  $|A_1|$  is odd, and therefore so is  $|M_1|$ . This proves (1).

(2) There do not exist  $M_1 \subseteq B_1$  and  $M_2 \subseteq B_2$  with  $g(B_2 \cup M_1), g(B_1 \cup M_2) \neq 0$  and with  $M_1, M_2$  both nonempty.

Suppose such sets  $M_1, M_2$  exist and choose them maximal. Let  $A_i = B_i \setminus M_i$  for  $i = 1, 2$ . By (1),  $g(B_2 \cup M_1), g(B_1 \cup M_2) = -1$ , and  $|M_1|, |M_2|$  are odd. Thus  $|A_1|$  and  $|A_2|$  are odd, and so  $g(M_1 \cup M_2, A_1 \cup A_2) = 2$  by 2.1, a contradiction. This proves (2).

(3)  $g(X) = 0$  for all  $X \subseteq B_1 \cup B_2$  with  $X \cap B_1, X \cap B_2$  both nonempty.

Suppose not; then from 4.3, and by exchanging  $C_1, C_2$  if necessary, we may assume that there exists  $M_1 \subseteq B_1$ , nonempty, with  $g(B_2 \cup M_1) \neq 0$ . Choose  $M_1$  maximal. By (1),  $g(B_2 \cup M_1) = -1$  and  $|M_1|$  is odd. Let  $A_1 = B_1 \setminus M_1$ , and choose  $u \in A_1$ . Choose  $v \in B_2$ . Then by 2.1, since  $A_1 \setminus \{u\} \neq \emptyset$ , it follows that  $g(M_1 \cup \{u\}, (A_1 \cup B_2) \setminus \{u, v\}) = -1$  and  $g(M_1 \cup \{v\}, (A_1 \cup B_2) \setminus \{u, v\}) = 1$ , contradicting that  $g$  is good. This proves (3).

From (3), 2.1 implies that  $g(\emptyset, B_1 \cup B_2) = -2$ , a contradiction. This proves 4.5. ■

**4.6** Let  $C_1, C_2$  be distinct components of  $H$ , and suppose there is a base for  $C_2$ . Let  $D_1, \dots, D_n$  be the induced partition of  $C_2$ . Then there is no  $i \in \{D_1, \dots, D_n\}$  such that every edge of  $G$  between  $C_2$  and  $V(G) \setminus (C_1 \cup C_2)$  has an end in  $D_i$ .



**Proof.** Suppose there is such a value of  $i$ , say  $i = 1$ . Let  $A_1$  be the set of vertices in  $C_1$  with neighbours in  $C_2$ . Now  $n \geq 4$  (by the second and last statements of 4.4); choose  $v \in D_2$ . Thus all neighbours of  $v$  belong to  $C_1$ , and hence to  $A_1$ . By the first statement of 4.4, there is a base for  $A_1$ . By 4.5, there is a set  $X$  that meets all edges between  $A_1$  and  $C_2$ , and  $X$  is either a class of the induced partition of  $C_2$  or a class of the induced partition of  $A_1$ . The first is impossible since there are at least four classes of the induced partition of  $C_2$ , and each such class different from  $D_1$  meets an edge between  $C_2$  and  $A_1$  (because it meets some edge, and it has no edge to  $V(G) \setminus (C_1 \cup C_2)$  from the choice of  $D_1$ ). Also the second is impossible, since each class of the induced partition of  $A_1$  has an edge to  $C_2$ , from the definition of  $A_1$ . This proves 4.6.  $\blacksquare$

Now we complete the proof of 1.6, which we restate:

**4.7** *If  $g$  is a good counter on a ternary graph  $G$ , then  $|g(\emptyset)| \leq 1$ .*

**Proof.** In the same notation as before, we know that  $H$  has two, three or four components. Suppose it has only two, say  $C_1, C_2$ . By the first statement of 4.4, there are bases for  $C_1$  and for  $C_2$ , contrary to 4.6.

Now suppose that  $H$  has exactly three components  $C_1, C_2, C_3$ . By 2.6 we may assume that some vertex  $v \in C_2$  has no neighbour in  $C_1$ , and so by 4.4, there is a base for  $C_3$ . Suppose that there is also a base for  $C_2$ . By 4.5, by exchanging  $C_2, C_3$  if necessary, we may assume that there is a class of the induced partition of  $C_2$  that meets all edges between  $C_2, C_3$ , contrary to 4.6. Thus, neither of  $C_1, C_2$  have bases. By 4.4, every vertex in  $C_1 \cup C_2$  has a neighbour in  $C_3$ . We recall that  $v \in C_2$  has no neighbour in  $C_1$ . Since  $C_1$  has no base, it follows that  $g(C_1) = 1$ , and so by 2.4,  $v$  has a neighbour,  $u$  say, with no neighbour in  $C_1$ . But then all neighbours of  $u$  are in  $C_2$ , and so by 4.4, there is a base for  $C_2$ , a contradiction.

Finally, suppose that  $H$  has four components  $C_1, \dots, C_4$ . Let  $K$  be the graph with vertex set  $\{1, \dots, 4\}$  in which distinct  $i, j$  are adjacent if there is an edge of  $G$  between  $C_i, C_j$ . Since  $G$  is connected, it follows that every vertex of  $K$  has nonzero degree. Suppose that  $K$  has a 2-edge matching; then by renumbering  $C_1, \dots, C_4$  we may assume that there exist  $u_i \in C_i$  for  $1 \leq i \leq 4$  such that  $u_1 u_2, u_3 u_4 \in E(G)$ . But then  $g(\emptyset, \{u_1, u_2, u_3, u_4\}) = -2$  by 2.1, a contradiction. Thus  $K$  has no 2-edge matching, and since every vertex of  $K$  has nonzero degree, we may assume that every edge of  $K$  is incident with 1, and so all edges of  $G$  have an end in  $C_1$ .

For  $i = 2, 3, 4$ , let  $X_i$  be the set of vertices in  $C_1$  with no neighbour in  $C_i$ . By the first statement of 4.4, there is a base for  $C_1$ . By 4.6, there is no base for  $C_2$ , and similarly none for  $C_3, C_4$ ; and so by the first statement of 4.4, every vertex of  $C_1$  has neighbours in at least two of  $C_2, C_3, C_4$ . In particular, for all distinct  $i, j \in \{2, 3, 4\}$  every vertex in  $X_i$  has a neighbour in  $C_j$ .

Since  $g(C_2) \neq 0$ , 2.4 implies that for all distinct  $i, j \in \{2, 3, 4\}$ , every vertex in  $C_i$  has a neighbour in  $X_j$ . Make a digraph  $J$  with vertex set  $C_2 \cup C_3 \cup C_4$  in which for  $i = 2, 3, 4$  and  $u \in C_i$  and  $v \in C_{i+1}$  (where  $C_5$  means  $C_2$ ), there is an edge of  $J$  from  $u$  to  $v$  if  $u, v$  has a common neighbour in  $X_{i-1}$  (where  $X_1$  means  $X_4$ ). Every vertex has positive outdegree in  $J$ , and so  $J$  has an induced directed cycle. Let  $K$  be such a cycle, with vertices (in order):

$$a_1, b_1, c_1, a_2, b_2, c_2, \dots, a_k, b_k, c_k, a_1$$

where  $a_1, \dots, a_k \in C_2$ ,  $b_2, \dots, b_k \in C_3$  and  $c_1, \dots, c_k \in C_4$ . For each  $i$  with  $1 \leq i \leq k$ , there exists  $x_i \in X_4$  adjacent in  $G$  to  $a_i, b_i$ , and  $y_i \in X_2$  adjacent to  $b_i, c_i$ , and  $z_i \in X_3$  adjacent to  $c_i, a_{i+1}$  (where

$a_{k+1}$  means  $a_1$ ). Also, for each such  $i$ ,  $x_i$  has no other neighbours in  $V(K)$ ; it is nonadjacent to each  $a_j$  because  $x_i \in X_4$ , and nonadjacent to the remaining vertices of  $V(K)$  since  $K$  is induced. A similar statement holds for the  $y_i$ 's and  $z_i$ 's. Consequently the subgraph of  $G$  induced on

$$\{a_i, b_i, c_i, x_i, y_i, z_i : 1 \leq i \leq k\}$$

is an induced cycle of length  $6k$ , contradicting that  $G$  is ternary. This proves that  $H$  does not have four components, and so proves 4.7 and hence 1.4. ■

## References

- [1] M. Bonamy, P. Charbit and S. Thomassé, “Graphs with large chromatic number induce  $3k$ -cycles”, [arXiv:1408.2172](https://arxiv.org/abs/1408.2172).
- [2] M. Bousquet-Mélou, S. Linusson and E. Nevo, “On the independence complex of square grids”, *J. Algebr. Comb.*, **27** (2008), 423–450.
- [3] G. Chen and A. Saito, “Graphs with a cycle of length divisible by three”, *J. Combinatorial Theory, Ser. B*, **60** (1994), 277–292.
- [4] P. Fendley, K. Schoutens and H. van Eerten, “Hard squares with negative activity”, *J. Phys. A*, **38** (2005), 31–322.
- [5] G. Gauthier, *Graphs with no cycle length divisible by three*, Ph. D. thesis, Princeton, 2017.
- [6] A. Hatcher, *Algebraic Topology*, Cambridge University Press (2009).
- [7] J. Jonsson, “Hard squares with negative activity and rhombus tilings of the plane”, *Electronic J. Combinatorics*, **13** (2006), #R67.
- [8] G. Kalai, “When do a few colors suffice?”, <https://gilkalai.wordpress.com/2014/12/19/when-a-few-colors-suffice>.
- [9] A. Scott and P. Seymour, “Induced subgraphs of graphs with large chromatic number. X. Holes with specific residue”, *Combinatorica*, to appear, [arXiv:1705.04609](https://arxiv.org/abs/1705.04609).
- [10] A. Scott and A. D. Sokal, “The repulsive lattice gas, the independent-set polynomial, and the Lovász local lemma”, *J. Statistical Physics*, **118** (2005), 1151–1261.