

CLUSTERED COLOURING OF GRAPH CLASSES WITH BOUNDED TREEDEPTH OR PATHWIDTH

Sergey Norin[†] Alex Scott[‡] David R. Wood[¶]

Abstract. The *clustered chromatic number* of a class of graphs is the minimum integer k such that for some integer c every graph in the class is k -colourable with monochromatic components of size at most c . We determine the clustered chromatic number of any minor-closed class with bounded treedepth, and prove a best possible upper bound on the clustered chromatic number of any minor-closed class with bounded pathwidth. As a consequence, we determine the fractional clustered chromatic number of every minor-closed class.

1 Introduction

This paper studies improper vertex colourings of graphs with bounded monochromatic degree or bounded monochromatic component size. This topic has been extensively studied recently [1, 3, 4, 7, 9, 11–20, 22–24]; see [25] for a survey.

A *k -colouring* of a graph G is a function that assigns one of k colours to each vertex of G . In a coloured graph, a *monochromatic component* is a connected component of the subgraph induced by all the vertices of one colour.

A colouring has *defect* d if each monochromatic component has maximum degree at most d . The *defective chromatic number* of a graph class \mathcal{G} , denoted by $\chi_{\Delta}(\mathcal{G})$, is the minimum integer k such that, for some integer d , every graph in \mathcal{G} is k -colourable with defect d .

A colouring has *clustering* c if each monochromatic component has at most c vertices. The *clustered chromatic number* of a graph class \mathcal{G} , denoted by $\chi_{\star}(\mathcal{G})$, is the minimum integer k such that, for some integer c , every graph in \mathcal{G} has a k -colouring with clustering c . We shall consider such colourings, where the goal is to minimise the number of colours, without optimising the clustering value.

Every colouring of a graph with clustering c has defect $c - 1$. Thus $\chi_{\Delta}(\mathcal{G}) \leq \chi_{\star}(\mathcal{G})$ for every class \mathcal{G} .

The following is a well-known and important example in defective and clustered graph colouring. Let T be a rooted tree. The *depth* of T is the maximum number of vertices

February 10, 2021

[†] Department of Mathematics and Statistics, McGill University, Montréal, Canada (snorin@math.mcgill.ca). Supported by NSERC grant 418520.

[‡] Mathematical Institute, University of Oxford, Oxford, U.K. (scott@maths.ox.ac.uk).

[¶] School of Mathematics, Monash University, Melbourne, Australia (david.wood@monash.edu). Supported by the Australian Research Council.

on a root-to-leaf path in T . The *closure* of T is obtained from T by adding an edge between every ancestor and descendant in T . For $h, k \geq 1$, let $C\langle h, k \rangle$ be the closure of the complete k -ary tree of depth h , as illustrated in Figure 1.




Figure 1: The standard example $C\langle 4, 2 \rangle$.

It is well known and easily proved (see [25]) that there is no $(h - 1)$ -colouring of $C\langle h, k \rangle$ with defect $k - 1$, which implies there is no $(h - 1)$ -colouring of $C\langle h, k \rangle$ with clustering k . This says that if a graph class \mathcal{G} includes $C\langle h, k \rangle$ for all k , then the defective chromatic number and the clustered chromatic number are at least h . Put another way, define the *tree-closure-number* of a graph class \mathcal{G} to be

$$\text{tcn}(\mathcal{G}) := \min\{h : \exists k C\langle h, k \rangle \notin \mathcal{G}\} = \max\{h : \forall k C\langle h, k \rangle \in \mathcal{G}\} + 1;$$

then

$$\chi_*(\mathcal{G}) \geq \chi_\Delta(\mathcal{G}) \geq \text{tcn}(\mathcal{G}) - 1.$$

Our main result, Theorem 1 below, establishes a converse result for minor-closed classes with bounded treedepth. First we explain these terms. A graph H is a *minor* of a graph G if a graph isomorphic to H can be obtained from some subgraph of G by contracting edges. A class of graphs \mathcal{M} is *minor-closed* if for every graph $G \in \mathcal{M}$ every minor of G is in \mathcal{M} , and \mathcal{M} is *proper* minor-closed if, in addition, some graph is not in \mathcal{M} . The *connected treedepth* of a graph H , denoted by $\overline{\text{td}}(H)$, is the minimum depth of a rooted tree T such that H is a subgraph of the closure of T . This definition is a variant of the more commonly used definition of the *treedepth* of H , denoted by $\text{td}(H)$, which equals the maximum connected treedepth of the connected components of H . (See [21] for background on treedepth.) If H is connected, then $\text{td}(H) = \overline{\text{td}}(H)$. In fact, $\text{td}(H) = \overline{\text{td}}(H)$ unless H has two connected components H_1 and H_2 with $\text{td}(H_1) = \text{td}(H_2) = \text{td}(H)$, in which case $\overline{\text{td}}(H) = \text{td}(H) + 1$. It is convenient to work with connected treedepth to avoid this distinction. A class of graphs has *bounded treedepth* if there exists a constant c such that every graph in the class has treedepth at most c .

Theorem 1. *For every minor-closed class \mathcal{G} with bounded treedepth,*

$$\chi_{\Delta}(\mathcal{G}) = \chi_{\star}(\mathcal{G}) = \text{tcn}(\mathcal{G}) - 1.$$

Our second result concerns pathwidth. A *path-decomposition* of a graph G consists of a sequence (B_1, \dots, B_n) , where each B_i is a subset of $V(G)$ called a *bag*, such that for every vertex $v \in V(G)$, the set $\{i \in [1, n] : v \in B_i\}$ is an interval, and for every edge $vw \in E(G)$ there is a bag B_i containing both v and w . Here $[a, b] := \{a, a+1, \dots, b\}$. The *width* of a path decomposition (B_1, \dots, B_n) is $\max\{|B_i| : i \in [1, n]\} - 1$. The *pathwidth* of a graph G is the minimum width of a path-decomposition of G . Note that paths (and more generally caterpillars) have pathwidth 1. A class of graphs has *bounded pathwidth* if there exists a constant c such that every graph in the class has pathwidth at most c .

Theorem 2. *For every minor-closed class \mathcal{G} with bounded pathwidth,*

$$\chi_{\Delta}(\mathcal{G}) \leq \chi_{\star}(\mathcal{G}) \leq 2 \text{tcn}(\mathcal{G}) - 2.$$

Theorems 1 and 2 are respectively proved in Sections 2 and 3. These results are best possible and partially resolve a number of conjectures from the literature, as we now explain.

Ossona de Mendez et al. [23] studied the defective chromatic number of minor-closed classes. For a graph H , let \mathcal{M}_H be the class of H -minor-free graphs (that is, not containing H as a minor). Ossona de Mendez et al. [23] proved the lower bound, $\chi_{\Delta}(\mathcal{M}_H) \geq \overline{\text{td}}(H) - 1$ and conjectured that equality holds.

Conjecture 3 ([23]). *For every graph H ,*

$$\chi_{\Delta}(\mathcal{M}_H) = \overline{\text{td}}(H) - 1.$$

Note that Conjecture 3 is known to hold in some special cases. Edwards et al. [9] proved it if $H = K_t$; that is, $\chi_{\Delta}(\mathcal{M}_{K_t}) = t - 1$, which can be thought of as a defective version of Hadwiger's Conjecture; see [24] for an improved bound on the defect in this case. Ossona de Mendez et al. [23] proved Conjecture 3 if $\overline{\text{td}}(H) \leq 3$ or if H is a complete bipartite graph. In particular, $\chi_{\Delta}(\mathcal{M}_{K_{s,t}}) = \min\{s, t\}$.

Norin et al. [22] studied the clustered chromatic number of minor-closed classes. They showed that for each $k \geq 2$, there is a graph H with treedepth k and connected treedepth k such that $\chi_{\star}(\mathcal{M}_H) \geq 2k - 2$. It is easily seen that the corresponding graphs have bounded pathwidth (at most $2k - 3$ to be precise). Thus the upper bound on $\chi_{\star}(\mathcal{G})$ in Theorem 2 is best possible.

Norin et al. [22] conjectured the following converse upper bound (analogous to Conjecture 3):

Conjecture 4 ([22]). *For every graph H ,*

$$\chi_{\star}(\mathcal{M}_H) \leq 2\overline{\text{td}}(H) - 2.$$

While Conjectures 3 and 4 remain open, Norin et al. [22] showed in the following theorem that $\chi_{\Delta}(\mathcal{M}_H)$ and $\chi_{\star}(\mathcal{M}_H)$ are controlled by the treedepth of H :

Theorem 5 ([22]). *For every graph H , $\chi_{\star}(\mathcal{M}_H)$ is tied to the (connected) treedepth of H . In particular,*

$$\overline{\text{td}}(H) - 1 \leq \chi_{\star}(\mathcal{M}_H) \leq 2^{\overline{\text{td}}(H)+1} - 4.$$

Theorem 1 gives a much more precise bound than Theorem 5 under the extra assumption of bounded treedepth.

Our third main result concerns fractional colourings. For real $t \geq 1$, a graph G is *fractionally t -colourable with clustering c* if there exist $Y_1, Y_2, \dots, Y_s \subseteq V(G)$ and $\alpha_1, \dots, \alpha_s \in [0, 1]$ such that¹:

- Every component of $G[Y_i]$ has at most c vertices,
- $\sum_{i=1}^s \alpha_i \leq t$,
- $\sum_{i:v \in Y_i} \alpha_i \geq 1$ for every $v \in V(G)$.

The *fractional clustered chromatic number* $\chi_{\star}^f(\mathcal{G})$ of a graph class \mathcal{G} is the infimum of $t > 0$ such that there exists $c = c(t, \mathcal{G})$ such that every $G \in \mathcal{G}$ is fractionally t -colourable with clustering c . *Fractional defective chromatic number* $\chi_{\Delta}^f(\mathcal{G})$ is defined in exactly the same way, except the condition on the component size is replaced by “the maximum degree of $G[Y_i]$ is at most d ”. The following theorem determines the fractional clustered chromatic number and fractional defective chromatic number of any proper minor-closed class.

Theorem 6. *For every proper minor-closed class \mathcal{G} ,*

$$\chi_{\Delta}^f(\mathcal{G}) = \chi_{\star}^f(\mathcal{G}) = \text{tcn}(\mathcal{G}) - 1.$$

This result is proved in Section 4.

We now give an interesting example of Theorem 6.

Corollary 7. *For every surface Σ , if \mathcal{G}_{Σ} is the class of graphs embeddable in Σ , then*

$$\chi_{\Delta}^f(\mathcal{G}_{\Sigma}) = \chi_{\star}^f(\mathcal{G}_{\Sigma}) = 3.$$

Proof. Note that $C\langle 3, k \rangle$ is planar for all k . Thus $\text{tcn}(\mathcal{G}_{\Sigma}) \geq 4$. Say Σ has Euler genus g . It follows from Euler’s formula that $K_{3,2g+3} \notin \mathcal{G}_{\Sigma}$. Since $K_{3,2g+3} \subseteq C\langle 4, 2g+3 \rangle$, we have $C\langle 4, 2g+3 \rangle \notin \mathcal{G}_{\Sigma}$. Thus $\text{tcn}(\mathcal{G}_{\Sigma}) \leq 4$. The result follows from Theorem 6. \square

In contrast to Corollary 7, Dvořák and Norin [7] proved that $\chi_{\star}(\mathcal{G}_{\Sigma}) = 4$. Note that Archdeacon [2] proved that $\chi_{\Delta}(\mathcal{G}_{\Sigma}) = 3$; see [5] for an improved bound on the defect.

¹ If $c = 1$, then this corresponds to a (proper) fractional t -colouring, and if the α_i are integral, then this yields a t -colouring with clustering c .

2 Treedepth

Say G is a subgraph of the closure of some rooted tree T . For each vertex $v \in V(T)$, let T_v be the subtree of T rooted at v (consisting of v and all its descendants), and let $G[T_v]$ be the subgraph of G induced by $V(T_v)$.

The *weak closure* of a rooted tree T is the graph G with vertex set $V(T)$, where two vertices $v, w \in V(T)$ are adjacent in G whenever v is a leaf of T and w is an ancestor of v in T . As illustrated in Figure 2, let $W\langle h, k \rangle$ be the weak closure of the complete k -ary tree of height h .




Figure 2: The weak closure $W\langle 4, 2 \rangle$.

Note that $W\langle h, k \rangle$ is a proper subgraph of $C\langle h, k \rangle$ for $h \geq 3$. On the other hand, Norin et al. [22] showed that $W\langle h, k \rangle$ contains $C\langle h, k - 1 \rangle$ as a minor for all $h, k \geq 2$. Therefore Theorem 1 is an immediate consequence of the following lemma.

Lemma 8. *For all $d, k \in \mathbb{N}$ there exists $c = c(d, k) \in \mathbb{N}$ such that for every $h \in \mathbb{N}$ and for every graph G with treedepth at most d , either G contains a $W\langle h, k \rangle$ -minor or G is $(h - 1)$ -colourable with clustering c .*

Proof. Throughout this proof, d is fixed, and we make no attempt to optimise c .

We may assume that G is connected. So G is a subgraph of the closure of some rooted tree of depth at most d . Choose a tree T of depth at most d rooted at some vertex r , such that G is a subgraph of the closure of T , and subject to this, $\sum_{v \in V(T)} \text{dist}_T(v, r)$ is minimal. Suppose that $G[T_v]$ is disconnected for some vertex v in T . Choose such a vertex v at maximum distance from r . Since G is connected, $v \neq r$. By the choice of v , for each child w of v , the subgraph $G[T_w]$ is connected. Thus, for some child w of v , there is no edge in G joining v and $G[T_w]$. Let u be the parent of v . Let T' be obtained from T by deleting the edge vw and adding the edge uw , so that w is a child of u in T' . Note that G is a subgraph of the closure of T' (since v has no neighbour in $G[T_w]$). Moreover, $\text{dist}_{T'}(x, r) = \text{dist}_T(x, r) - 1$ for every

vertex $x \in V(T_w)$, and $\text{dist}_{T'}(y, r) = \text{dist}_T(y, r)$ for every vertex $y \in V(T) \setminus V(T_w)$. Hence $\sum_{v \in V(T')} \text{dist}_{T'}(v, r) < \sum_{v \in V(T)} \text{dist}_T(v, r)$, which contradicts our choice of T . Therefore $G[T_v]$ is connected for every vertex v of T .

Consider each vertex $v \in V(T)$. Define the *level* $\ell(v) := \text{dist}_T(r, v) \in [0, d-1]$. Let T_v^+ be the subtree of T consisting of T_v plus the vr -path in T , and let $G[T_v^+]$ be the subgraph of G induced by $V(T_v^+)$. For a subtree X of T rooted at vertex v , define the *level* $\ell(X) := \ell(v)$.

A *ranked graph* (for fixed d) is a triple (H, L, \prec) where:

- H is a graph,
- $L : V(H) \rightarrow [0, d-1]$ is a function,
- \prec is a partial order on $V(H)$ such that $L(v) < L(w)$ whenever $v \prec w$.

Up to isomorphism, the number of ranked graphs on n vertices is at most $2^{\binom{n}{2}} d^n 3^{\binom{n}{2}}$. For a vertex v of T , a ranked graph (H, L, \prec) is said to be *contained in* $G[T_v^+]$ if there is an isomorphism ϕ from H to some subgraph of $G[T_v^+]$ such that:

- (A) for each vertex $v \in V(H)$ we have $L(v) = \ell(\phi(v))$, and
- (B) for all distinct vertices $v, w \in V(H)$ we have that $v \prec w$ if and only if $\phi(v)$ is an ancestor of $\phi(w)$ in T .

If (H, L, \prec) is a ranked graph and $i \in [0, d-1]$, then define the *i -splice* of (H, L, \prec) to be the ranked graph (H', L', \prec') obtained from (H, L, \prec) by taking k copies of the subgraph at levels greater than i . More formally, let

$$\begin{aligned} V(H') &:= \{(v, 0) : v \in V(H), L(v) \in [0, i]\} \cup \\ &\quad \{(v, j) : v \in V(H), L(v) \in [i+1, d], j \in [1, k]\}. \\ E(H') &:= \{(v, 0)(w, 0) : vw \in E(H), L(v) \in [0, i], L(w) \in [0, i]\} \cup \\ &\quad \{(v, 0)(w, j) : vw \in E(H), L(v) \in [0, i], L(w) \in [i+1, d], j \in [1, k]\} \cup \\ &\quad \{(v, j)(w, j) : vw \in E(H), L(v) \in [i+1, d], L(w) \in [i+1, d], j \in [1, k]\}. \end{aligned}$$

Define $L'((v, j)) := L(v)$ for every vertex $(v, j) \in V(H')$. Now define the following partial order \prec' on $V(H')$:

- If $v \prec w$ and $L(v), L(w) \in [0, i]$, then $(v, 0) \prec' (w, 0)$.
- If $v \prec w$ and $L(v) \in [0, i]$ and $L(w) \in [i+1, d]$, then $(v, 0) \prec' (w, j)$ for all $j \in [1, k]$.
- If $v \prec w$ and $L(v), L(w) \in [i+1, d]$, then $(v, j) \prec' (w, j)$ for all $j \in [1, k]$.

Note that if $(v, a) \prec' (w, b)$, then $a \leq b$ and $v \prec w$ (implying $L(v) < L(w)$). It follows that \prec' is a partial order on $V(H')$ such that $L'((v, a)) < L'((w, b))$ whenever $(v, a) \prec' (w, b)$. Thus (H', L', \prec') is a ranked graph.

For $\ell \in [0, d-1]$, let

$$N_\ell := (d+1)(h-1)(k+1)^{d-1-\ell}.$$

For each vertex v of T , define the *profile* of v to be the set of all ranked graphs (H, L, \prec) contained in $G[T_v^+]$ such that $|V(H)| \leq N_{\ell(v)}$. Note that if v is a descendant of u , then the profile of v is a subset of the profile of u . For $\ell \in [0, d-1]$, if $N = N_\ell$ then let

$$M_\ell := 2^{2^{\binom{N}{2}}} d^N 3^{\binom{N}{2}}.$$

Then there are at most M_ℓ possible profiles of a vertex at level ℓ .

We now partition $V(T)$ into subtrees. Each subtree is called a *group*. (At the end of the proof, vertices in a single group will be assigned the same colour.) We assign vertices to groups in non-increasing order of their distance from the root. Initialise this process by placing each leaf v of T into a singleton group. We now show how to determine the group of a non-leaf vertex. Let v be a vertex not assigned to a group at maximum distance from r . So each child of v is assigned to a group. Let Y_v be the set of children y of v , such that the number of children of v that have the same profile as y is in the range $[1, k-1]$. If $Y_v = \emptyset$ start a new singleton group $\{v\}$. If $Y_v \neq \emptyset$ then merge all the groups rooted at vertices in Y_v into one group including v . This defines our partition of $V(T)$ into groups. Each group X is *rooted* at the vertex in X closest to r in T . A group Y is *above* a distinct group X if the root of Y is on the path in T from the root of X to r .

The next claim is the key to the remainder of the proof.

Claim 1. *Let $uv \in E(T)$ where u is the parent of v , and u is in a different group to v . Then for every ranked graph (H, L, \prec) in the profile of v , the $\ell(u)$ -splice of (H, L, \prec) is in the profile of u .*

Proof. Since (H, L, \prec) is in the profile of v , there is an isomorphism ϕ from H to some subgraph of $G[T_v^+]$ such that for each vertex $x \in V(H)$ we have $L(x) = \ell(\phi(x))$, and for all distinct vertices $x, y \in V(H)$ we have that $x \prec y$ if and only if $\phi(x)$ is an ancestor of $\phi(y)$ in T .

Since u and v are in different groups, there are k children y_1, \dots, y_k of u (one of which is v) such that the profiles of y_1, \dots, y_k are equal. Thus (H, L, \prec) is in the profile of each of y_1, \dots, y_k . That is, for each $j \in [1, k]$, there is an isomorphism ϕ_j from H to some subgraph of $G[T_{y_j}^+]$ such that for each vertex $x \in V(H)$ we have $L(x) = \ell(\phi_j(x))$, and for all distinct vertices $x, y \in V(H)$ we have that $x \prec y$ if and only if $\phi_j(x)$ is an ancestor of $\phi_j(y)$ in T .

Let (H', L', \prec') be the $\ell(u)$ -splice of (H, L, \prec) . We now define a function ϕ' from $V(H')$ to $V(G[T_u^+])$. For each vertex $(x, 0)$ of H' (thus with $x \in V(H)$ and $L(x) \in [0, \ell(u)]$), define $\phi'((x, 0)) := \phi(x)$. For every other vertex (x, j) of H' (thus with $x \in V(H)$ and $L(x) \in [\ell(u) + 1, d-1]$ and $j \in [1, k]$), define $\phi'((x, j)) := \phi_j(x)$.

We now show that ϕ' is an isomorphism from H' to a subgraph of $G[T_u^+]$. Consider an edge $(x, a)(y, b)$ of H' . Thus $xy \in E(H)$. It suffices to show that $\phi'((x, a))\phi'((y, b)) \in E(G[T_u^+])$. First suppose that $a = b = 0$. So $L(x) \in [0, \ell(u)]$ and $L(y) \in [0, \ell(u)]$.

Thus $\phi'((x, a)) = \phi(x)$ and $\phi'((y, b)) = \phi(y)$. Since ϕ is an isomorphism to a subgraph of $G[T_v^+]$, we have $\phi(x)\phi(y) \in E(G[T_v^+])$, which is a subgraph of $G[T_u^+]$. Hence $\phi'((x, a))\phi'((y, b)) \in E(G[T_u^+])$, as desired. Now suppose that $a = 0$ and $b \in [1, k]$. Thus $\phi'((x, a)) = \phi(x)$ and $\phi'((y, b)) = \phi_b(y)$. Moreover, both $\ell(\phi(x))$ and $\ell(\phi_b(x))$ equal $L(x) \in [0, \ell(u)]$. There is only vertex z in T_v^+ with $\ell(z)$ equal to a specific number in $[0, \ell(u)]$. Thus $\phi'((x, a)) = \phi(x) = \phi_b(x) (= z)$. Since ϕ_b is an isomorphism to a subgraph of $G[T_{y_b}^+]$, we have $\phi_b(x)\phi_b(y) \in E(G[T_{y_b}^+])$, which is a subgraph of $G[T_u^+]$. Hence $\phi'((x, a))\phi'((y, b)) \in E(G[T_u^+])$, as desired. Finally, suppose that $a = b \in [1, k]$. Thus $\phi'((x, a)) = \phi_a(x)$ and $\phi'((y, b)) = \phi_b(y) = \phi_a(y)$. Since ϕ_a is an isomorphism to a subgraph of $G[T_{y_a}^+]$, we have $\phi_a(x)\phi_a(y) \in E(G[T_{y_a}^+])$, which is a subgraph of $G[T_u^+]$. Hence $\phi'((x, a))\phi'((y, b)) \in E(G[T_u^+])$, as desired. This shows that ϕ' is an isomorphism from H' to a subgraph of $G[T_u^+]$.

We now verify property (A) for (H', L', \prec') . For each vertex $(x, 0)$ of H' (thus with $x \in V(H)$ and $L(x) \in [0, \ell(u)]$) we have $L'((x, 0)) = L(x) = \ell(\phi(x)) = \ell(\phi'((x, 0)))$, as desired. For every other vertex (x, j) of H' (thus with $x \in V(H)$ and $L(x) \in [\ell(u) + 1, d - 1]$ and $j \in [1, k]$) we have $L'((x, j)) = L(x) = \ell(\phi_j(x)) = \ell(\phi'((x, j)))$, as desired. Hence property (A) is satisfied for (H', L', \prec') .

We now verify property (B) for (H', L', \prec') . Consider distinct vertices $(x, a), (y, b) \in V(H')$. First suppose that $a = 0$ and $b = 0$. Then $(x, a) \prec' (y, b)$ if and only if $x \prec y$ if and only if $\phi(x)$ is an ancestor of $\phi(y)$ in T if and only if $\phi'((x, a))$ is an ancestor of $\phi'((y, b))$ in T , as desired. Now suppose that $a = 0$ and $b \in [1, k]$. Then $(x, a) \prec' (y, b)$ if and only if $x \prec y$ if and only if $\phi(x)$ is an ancestor of $\phi_b(y)$ in T if and only if $\phi'((x, a))$ is an ancestor of $\phi'((y, b))$ in T , as desired. Now suppose that $a = b \in [1, k]$. Then $(x, a) \prec' (y, b)$ if and only if $x \prec y$ if and only if $\phi_a(x)$ is an ancestor of $\phi_b(y)$ in T if and only if $\phi'((x, a))$ is an ancestor of $\phi'((y, b))$ in T , as desired. Finally, suppose that $a, b \in [1, k]$ and $a \neq b$. Then (x, a) and (y, b) are incomparable under \prec' , and $\phi'((x, a))$ and $\phi'((y, b))$ in T are unrelated in T , as desired. Hence property (B) is satisfied for (H', L', \prec') .

So ϕ' is an isomorphism from H' to a subgraph of $G[T_u^+]$ satisfying properties (A) and (B). Thus (H', L', \prec') is contained in $G[T_u^+]$, as desired. Since (H, L, \prec) is in the profile of v , we have $|V(H)| \leq (d + 1)(h - 1)(k + 1)^{h - \ell(v)}$. Since $|V(H')| \leq (k + 1)|V(H)|$ and $\ell(u) = \ell(v) - 1$, we have $|V(H')| \leq (d + 1)(h - 1)(k + 1)^{h + 1 - \ell(v)} = (d + 1)(h - 1)(k + 1)^{h - \ell(u)}$. Thus (H', L', \prec') is in the profile of u . \square

The proof now divides into two cases. If some group X_0 is adjacent in G to at least $h - 1$ other groups above X_0 , then we show that G contains $W\langle h, k \rangle$ as a minor. Otherwise, every group X is adjacent in G to at most $h - 2$ other groups above X , in which case we show that G is $(h - 1)$ -colourable with bounded clustering.

Finding the Minor

Suppose that some group X_0 is adjacent in G to at least $h - 1$ other groups X_1, \dots, X_{h-1} above X_0 . We now show that G contains $W\langle h, k \rangle$ as a minor; refer to Figure 3.

For $i \in [0, h - 1]$, let v_i be the root of X_i . For $i \in [1, h - 1]$, let w_i be a vertex in X_i adjacent to some vertex z_i in X_0 ; since G is a subgraph of the closure of T , w_i and thus v_i are on the $v_0 r$ -path in T . For $i \in [0, h - 2]$, let u_i be the parent of v_i in T (which exists since $v_{h-2} \neq r$). So u_i is not in X_i (but may be in X_{i+1}). Note that $v_0, u_0, w_1, v_1, u_1, \dots, w_{h-2}, v_{h-2}, u_{h-2}, w_{h-1}, v_{h-1}$ appear in this order on the $v_0 r$ -path in T , where v_0, v_1, \dots, v_{h-1} are distinct (since they are in distinct groups).




Figure 3: Construction of a $W\langle 4, k \rangle$ minor (where u_i might be in X_{i+1}).

Let P_j be the $z_j r$ -path in T for $j \in [1, h - 1]$. Let H_0 be the graph with $V(H_0) := V(P_1 \cup \dots \cup P_{h-1})$ and $E(H_0) := \{z_j w_j : j \in [1, h - 1]\}$. Define the function $L_0 : V(H_0) \rightarrow [0, d - 1]$ by $L_0(x) := \ell(x)$ for each $x \in V(H_0)$. Define the partial order \prec_0 on $V(H_0)$, where $x \prec_0 y$ if and only if x is ancestor of y in T . Thus (H_0, L_0, \prec_0) is a ranked graph. By construction, (H_0, L_0, \prec_0) is contained in $G[T_{v_0}^+]$. Since H_0 has less than $(d + 1)(h - 1)$ vertices, H_0 is in the profile of v_0 . For $i = 0, 1, \dots, h - 2$, let $(H_{i+1}, L_{i+1}, \prec_{i+1})$ be the $\ell(u_i)$ -splice of (H_i, L_i, \prec_i) .

By induction, using Claim 1 at each step and since $G[T_{u_i}^+] \subseteq G[T_{v_{i+1}}^+]$, we conclude that for each $i \in [0, h - 1]$, the ranked graph (H_i, L_i, \prec_i) is in the profile of v_i . In particular, $(H_{h-1}, L_{h-1}, \prec_{h-1})$ is in the profile of v_{h-1} , and H_{h-1} is isomorphic to a subgraph of

G . Note that each vertex of H_{h-1} is of the form $((\dots(x, d_1), d_2), \dots), d_{h-1})$ for some $x \in V(H_0)$ and $d_1, \dots, d_{h-1} \in [0, k]$. For brevity, call such a vertex $x\langle d_1, \dots, d_{h-1} \rangle$. Note that if $x = w_j$ for some $j \in [1, h-1]$, then $d_1 = \dots = d_j = 0$ (since w_j is above u_i whenever $i < j$, and $(H_{i+1}, L_{i+1}, \prec_{i+1})$ is the $\ell(u_i)$ -splice of (H_i, L_i, \prec_i)).

For $x \in V(H_0)$, let Λ_x be the set of vertices $x\langle d_1, \dots, d_{h-1} \rangle$ in H_{h-1} . By construction, no two vertices in Λ_x are comparable under \prec_{h-1} . Therefore, by property (B), $V(T_a) \cap V(T_b) = \emptyset$ for all distinct $a, b \in \Lambda_x$. In particular, $V(T_a) \cap V(T_b) = \emptyset$ for all distinct $a, b \in \Lambda_{v_0}$. As proved above, $G[T_a]$ is connected for each $a \in V(T)$. Let G' be the graph obtained from G by contracting $G[T_a]$ into a single vertex $\alpha\langle d_1, \dots, d_{h-1} \rangle$, for each $a = v_0\langle d_1, \dots, d_{h-1} \rangle \in \Lambda_{v_0}$. So G' is a minor of G .

Let U be the tree with vertex set $\{\langle d_1, \dots, d_{h-1} \rangle : \exists j \in [0, h-1] \ d_1 = \dots = d_j = 0 \text{ and } d_{j+1}, \dots, d_{h-1} \in [1, k]\}$, where the parent of $(0, \dots, 0, d_{j+1}, d_{j+2}, \dots, d_{h-1})$ is $(0, \dots, 0, d_{j+2}, \dots, d_{h-1})$. Then U is isomorphic to the complete k -tree of height h rooted at $\langle 0, \dots, 0 \rangle$. We now show that the weak closure of U is a subgraph of G' , where each vertex $\langle 0, \dots, 0, d_{j+1}, \dots, d_{h-1} \rangle$ of U with $j \in [1, h-1]$ is mapped to vertex $w_j\langle 0, \dots, 0, d_{j+1}, \dots, d_{h-1} \rangle$ of G' , and each other vertex $\langle d_1, \dots, d_{h-1} \rangle$ of U is mapped to $\alpha\langle d_1, \dots, d_{h-1} \rangle$ of G' . For all $d_1, \dots, d_{h-1} \in [1, k]$ and $j \in [1, h-1]$ the vertex $z_j\langle d_1, \dots, d_{h-1} \rangle$ of G is contracted into the vertex $\alpha\langle d_1, \dots, d_{h-1} \rangle$ of G' . By construction, $z_j\langle d_1, \dots, d_{h-1} \rangle$ is adjacent to $w_j\langle 0, \dots, 0, d_{j+1}, \dots, d_{h-1} \rangle$ in G . So $\alpha\langle d_1, \dots, d_{h-1} \rangle$ is adjacent to $w_j\langle 0, \dots, 0, d_{j+1}, \dots, d_{h-1} \rangle$ in G' . This implies that the weak closure of U (that is, $W\langle h, k \rangle$) is isomorphic to a subgraph of G' , and is therefore a minor of G .

Finding the Colouring

Now assume that every group X is adjacent in G to at most $h-2$ other groups above X . Then $(h-1)$ -colour the groups in order of distance from the root, such that every group X is assigned a colour different from the colours assigned to the neighbouring groups above X . Assign each vertex within a group the same colour as that assigned to the whole group. This defines an $(h-1)$ -colouring of G .

Consider the function $s : [0, d-1] \rightarrow \mathbb{N}$ recursively defined by

$$s(\ell) := \begin{cases} 1 & \text{if } \ell = d-1 \\ (k-1) \cdot M_{\ell+1} \cdot s(\ell+1) & \text{if } \ell \in [0, d-2]. \end{cases}$$

Then every group at level ℓ has at most $s(\ell)$ vertices. By construction, our $(h-1)$ -colouring of G has clustering $s(0)$, which is bounded by a function of d and k , as desired. \square

3 Pathwidth

The following lemma of independent interest is the key to proving Theorem 2. Note that Eppstein [10] independently discovered the same result (with a slightly weaker

bound on the path length). The decomposition method in the proof has been previously used, for example, by Dujmović, Joret, Kozik, and Wood [6, Lemma 17].

Lemma 9. *Every graph with pathwidth at most w has a vertex 2-colouring such that each monochromatic path has at most $(w + 3)^w$ vertices.*

Proof. We proceed by induction on $w \geq 1$. Every graph with pathwidth 1 is a caterpillar, and is thus properly 2-colourable. Now assume $w \geq 2$ and the result holds for graphs with pathwidth at most $w - 1$. Let G be a graph with pathwidth at most w . Let (B_1, \dots, B_n) be a path-decomposition of G with width at most w . Let $t_0, t_1, t_2, \dots, t_m$ be a maximal sequence such that $t_0 = 0$, $t_1 = 1$, and for each $i \geq 2$, t_i is the minimum integer such that $B_{t_i} \cap B_{t_{i-1}} = \emptyset$. For odd i , colour every vertex in B_{t_i} 'red'. For even i , colour every vertex in B_{t_i} 'blue'. Since $B_{t_i} \cap B_{t_{i-1}} = \emptyset$, no vertex is coloured twice. Let G' be the subgraph of G induced by the uncoloured vertices. By the choice of B_{t_i} , each bag B_j with $j \in [t_{i-1} + 1, t_i - 1]$ intersects $B_{t_{i-1}}$. Thus $(B_1 \cap V(G'), \dots, B_n \cap V(G'))$ is a path-decomposition of G' of width at most $w - 1$. By induction, G' has a vertex 2-colouring such that each monochromatic path has at most $(w + 3)^{w-1}$ vertices. Since $B_{t_i} \cup B_{t_{i+2}}$ separates $B_{t_{i+1}} \cup \dots \cup B_{t_{i+2}-1}$ from the rest of G , each monochromatic component of G is contained in $B_{t_{i+1}} \cup \dots \cup B_{t_{i+2}-1}$ for some $i \in [0, n - 2]$. Consider a monochromatic path P in $G[B_{t_{i+1}} \cup \dots \cup B_{t_{i+2}-1}]$. Then P has at most $w + 1$ vertices in $B_{t_{i+1}}$. Note that $P - B_{t_{i+1}}$ is contained in G' . Thus P consists of up to $w + 2$ monochromatic subpaths in G' plus $w + 1$ vertices in $B_{t_{i+1}}$. Hence P has at most $(w + 2)(w + 3)^{w-1} + (w + 1) < (w + 3)^w$ vertices. \square

Nešetřil and Ossona de Mendez [21] showed that if a graph G contains no path on k vertices, then $\text{td}(G) < k$ (since G is a subgraph of the closure of a DFS spanning tree with height at most k). Thus Lemma 9 implies:

Corollary 10. *Every graph with pathwidth at most w has a vertex 2-colouring such that each monochromatic component has treedepth at most $(w + 3)^w$.*

Proof of Theorem 2. Let \mathcal{G} be a minor-closed class of graphs, each with pathwidth at most w . Let h be the minimum integer such that $C\langle h, k \rangle \notin \mathcal{G}$ for some $k \in \mathbb{N}$. Consider $G \in \mathcal{G}$. By Corollary 10, G has a vertex 2-colouring such that each monochromatic component H of G has treedepth at most $(w + 3)^w$. Since $C\langle h, k \rangle$ is not a minor of H , by Lemma 8, H is $(h - 1)$ -colourable with clustering $c((w + 3)^w, k)$. Taking a product colouring, G is $(2h - 2)$ -colourable with clustering $c((w + 3)^w, k)$. Hence $\chi_{\Delta}(\mathcal{G}) \leq \chi_{\star}(\mathcal{G}) \leq 2h - 2$. \square

Note that Lemma 9 cannot be extended to the setting of bounded tree-width graphs: Esperet and Joret (see [16, Theorem 4.1]) proved that for all positive integers w and d there exists a graph G with tree-width at most w such that for every w -colouring of G there exists a monochromatic component of G with diameter greater than d (and thus with a monochromatic path on more than d vertices, and thus with treedepth at least $\log_2 d$).

4 Fractional Colouring

This section proves Theorem 6. The starting point is the following key result of Dvořák and Sereni [8].²

Theorem 11 ([8]). *For every proper minor-closed class \mathcal{G} and every $\delta > 0$ there exists $d \in \mathbb{N}$ satisfying the following. For every $G \in \mathcal{G}$ there exist $s \in \mathbb{N}$ and $X_1, X_2, \dots, X_s \subseteq V(G)$ such that:*

- $\text{td}(G[X_i]) \leq d$, and
- every $v \in V(G)$ belongs to at least $(1 - \delta)s$ of these sets.

We now prove a lower bound on the fractional defective chromatic number of the closure of complete trees of given height.

Lemma 12. *Let $\mathcal{C}_h := \{C\langle h, k \rangle\}_{k \in \mathbb{N}}$. Then $\chi_{\Delta}^f(\mathcal{C}_h) \geq h$.*

Proof. We show by induction on h that if $C\langle h, k \rangle$ is fractionally t -colourable with defect d , then $t \geq h - (h - 1)d/k$. This clearly implies the lemma. The base case $h = 1$ is trivial.

For the induction step, suppose that $G := C\langle h, k \rangle$ is fractionally t -colourable with defect d . Thus there exist $Y_1, Y_2, \dots, Y_s \subseteq V(G)$ and $\alpha_1, \dots, \alpha_s \in [0, 1]$ such that:

- every component of $G[Y_i]$ has maximum degree at most d ,
- $\sum_{i=1}^s \alpha_i \leq t$, and
- $\sum_{i: v \in Y_i} \alpha_i \geq 1$ for every $v \in V(G)$.

Let r be the vertex of G corresponding to the root of the complete k -ary tree and let H_1, \dots, H_k be the components of $G - r$. Then each H_i is isomorphic to $C\langle h - 1, k \rangle$. Let $J_0 := \{j : r \in Y_j\}$, and let $J_i := \{j : Y_j \cap V(H_i) \neq \emptyset\}$ for $i \in [1, k]$. Denote $\sum_{j \in J_i} \alpha_j$ by $\alpha(J_i)$ for brevity. Thus $\alpha(J_0) \geq 1$. For $i \in [1, k]$, the subgraph H_i is $\alpha(J_i)$ -colourable with defect d , and thus $\alpha(J_i) \geq h - 1 - (h - 2)d/k$ by the induction hypothesis. Thus

$$(k - d)\alpha(J_0) + \sum_{i=1}^k \alpha(J_i) \geq (k - d) + k(h - 1) - (h - 2)d = kh - (h - 1)d.$$

If $j \in J_0$ then Y_j intersects at most d of H_1, \dots, H_k (since $G[Y_j]$ has maximum degree at most d). Thus every α_j appears with coefficient at most k in the left side of the above inequality, implying

$$(k - d)\alpha(J_0) + \sum_{i=1}^k \alpha(J_i) \leq k \sum_{i=1}^s \alpha_i \leq kt.$$

² Dvořák and Sereni [8] expressed their result in the terms of “treedepth fragility”. The sentence “proper minor-closed classes are fractionally treedepth-fragile” after Theorem 31 in [8] is equivalent to Theorem 11. Informally speaking, Theorem 11 shows that the fractional “treedepth” chromatic number of every minor-closed class equals 1.

Combining the above inequalities yields the claimed bound on t . \square

Proof of Theorem 6. By Lemma 12,

$$\chi_{\star}^f(\mathcal{G}) \geq \chi_{\Delta}^f(\mathcal{G}) \geq \text{tcn}(\mathcal{G}) - 1.$$

It remains to show that $\chi_{\star}^f(\mathcal{G}) \leq \text{tcn}(\mathcal{G}) - 1$. Equivalently, we need to show that for all $h, k \in \mathbb{N}$ and $\varepsilon > 0$, if $C\langle h, k \rangle \notin \mathcal{G}$ then there exists c such that every graph in \mathcal{G} is fractionally $(h - 1 + \varepsilon)$ -colourable with clustering c . This is trivial for $h = 1$, and so we assume $h \geq 2$.

Let $d \in \mathbb{N}$ satisfy the conclusion of Theorem 11 for the class \mathcal{G} and $\delta = 1 - \frac{1}{1+\varepsilon/(h-1)}$. Choose $c = c(d, k + 1)$ to satisfy the conclusion of Lemma 8. We show that c is as desired.

Consider $G \in \mathcal{G}$. By the choice of d there exists $s \in \mathbb{N}$ and $X_1, X_2, \dots, X_s \subseteq V(G)$ such that:

- $\text{td}(G[X_i]) \leq d$, and
- every $v \in V(G)$ belongs to at least $(1 - \delta)s$ of these sets.

Since $C\langle h, k \rangle \notin \mathcal{G}$, we have $W\langle h, k + 1 \rangle \notin \mathcal{G}$, and by the choice of c , for each $i \in [1, s]$ there exists a partition $(Y_i^1, Y_i^2, \dots, Y_i^{h-1})$ of X_i such that every component of $G[Y_i^j]$ has at most c vertices. Every vertex of G belongs to at least $(1 - \delta)s$ sets Y_i^j where $i \in [1, s]$ and $j \in [1, h - 1]$. Considering these sets with equal coefficients $\alpha_i^j := \frac{1}{(1-\delta)s}$, we conclude that G is fractionally $\frac{h-1}{1-\delta}$ -colourable with clustering c , as desired (since $\frac{h-1}{1-\delta} = h - 1 + \varepsilon$). \square

Acknowledgement

This work was partially completed while SN was visiting Monash University supported by a Robert Bartnik Visiting Fellowship. SN thanks the School of Mathematics at Monash University for its hospitality.

References

- [1] NOGA ALON, GUOLI DING, BOGDAN OPOROWSKI, AND DIRK VERTIGAN. [Partitioning into graphs with only small components](#). *J. Combin. Theory Ser. B*, 87(2):231–243, 2003.
- [2] DAN ARCHDEACON. [A note on defective colorings of graphs in surfaces](#). *J. Graph Theory*, 11(4):517–519, 1987.
- [3] NICOLAS BROUTIN AND ROSS J. KANG. [Bounded monochromatic components for random graphs](#). *J. Comb.*, 9(3):411–446, 2018. arXiv:1407.3555.
- [4] ILKYOO CHOI AND LOUIS ESPERET. [Improper coloring of graphs on surfaces](#). *J. Graph Theory*, 91(1):16–34, 2019.

- [5] LENORE COWEN, WAYNE GODDARD, AND C. ESTHER JESURUM. [Defective coloring revisited](#). *J. Graph Theory*, 24(3):205–219, 1997.
- [6] VIDA DUJMOVIĆ, GWENAËL JORET, JAKUB KOZIK, AND DAVID R. WOOD. [Nonrepetitive colouring via entropy compression](#). *Combinatorica*, 36(6):661–686, 2016.
- [7] ZDENĚK DVOŘÁK AND SERGEY NORIN. [Islands in minor-closed classes. I. Bounded treewidth and separators](#). 2017, arXiv:1710.02727.
- [8] ZDENĚK DVOŘÁK AND JEAN-SÉBASTIEN SERENI. [On fractional fragility rates of graph classes](#). *Electronic J. Combinatorics*, 27:P4.9, 2020.
- [9] KATHERINE EDWARDS, DONG YEAP KANG, JAEHOON KIM, SANG-IL OUM, AND PAUL SEYMOUR. [A relative of Hadwiger’s conjecture](#). *SIAM J. Discrete Math.*, 29(4):2385–2388, 2015.
- [10] DAVID EPPSTEIN. [Pathbreaking for intervals](#). In *11011110*. 2020.
- [11] LOUIS ESPERET AND GWENAËL JORET. [Colouring planar graphs with three colours and no large monochromatic components](#). *Combinatorics, Probability & Computing*, 23(4):551–570, 2014.
- [12] PENNY HAXELL, TIBOR SZABÓ, AND GÁBOR TARDOS. [Bounded size components—partitions and transversals](#). *J. Combin. Theory Ser. B*, 88(2):281–297, 2003.
- [13] DONG YEAP KANG AND SANG-IL OUM. [Improper coloring of graphs with no odd clique minor](#). *Combin. Probab. Comput.*, 28(5):740–754, 2019.
- [14] KEN-ICHI KAWARABAYASHI. [A weakening of the odd Hadwiger’s conjecture](#). *Combin. Probab. Comput.*, 17(6):815–821, 2008.
- [15] KEN-ICHI KAWARABAYASHI AND BOJAN MOHAR. [A relaxed Hadwiger’s conjecture for list colorings](#). *J. Combin. Theory Ser. B*, 97(4):647–651, 2007.
- [16] CHUN-HUNG LIU AND SANG-IL OUM. [Partitioning \$H\$ -minor free graphs into three subgraphs with no large components](#). *J. Combin. Theory Ser. B*, 128:114–133, 2018.
- [17] CHUN-HUNG LIU AND DAVID R. WOOD. [Clustered coloring of graphs excluding a subgraph and a minor](#). 2019, arXiv:1905.09495.
- [18] CHUN-HUNG LIU AND DAVID R. WOOD. [Clustered graph coloring and layered treewidth](#). 2019, arXiv:1905.08969.
- [19] CHUN-HUNG LIU AND DAVID R. WOOD. [Clustered variants of Hajós’ conjecture](#). 2019, arXiv:1908.05597.
- [20] BOJAN MOHAR, BRUCE REED, AND DAVID R. WOOD. [Colourings with bounded monochromatic components in graphs of given circumference](#). *Australas. J. Combin.*, 69(2):236–242, 2017.
- [21] JAROSLAV NEŠETŘIL AND PATRICE OSSONA DE MENDEZ. [Sparsity](#), vol. 28 of *Algorithms and Combinatorics*. Springer, 2012.
- [22] SERGEY NORIN, ALEX SCOTT, PAUL SEYMOUR, AND DAVID R. WOOD. [Clustered colouring in minor-closed classes](#). *Combinatorica*, 39(6):1387–1412, 2019.
- [23] PATRICE OSSONA DE MENDEZ, SANG-IL OUM, AND DAVID R. WOOD. [Defective colouring of graphs excluding a subgraph or minor](#). *Combinatorica*, 39(2):377–410, 2019.
- [24] JAN VAN DEN HEUVEL AND DAVID R. WOOD. [Improper colourings inspired by Hadwiger’s conjecture](#). *J. London Math. Soc.*, 98:129–148, 2018.

- [25] DAVID R. WOOD. [Defective and clustered graph colouring](#). *Electron. J. Combin.*, DS23, 2018. Version 1.