

Elliptic Curves. Sheet 4.

1. Find the torsion group over $\mathbb{Q}$ for each of:

(a). $Y^2 = X^3 + 1$. (b). $Y^2 = X(X - 1)(X - 2)$. (c). $Y^2 = X^3 + 1/3^6$.

2. Let, as usual, $\mathcal{C} : Y^2 = X(X^2 + aX + b)$ and $\mathcal{D} : Y^2 = X(X^2 + a_1X + b_1)$, where $a, b \in \mathbb{Z}$, $a_1 = -2a$, $b_1 = a^2 - 4b$ and $b(a^2 - 4b) \neq 0$. Let $\mathcal{C}_{\text{oddtors}}(\mathbb{Q})$ denote the set of torsion elements of $\mathcal{C}(\mathbb{Q})$ which have odd order, and let $\mathcal{D}_{\text{oddtors}}(\mathbb{Q})$ denote the set of torsion elements of $\mathcal{D}(\mathbb{Q})$ which have odd order. Show that $\mathcal{C}_{\text{oddtors}}(\mathbb{Q})$ and $\mathcal{D}_{\text{oddtors}}(\mathbb{Q})$ are isomorphic.

3. Let $\mathcal{C}$ and $\mathcal{D}$ be as in question 2. Let the homomorphisms $\phi, \hat{\phi}$ be defined as usual by

$$\phi : \mathcal{C} \rightarrow \mathcal{D} : (x, y) \mapsto \left(\left(\frac{y}{x} \right)^2, y - \frac{by}{x^2} \right), \quad \hat{\phi} : \mathcal{D} \rightarrow \mathcal{C} : (u, v) \mapsto \left(\frac{1}{4} \left(\frac{v}{u} \right)^2, \frac{1}{8} \left(v - \frac{b_1 v}{u^2} \right) \right).$$

What are the preimages of $(0, 0)$ under $\hat{\phi}$? Show that $(0, 0) \in 2\mathcal{C}(\mathbb{Q})$ if and only if there exist $m, n \in \mathbb{Z}$ such that $b = m^2$ and $a + 2m = n^2$.

4. Find the ranks of the following elliptic curves.

(a). $Y^2 = X(X^2 + 2X + 3)$.

(b). $Y^2 = X(X^2 + 14X + 1)$.

5. Let $A, +$ be an Abelian group. Let $h : A \rightarrow \mathbb{R}_{\geq 0}$ satisfy:

(I) There exists a constant C , independent of P, Q , such that

$$|h(P + Q) + h(P - Q) - 2h(P) - 2h(Q)| \leq C, \text{ for all } P, Q \in A,$$

(II) For any $B \in \mathbb{R}$, the set $\{P \in A : h(P) \leq B\}$ is finite.

Show that h is a height function on A . Show also that there exists a constant C_3 , independent of P , such that $|h(3P) - 9h(P)| \leq C_3$, for all $P \in A$.

$[\mathbb{R}_{\geq 0}]$ denotes $\{x \in \mathbb{R} : x \geq 0\}$.

6. A four-letter word $L_1L_2L_3L_4$ has been divided into two pairs: L_1L_2 and L_3L_4 . Each of these pairs has been converted into an integer (of at most 4 digits) via the standard map: $A \mapsto 01, B \mapsto 02, \dots, Z \mapsto 26$. These integers have been encoded by taking each to the power of $d = 4085$, modulo $N = 10481$. The encoded message reads: **6012, 3236**. You may assume that N is the product of two primes. You should show, in your calculations, how you are only using numbers of length at most 9 digits.

(a) Find a proper factor of N (that is, a factor d of N satisfying $1 < d < N$) by applying Pollard's "p - 1" method, using base 2 and exponent 46.

(b) Factorise N by applying the Elliptic Curve Method, using the curve $\mathcal{E} : Y^2 = X^3 - X + 1$ and $3P$, where $P = (5, 11)$.

(c) Use the factorisation of N to decode the message (which is the name of the town famous for being the country music capital of New Zealand).

The following question is compulsory for students taking the MSc in MFoCS (Mathematics and the Foundations of Computer Science). For everyone else, it is optional.

7. Show that any elliptic curve over $\mathbb{Q}$ with a rational point of order 4 is birationally equivalent to: $Y^2 + XY + vY = X^3 + vX^2$, for some $v \in \mathbb{Q}$.