

Feynman periods - on graphs, integrals, polytopes and tropical physics


Erik Panzer

All Souls College (Oxford)


18th June 2019

Felix Klein Kolloquium, TU Kaiserslautern


Perturbative Quantum Field Theory




Perturbative Quantum Field Theory




Perturbative Quantum Field Theory




Perturbative Quantum Field Theory




Perturbative Quantum Field Theory




Perturbative Quantum Field Theory




Perturbative Quantum Field Theory




Perturbative Quantum Field Theory




Perturbative Quantum Field Theory




- Feynman graph $G \mapsto$ Feynman integral $\Phi(G, \{m_i^2, \vec{p}_i \cdot \vec{p}_j\})$
- compute more graphs $\sum_G \Phi(G) \Rightarrow$ higher precision



not spanning



not connected



has a loop

Definition

A **spanning tree** $T \subset G$ is a spanning, simply connected subgraph.

$$\text{ST} \left(\begin{array}{c} \text{graph with 4 vertices and 5 edges} \end{array} \right) = \left\{ \begin{array}{c} \text{graph with 4 vertices and 5 edges} \end{array}, \begin{array}{c} \text{graph with 4 vertices and 5 edges} \end{array}, \dots \right\}$$

Definition

The **graph polynomial** Ψ and **Feynman period** of G are

$$\Psi = \sum_{T \in \text{ST}(G)} \prod_{e \notin T} x_e \quad \text{and} \quad \mathcal{P}(G) = \left(\prod_{e \geq 1} \int_0^\infty dx_e \right) \frac{1}{\Psi^2|_{x_1=1}}$$

$$G = \text{circle with two vertices} \Rightarrow \Psi = x_1 + x_2 \quad \text{and} \quad \mathcal{P}(\text{circle with two vertices}) = \int_0^\infty \frac{dx_2}{(1+x_2)^2} = 1$$

Definition

The **graph polynomial** Ψ and **Feynman period** of G are

$$\Psi = \sum_{T \in \text{ST}(G)} \prod_{e \notin T} x_e \quad \text{and} \quad \mathcal{P}(G) = \left(\prod_{e \geq 1} \int_0^\infty dx_e \right) \frac{1}{\Psi^2|_{x_1=1}}$$

$$G = \text{circle with two vertices} \Rightarrow \Psi = x_1 + x_2 \quad \text{and} \quad \mathcal{P} \left(\text{circle with two vertices} \right) = \int_0^\infty \frac{dx_2}{(1+x_2)^2} = 1$$

Assumptions:

① logarithmic divergence: $\omega(G) := |E(G)| - 2 \cdot \ell(G) \stackrel{!}{=} 0$

$$\ell(G) = h_1(G) = |E(G)| - |V(G)| + 1 \quad (\text{loop number})$$


② no subdivergences: $\omega(\gamma) > 0$ for all $\emptyset \neq \gamma \subsetneq G$

All such periods contribute to the β -function of the field theory.

\Rightarrow *renormalization constants, running coupling, critical exponents*

- These are periods in the sense of Kontsevich and Zagier
 \Rightarrow interesting transcendental numbers, motivic Galois theory


Example

$$\mathcal{P} \left(\text{Diagram} \right) = \int_{\mathbb{R}_+^5} \frac{dx_2 dx_3 dx_4 dx_5 dx_6}{(x_1 x_2 x_3 + 15 \text{ more terms})^2} = 6\zeta(3) = 6 \sum_{n=1}^{\infty} \frac{1}{n^3}$$


- Sometimes expressible as **multiple zeta values**

$$\zeta(s_1, \dots, s_d) = \sum_{0 < n_1 < \dots < n_d} \frac{1}{n_1^{s_1} \dots n_d^{s_d}}$$

Example (Broadhurst & Schnetz)


$$\mathcal{P} \left(\text{Diagram} \right) = \frac{92943}{160} \zeta(11) + 896 \zeta(3) \left(\frac{27}{80} \zeta(3, 5) + \frac{45}{64} \zeta(3) \zeta(5) - \frac{261}{320} \zeta(8) \right) + \frac{3381}{20} (\zeta(3, 5, 3) - \zeta(3, 5) \zeta(3)) - \frac{1155}{4} \zeta(3)^2 \zeta(5)$$


- These integrals are very hard to compute (**even numerically**).


Only two infinite families of periods are known: wheels and zigzags




W_3



W_4




W_5




W_6

Theorem (Broadhurst 1985)


$$\mathcal{P}(W_n) = \binom{2n-2}{n-1} \zeta(2n-3)$$



ZZ_3



ZZ_4



ZZ_5



ZZ_6




Theorem (Brown & Schnetz 2012)

$$\mathcal{P}(ZZ_n) = 4 \frac{(2n-2)!}{n!(n-1)!} \left(1 - \frac{1 - (-1)^n}{2^{2n-3}} \right) \zeta(2n-3)$$



> 1000 more periods are known [Broadhurst, Schnetz, Panzer]

When is $\mathcal{P}(G_1) = \mathcal{P}(G_2)$?


1 Product:

$$\mathcal{P} \left(\text{Diagram 1} \right) = \mathcal{P} \left(\text{Diagram 2} \right) \cdot \mathcal{P} \left(\text{Diagram 3} \right)$$




Example

$$\mathcal{P} \left(\text{Diagram 4} \right) = \mathcal{P} \left(\text{Diagram 5} \right)^2 = (6\zeta(3))^2$$



2 Planar duality: $\mathcal{P}(G) = \mathcal{P}(G^{\text{dual}})$




- ③ Completion: If G is 4-regular and v, w are vertices G , then

$$\mathcal{P}(G \setminus v) = \mathcal{P}(G \setminus w)$$

Example

$$\mathcal{P}\left(\begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \end{array}\right) = \mathcal{P}\left(\begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \\ \quad \quad \quad \bullet \\ \quad \quad \quad \diagup \quad \diagdown \\ \quad \quad \quad \bullet \quad \bullet \\ \quad \quad \quad \diagdown \quad \diagup \\ \quad \quad \quad \bullet \end{array} \setminus v\right) = \mathcal{P}\left(\begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \\ \quad \quad \quad \bullet \\ \quad \quad \quad \diagup \quad \diagdown \\ \quad \quad \quad \bullet \quad \bullet \\ \quad \quad \quad \diagdown \quad \diagup \\ \quad \quad \quad \bullet \end{array} \setminus w\right) = \mathcal{P}\left(\begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \end{array}\right)$$

- ④ Twist:



[drawing by Crump]

Goal:

Construct simpler graph invariants with those symmetries.

$$c_2(G)(p) = \frac{1}{p^2} \left| \left\{ \vec{x} \in (\mathbb{Z}/p\mathbb{Z})^N : \Psi(\vec{x}) = 0 \right\} \right|$$

$P_{7,11}$

p	2	3	5	7	11	13	17	19	23
$c_2(p)$	1	0	1	-1	1	-1	1	-1	1
$\text{Perm}(p)$		0	1	1	1	11	5	0	22

Goal:

Construct simpler graph invariants with those symmetries.

$$c_2(G)(p) = \frac{1}{p^2} \left| \left\{ \vec{x} \in (\mathbb{Z}/p\mathbb{Z})^N : \Psi(\vec{x}) = 0 \right\} \right|$$

$P_{7,11}$

p	2	3	5	7	11	13	17	19	23
$c_2(p)$	1	0	1	-1	1	-1	1	-1	1
$\text{Perm}(p)$		0	1	1	1	11	5	0	22

	product	duality	completion	twist
--	---------	---------	------------	-------

c_2 [Schnetz]	yes	yes [Doryn]	for $p = 2$ [Yeats, Doryn]	open	few values, sees number theory
--------------------	-----	----------------	-------------------------------	------	-----------------------------------


permanent [Crump]	yes	yes	yes	yes	almost faithful
----------------------	-----	-----	-----	-----	-----------------

\mathcal{H}	yes	yes	yes	yes	faithful (conj.), sees magnitude
---------------	-----	-----	-----	-----	-------------------------------------

Hepp bound

$$\mathcal{H}(G) := \left(\prod_{e \in G} \int_0^\infty dx_e \right) \frac{1}{\Psi_{\max}^2 |x_1=1|} \quad \text{where} \quad \Psi_{\max} := \max_{T \in \text{ST}(G)} \prod_{e \notin T} x_e$$

Example


$$\mathcal{H} \left(\text{circle with two vertices} \right) = \int_0^\infty \frac{dx_1}{(\max\{x_1, 1\})^2} = \int_0^1 dx_1 + \int_1^\infty \frac{dx_1}{x_1^2} = 2$$

- $\mathcal{H}(G) > \mathcal{P}(G) > \mathcal{H}(G)/|\text{ST}(G)|^2$
- fulfils the four symmetries
- $\mathcal{H}(G) \in \mathbb{Q}_{>0}$
- can be computed very efficiently


Theorem

$$\mathcal{H}(G) = \sum_{\substack{\gamma_1 \subset \gamma_2 \subset \dots \subset \gamma_{\ell(G)} = G \\ \text{each } \gamma_i \text{ is 1PI}}} \frac{|\gamma_1| \cdot |\gamma_2 \setminus \gamma_1| \cdots |G \setminus \gamma_{\ell(G)-1}|}{\omega(\gamma_1) \cdots \omega(\gamma_{\ell(G)-1})}$$

γ_1	\subset	γ_2	summand	#	Σ
	\subset		$\frac{3 \cdot 2 \cdot 1}{1 \cdot 1} = 6$	12	72
	\subset		$\frac{4 \cdot 1 \cdot 1}{2 \cdot 1} = 2$	6	12

$\left. \vphantom{\begin{matrix} \text{summand} \\ \# \\ \Sigma \end{matrix}} \right\} \Rightarrow \mathcal{H} \left(\text{triangle} \right) = 84$

Hepp-Period correlation



Conjecture


$$\mathcal{H}(G_1) = \mathcal{H}(G_2) \quad \Leftrightarrow \quad \mathcal{P}(G_1) = \mathcal{P}(G_2)$$

Conjecture

$$\mathcal{H}(G_1) = \mathcal{H}(G_2) \quad \Leftrightarrow \quad \mathcal{P}(G_1) = \mathcal{P}(G_2)$$

For example, we find a pair of unknown 8 loop periods with:

- $\mathcal{H}(P_{8,30}) = \frac{1724488}{3} = \mathcal{H}(P_{8,36})$
- $\mathcal{P}(P_{8,30}) \approx 505.5 \approx \mathcal{P}(P_{8,36})$



The Hepp bound is the volume of a polytope

Spanning tree polytope:

$$\mathcal{N}_G := \text{conv} \left\{ \vec{T} - \vec{T}^c : T \text{ spanning tree of } G \right\} \subset \left\{ \vec{G} \cdot \vec{a} = 0 \right\} \subset \mathbb{R}^{E_G}$$

Facets of \mathcal{N}_G are indexed by certain subgraphs:

$$A_G := \{ \gamma \subset G : \gamma \text{ and } G/\gamma \text{ are 2-vertex connected} \}$$

Factorisation of the facets:

$$F_\gamma := \mathcal{N}_G \cap \{ \vec{\gamma} \cdot \vec{a} = \omega_\gamma \} \cong \mathcal{N}_\gamma \times \mathcal{N}_{G/\gamma}$$

Lemma

$$\mathcal{H}(G) = (E_G - 1)! \cdot \text{Vol}(\mathcal{N}_G^\circ)$$

$$\mathcal{N}_G^\circ = \bigcap_{T \text{ spanning tree of } G} \left\{ \vec{a} : \vec{a} \cdot (\vec{T} - \vec{T}^c) \leq 1 \right\}$$

Multivariate version & canonical form

Now consider arbitrary indices:

$$\mathcal{H}(G; \vec{\nu}) := \int \frac{\delta(1 - x_i) d^E \vec{x}}{\psi_{\max}^{D/2}} \prod_e x_e^{\nu_e - 1}$$

The dimension is fixed by $\omega(G) = \sum_e \nu_e - (D/2) \cdot \ell(G) \stackrel{!}{=} 0$.

Example

The flag formula generalizes to this case, e.g.

$$\mathcal{H} \left(\begin{array}{c} \text{Diagram: A triangle with vertices } v_1, v_2, v_3 \text{ and edges } (v_1, v_2) \text{ labeled } 1, (v_1, v_3) \text{ labeled } 2, (v_2, v_3) \text{ labeled } 3. \text{ There is an additional vertex } v_4 \text{ connected to } v_3 \text{ by an edge labeled } 4. \end{array} ; \vec{\nu} \right) = \frac{1}{\nu_1 \nu_2 \nu_3 \nu_4} \times \left\{ \frac{(\nu_1 + \nu_2 + \nu_3) \nu_4}{\nu_1 + \nu_2 + \nu_3 - D/2} + \frac{(\nu_1 + \nu_2 + \nu_4) \nu_3}{\nu_1 + \nu_2 + \nu_4 - D/2} + \frac{(\nu_3 + \nu_4)(\nu_1 + \nu_2)}{\nu_3 + \nu_4 - D/2} \right\}$$

Consider the Hepp bound $\mathcal{H}(G; \vec{\nu})$:

- it is a rational function in $\vec{\nu}$
- it has simple poles
- at hyperplanes $\omega(\gamma) = 0$ for 1PI subgraphs γ

Lemma (factorization of residues)

$$\text{Res}_{\omega(\gamma)=0} \mathcal{H}(G; \vec{\nu}) = \mathcal{H}(\gamma; \vec{\nu}_\gamma) \Big|_{\omega(\gamma)=0} \cdot \mathcal{H}(G/\gamma; \vec{\nu}_{G/\gamma}) \Big|_{\omega(G/\gamma)=0}$$

Example

$$\text{Res}_{\nu_e=0} \mathcal{H}(G; \vec{\nu}) = \mathcal{H}(G/e; \vec{\nu}_{G/e})$$

- it is the volume of a polytope:

$$\mathcal{H}(G; \vec{\nu}) = (E - 1)! \cdot \text{Vol} \left(\left(\mathcal{N}_G + (\vec{\nu} - \vec{1}) \right)^\circ \right)$$

Summary

- There is a rational version of Feynman periods.
- It captures identities and gives numeric estimates.
- Relates to a polytope and a function with factorizing residues.
- Generalizes to matroids.

Some questions

- add kinematics
- study divergences
- Can this be turned into an approximation scheme?

Thanks

Thank you for your attention!