

Induced subgraphs of graphs with large chromatic number.
XII. Distant stars

Maria Chudnovsky¹
Princeton University, Princeton, NJ 08544

Alex Scott²
Oxford University, Oxford, UK

Paul Seymour³
Princeton University, Princeton, NJ 08544

¹Supported by NSF grant DMS-1550991 and US Army Research Office Grant W911NF-16-1-0404.

²Supported by a Leverhulme Trust Research Fellowship.

³Supported by ONR grant N00014-14-1-0084 and NSF grant DMS-1265563.

Abstract

The Gyárfás-Sumner conjecture asserts that if H is a tree then every graph with bounded clique number and very large chromatic number contains H as an induced subgraph. This is still open, although it has been proved for a few simple families of trees, including trees of radius two, some special trees of radius three, and subdivided stars. These trees all have the property that their vertices of degree more than two are clustered quite closely together. In this paper, we prove the conjecture for two families of trees which do not have this restriction. As special cases, these families contain all double-ended brooms and two-legged caterpillars.

1 Introduction

All graphs in this paper are finite and simple. If G is a graph, then $\chi(G)$ denotes its chromatic number, and $\omega(G)$ denotes its clique number, that is, the cardinality of the largest clique of G .

Let H be a graph. When is there a function f such that $\chi(G) \leq f(\omega(G))$ for every graph G not containing H as an induced subgraph? Let us call such a graph H χ -*bounding*. Every χ -bounding graph H is a forest, because we could take G to have large girth and large chromatic number, and every such graph G should contain H . The Gyárfás-Sumner conjecture [2, 9] asserts that the converse holds:

1.1 Conjecture: *Every forest is χ -bounding.*

It is easy to see that a forest is χ -bounding if and only if all its components are χ -bounding, so the question reduces to trees. Despite considerable attention, there are still only a few families of trees for which the Gyárfás-Sumner conjecture has been proved. The only trees that have been shown to be χ -bounding so far are:

- trees of radius at most two (Gyárfás, Szemerédi and Tuza [4] in the triangle-free case; Kierstead and Penrice [5] in the general case);
- trees that can be obtained from a tree of radius at most two by subdividing once *every* edge incident with the root (Kierstead and Zhu [6]); and
- subdivisions of stars (this follows from the “topological” version of the Gyárfás-Sumner conjecture proved in [7]: for every tree T there is a function f such that $\chi(G) \leq f(\omega(G))$ for every graph G containing no subdivision of T as an induced subgraph. In fact, it is enough to exclude the finite family of subdivisions of T such that each edge is subdivided at most c_T times, where c_T is a constant depending only on the radius of T).

In addition, two of us hope to show in a later paper [8] that every tree is χ -bounding that can be obtained from a tree of radius at most two by subdividing once *some of* the edges incident with the root, thus unifying the first two classes above; but the proof of that is long and difficult.

All the trees mentioned so far have the property that their vertices of degree greater than two are all clustered closely together. However, the conjecture is not known for any tree that contains a distant pair of vertices with degree more than two. The aim of this paper is to show the existence of such trees.

We begin with two special cases. Take a six-vertex path, and for each of its two middle vertices v say, add another vertex adjacent to v . We obtain a tree with eight vertices, and it was not previously known whether this tree is χ -bounding. More generally let us say a *two-legged caterpillar* is a tree obtained from a path by adding two more vertices, each with one neighbour in the path. We will prove:

1.2 *Every two-legged caterpillar is χ -bounding.*

A *star* is a tree in which one vertex is adjacent to all the others, and a *broom* is a tree obtained from a star by replacing one of its edges by a path of arbitrary length. A tree is a *subdivided star* if it has at most one vertex of degree at least three. (All brooms are χ -bounding [3], and indeed they are subdivided stars.) A *double broom* is a tree obtained from two disjoint stars by adding a path between the centres of the stars. We will show:




Figure 1: A 2-legged caterpillar and a double broom

1.3 *Every double broom is χ -bounding.*

Our two main theorems say that certain types of trees are χ -bounding, and the two results above are special cases. The first, implying 1.2, is

1.4 *Let H be a tree obtained by adding one vertex to a subdivided star. Then H is χ -bounding.*

The second, implying 1.3, is:

1.5 *Let H be a tree obtained from a subdivided star and a star by adding a path joining their centres. Then H is χ -bounding.*

The proofs of the two results are almost the same, and we will prove them together. The trees of 1.4 and 1.5 both have the property that they have only two vertices with degree more than two, and the length (k say) of the path between these two vertices turns out to be the key parameter. The proof method depends on whether there is a ball of radius at most k and large chromatic number or not, so let us make that precise. If v is a vertex of a graph G , $N^k(v)$ or $N_G^k(v)$ denotes the set of vertices of G with distance exactly k from v , and $N^k[v]$ or $N_G^k[v]$ denotes the set with distance at most k from v . We sometimes write $\chi(X)$ for $\chi(G[X])$ when there is no risk of ambiguity. If G is a nonnull graph and $k \geq 1$, we define $\chi^k(G)$ to be the maximum of $\chi(N^k[v])$ taken over all vertices v of G . (For the null graph G we define $\chi^k(G) = 0$.) If H is a subgraph of G and $u, v \in V(H)$, the distance between u, v in H may be greater than the distance between u, v in G , and the H -distance between u, v means the distance between u, v in H .

Let $d \geq 1$, and take a d -star (that is, a copy of the complete bipartite graph $K_{1,d}$). Now subdivide each of its edges $d - 1$ times; that is, replace each edge by a path of length d , joining the same pair of vertices, and internally pairwise disjoint. This produces a subdivided star and we call it a d -superstar. Let $k \geq 1$; we define a “ (k, d) -binary star” and a “ (k, d) -bristled star” as follows.

- Take the disjoint union of a d -superstar and a d -star, and join their centres with a path of length k . We call this tree a (k, d) -binary star.
- Take the disjoint union of a d -superstar S and a path T of length $d + 1$, and join the centre of S and the second vertex of T with a path of length k . We call this tree a (k, d) -bristled star.

Every tree H as in 1.5 is an induced subgraph of a (k, d) -binary star for some $k, d \geq 1$, and every tree H as in 1.4 is an induced subgraph of a (k, d) -bristled star for some $k, d \geq 1$. It therefore suffices to prove 1.5 and 1.4 for trees H that are (k, d) -binary stars, and for those that are (k, d) -bristled stars. Let us say a graph is (k, d) -starry if it has an induced subgraph that is a (k, d) -binary star and one that is a (k, d) -bristled star. The following is our main result, implying both 1.4 and 1.5.




Figure 2: A $(3, 2)$ -bristled star and a $(3, 2)$ -binary star

1.6 For all $\kappa \geq 0$ and $k, d \geq 1$, there exists $c \geq 0$ such that every graph G with $\omega(G) \leq \kappa$ and $\chi(G) > c$ is (k, d) -starry.

The proof of 1.6 is given at the end of section six.

2 Using criticality

Our main tool is a set of lemmas proved in this section, that if X is a subset of $V(G)$ of small chromatic number, and G itself has large chromatic number, and deleting X from G reduces the chromatic number, then there are useful subgraphs rooted at some vertex in X and growing out into $G \setminus X$. We begin with:

2.1 Let $d \geq 0$ be an integer, let G be a graph with chromatic number more than d , and let $X \subseteq V(G)$ be stable, such that $\chi(G \setminus X) < \chi(G)$. Then some vertex in X has at least d neighbours in $V(G) \setminus X$.

Proof. Let $\chi(G) = k + 1$, and so $k \geq d$. Let $\phi : V(G) \setminus X \rightarrow \{1, \dots, k\}$ be a k -colouring of $G \setminus X$. For each $x \in X$, if x has at most $k - 1$ neighbours in $V(G) \setminus X$ then we may choose $\phi(x) \in \{1, \dots, k\}$, different from $\phi(v)$ for each neighbour $v \in V(G) \setminus X$ of x ; and this extends ϕ to a k -colouring of G , which is impossible. Thus for some $x \in X$, x has at least $k \geq d$ neighbours in $V(G) \setminus X$. This proves 2.1. ■

If $X \subseteq V(G)$, let us say an X -split is a triple (x, y, Z) , where

- $x \in X$, $y \in V(G) \setminus X$, and $Z \subseteq V(G) \setminus (X \cup \{y\})$;
- x is adjacent to y and has at least one neighbour in Z ;
- y has no neighbours in Z ; and
- $G[Z]$ is connected.

Let us say $\chi(Z)$ is the *chromatic number* of an X -split (x, y, Z) . Next we need:

2.2 For all $c, \tau \geq 0$ there exists c' with the following property. Let G be a graph with chromatic number more than c' , such that $\chi^1(G) \leq \tau$; and let $X \subseteq V(G)$ be stable, such that $\chi(G \setminus X) < \chi(G)$. Then there is an X -split in G with chromatic number more than c .

Proof. Let $c' = (2c + 3\tau + 3)\tau$, let G be a graph with chromatic number more than c' and $\chi^1(G) \leq \tau$, and let $X \subseteq V(G)$ be stable, such that $\chi(G \setminus X) < \chi(G)$. We prove the result by induction on $|X|$.

Choose $x \in X$. If $\chi(G \setminus x) = \chi(G)$, let $G' = G \setminus x$, and $X' = X \setminus \{x\}$; then $\chi(G' \setminus X') < \chi(G')$, and the result follows from the inductive hypothesis. Thus we may assume that $\chi(G \setminus x) < \chi(G)$.

Let $k = \chi(G) - 1$, and let $\phi : V(G) \setminus \{x\} \rightarrow \{1, \dots, k\}$ be a k -colouring of $G \setminus x$. Let N denote the set of all neighbours of x . Then, since G does not admit a k -colouring, it follows that for $1 \leq i \leq k$ there exists $n \in N$ with $\phi(n) = i$. Let \overline{G} be the complement graph of G , and let N_1, \dots, N_t be the vertex sets of the components of $\overline{G}[N]$. Now $\chi(N) \leq \tau$, since $\chi^1(G) \leq \tau$. But for $1 \leq i < j \leq t$, every vertex of N_i is adjacent to every vertex of N_j , and so $\chi(N) = \sum_{1 \leq i \leq t} \chi(N_i)$. Consequently $\sum_{1 \leq i \leq t} \chi(N_i) \leq \tau$, and in particular, $t \leq \tau$ since each $\chi(N_i) > 0$. For $1 \leq i \leq t$ let D_i be the set $\{\phi(v) : v \in N_i\}$, that is, the set of colours that appear in N_i . Now D_1, \dots, D_t are pairwise disjoint and have union $\{1, \dots, k\}$, so we may assume that $|D_1| \geq k/t \geq 2c + 3\tau + 3$.

Let Y be the set of vertices of $G \setminus N_1$ that are adjacent to every vertex in N_1 ; thus $x \in Y$, and $N \setminus N_1 \subseteq Y$. Let W be the set of vertices of $G \setminus (N_1 \cup Y)$ that have neighbours in N_1 .

(1) *If $\chi(W) > c + 2\tau + 1$ then the theorem holds.*

We assume that $\chi(W) > c + 2\tau + 1$. Since X is stable, it follows that $\chi(W \setminus X) > c + 2\tau$; and since $\chi(N) \leq \tau$, $\chi(W \setminus (N \cup X)) > c + \tau$. Choose $q \in N_1$; then since $\chi(N^1(q)) \leq \tau$, the set of vertices in $W \setminus (N \cup X)$ that are nonadjacent to q has chromatic number more than c , and so there exists $Z \subseteq W \setminus (N \cup X)$ with $\chi(Z) > c$, such that $G[Z]$ is connected and q has no neighbour in Z . Let P be the set of vertices in N_1 that have a neighbour in Z , and $Q = N_1 \setminus P$; then $P \neq \emptyset$, since every vertex of Z has a neighbour in N_1 , and $Q \neq \emptyset$, since $q \in Q$. Since $\overline{G}[N_1]$ is connected, there exist $y, z \in N_1$, nonadjacent, such that $z \in P$ and $y \in Q$. But then $(x, y, Z \cup \{z\})$ is an X -split satisfying the theorem. This proves (1).

In view of (1), we assume henceforth that $\chi(W) \leq c + 2\tau + 1$.

(2) *Let C be the vertex set of a component of $G \setminus (N_1 \cup Y)$. If $C \cap W \neq \emptyset$ and $\chi(C) > 2c + 2\tau + 2$ then the theorem holds.*

For then $\chi(C \setminus (W \cup X)) > c$, and so there is a subset $Z \subseteq C \setminus (W \cup X)$ with $\chi(Z) > c$ such that $G[Z]$ is connected. Since $Z \subseteq C$ and $C \cap W \neq \emptyset$, there is a path of $G[C]$ between W and Z ; choose such a path, P say, minimal. Now P has length at least one, since $Z \cap W = \emptyset$. Let w be the end of P in W . It follows that no vertex of P different from w has a neighbour in N_1 . Since $\overline{G}[N_1]$ is connected and w has a neighbour and a non-neighbour in N_1 , there exist nonadjacent $y, z \in N_1$ such that w is adjacent to z and not to y .

Suppose first that no vertex of P belongs to X . Then $(x, y, Z \cup V(P) \cup \{z\})$ is the desired X -split. We may assume therefore that some vertex of P belongs to X . Choose $x' \in X \cap V(P)$ such that the subpath of P (P' say) between x' and Z is minimal. If $x' \neq w$, let y' be the vertex of P adjacent to x' that does not belong to $V(P')$, and if $x' = w$, let $y' = z$. In either case, $y' \notin X$, since X is stable. Then $\{x', y', (V(P') \setminus \{x'\}) \cup Z\}$ is the desired X -split. This proves (2).

Let U be the union of the vertex sets of all components of $G \setminus (N_1 \cup Y)$ that have nonempty intersection with W .

(3) *If $v \in V(G) \setminus (N_1 \cup U \cup \{x\})$ and has a neighbour $u \in N_1 \cup U \cup \{x\}$, then $\phi(v) \notin D_1$.*

We claim that $v \in Y$; for if $u \in N_1$ then v has a neighbour in N_1 and $v \notin U \cup N_1 \cup \{x\}$, so $v \in Y$; if $u \in U$ then $v \in N_1 \cup Y$ since $v \notin U$, and so again $v \in Y$; and if $u = x$ then $v \in N^1(x) \subseteq N_1 \cup Y$, and again $v \in Y$. Thus $v \in Y$. Consequently $\phi(v)$ is different from $\phi(w)$ for every $w \in N_1$; and so from the definition of D_1 it follows that $\phi(v) \notin D_1$. This proves (3).

By (2) we may assume (for a contradiction) that $\chi(U) \leq 2c + 2\tau + 2$. It follows that $\chi(N_1 \cup U \cup \{x\}) \leq 2c + 3\tau + 3$. Consequently $G[N_1 \cup U \cup \{x\}]$ admits a colouring ψ using only the colours in D_1 , since $|D_1| \geq 2c + 3\tau + 3$. For each $v \in V(G)$, define $\phi'(v) = \phi(v)$ if $v \notin N_1 \cup U \cup \{x\}$, and $\phi'(v) = \psi(v)$ if $v \in N_1 \cup U \cup \{x\}$. By (3) this gives a k -colouring of G , which is impossible. This proves 2.2. ■

To use this, we combine it with a version of Gyárfás' path theorem (see [1] for this version):

2.3 *Let G be a graph, let $k \geq 0$, let $C \subseteq V(G)$, and let $x_0 \in V(G) \setminus C$, such that $G[C]$ is connected, x_0 has a neighbour in C , and $\chi(C) > k\chi^1(G)$. Then there is an induced path $x_0 \cdots x_k$ of G where $x_1, \dots, x_k \in C$, and a subset C' of C , with the following properties:*

- $x_0, \dots, x_k \notin C'$;
- $G[C']$ is connected;
- x_k has a neighbour in C' , and x_0, \dots, x_{k-1} have no neighbours in C' ; and
- $\chi(C') \geq \chi(C) - k\chi^1(G)$.

We deduce:

2.4 *For all $d, \tau \geq 0$ there exists c' with the following property. Let G be a graph with chromatic number more than c' , such that $\chi^1(G) \leq \tau$; and let $X \subseteq V(G)$ be stable, such that $\chi(G \setminus X) < \chi(G)$. Then there is a vertex $x \in X$, an induced path P of length d with one end x and no other vertices in X , and a vertex $y \in V(G) \setminus X$ that is adjacent to x and has no other neighbour in $V(P)$.*

Proof. Let $c = d\tau$ and let c' satisfy 2.2. We claim that c' satisfies 2.4. For let G, X be as in the theorem, with $\chi(G) > c'$. By 2.2 there is an X -split (x, y, Z) with chromatic number more than $c = d\tau$. By 2.3 there is an induced path P of $G[Z \cup \{x\}]$ with one end x , of length d . But then x, y, P satisfy 2.4. This proves 2.4. ■

We combine 2.1 and 2.4 in the following. If $X \subseteq V(G)$, we say that $x \in X$ is *d -equipped in $V(G) \setminus X$* if x has at least d neighbours in $V(G) \setminus X$, pairwise nonadjacent, and there is an induced path P of length d with one end x and no other vertices in X , and a vertex $y \in V(G) \setminus (X \cup V(P))$ that is adjacent to x and has no other neighbour in $V(P)$. We speak of the set of d pairwise nonadjacent neighbours, the path P and the vertex y as the *parts of the equipment* of x .

2.5 *For all $d, \tau \geq 0$ there exists c with the following property. Let G be a graph with chromatic number more than c , such that $\chi^1(G) \leq \tau$; and let $X \subseteq V(G)$ be stable, such that $\chi(G \setminus X) < \chi(G)$. Then there is a vertex $x \in X$ that is d -equipped in $V(G) \setminus X$.*

Proof. Choose $d' \geq 0$ such that every graph with d' vertices has either a clique of cardinality $\tau + 1$ or a stable set of size d . Choose $c \geq d'$ such that 2.4 holds with c' replaced by c . Let X_1 be the set of vertices in X that have at least d' neighbours in $V(G) \setminus X$; and let X_2 be the set of vertices $x \in X$ such that there is an induced path P of length d with one end x and no other vertices in X , and a vertex $y \in V(G) \setminus X$ that is adjacent to x and has no other neighbour in $V(P)$. Suppose for a contradiction that $X_1 \cap X_2 = \emptyset$. Since X is stable, it follows that no vertex in X_2 has d' neighbours in $V(G) \setminus X_2$; and so by 2.1, it follows that $\chi(G) = \chi(G \setminus X_2)$. By 2.4 applied to $X \setminus X_2$ in the graph $G \setminus X_2$, it follows that

$$\chi(G \setminus X_2) = \chi((G \setminus X_2) \setminus (X \setminus X_2)),$$

contradicting that $\chi(G \setminus X) < \chi(G)$. Hence there exists $x \in X_1 \cap X_2$. The choice of d' implies that x has d pairwise nonadjacent neighbours in $V(G) \setminus X$, so x is d -equipped in $V(G) \setminus X$. This proves 2.5. \blacksquare

Next we need a version in which X may not be stable:

2.6 *For all $a, d, \tau \geq 0$ there exists c with the following property. Let G be a graph with chromatic number more than c , such that $\chi^1(G) \leq \tau$; and let $X \subseteq V(G)$ with $\chi(X) \leq a$, such that $\chi(G \setminus X) < \chi(G)$. Then there is a vertex $x \in X$ that is d -equipped in $V(G) \setminus X$.*

Proof. Choose b such that 2.5 holds with c replaced by b ; and let $c = ab$. We claim that c satisfies the theorem. Let G be a graph with chromatic number more than c , and let $X \subseteq V(G)$ with $\chi(X) \leq a$, such that $\chi(G \setminus X) < \chi(G)$. Let $\chi(G) = k + 1$, and let $\phi : V(G) \setminus X \rightarrow \{1, \dots, k\}$ be a k -colouring of $G \setminus X$. Let (X_1, \dots, X_a) be a partition of X into a stable sets. For $1 \leq i \leq a$ let $A_i = \{(i-1)b + 1, \dots, ib\}$, and let Y_i be the set of vertices $v \in V(G) \setminus X$ with $\phi(v) \in A_i$. If $\chi(X_i \cup Y_i) \leq b$ for each i then $\chi(G) = \chi(G \setminus X)$, a contradiction; so we may assume that $\chi(X_1 \cup Y_1) > b$. Since $\chi(Y_1) \leq b$, and X_1 is stable, the choice of b implies that there is a vertex $x \in X_1$ that is d -equipped in $Y_1 \setminus X_1$ and hence in $V(G) \setminus X$. This proves 2.6. \blacksquare

We can iterate this (and we also throw in a bounded set of “forbidden vertices” B , but for most of the applications $B = \emptyset$).

2.7 *For all $a, b, d, \tau \geq 0$ there exists c with the following property. Let G be a graph with chromatic number more than c , such that $\chi^1(G) \leq \tau$; and let $X \subseteq V(G)$ such that $\chi(G \setminus X) < \chi(G)$ and $\chi(X) \leq a$. Let $B \subseteq V(G) \setminus X$ with $|B| \leq b$. Let M denote the set of vertices in $V(G) \setminus X$ with a neighbour in X . Then there is a vertex $x \in X$ that is d -equipped in $V(G) \setminus X$, and there is a neighbour x' of x in $M \setminus B$, such that x' is d -equipped in $B \cup (V(G) \setminus (X \cup N^1(x)))$.*

Proof. By increasing d , we may assume that $d > b$. Choose c_1 such that 2.6 holds with a, c replaced by τ, c_1 . Choose c such that 2.6 holds with a, c replaced by $c_1 + a + b, c$. Now let G, X, M be as in the theorem with $\chi(G) > c$. Let X_1 be the set of vertices in X that are d -equipped in $V(G) \setminus X$, and let M_1 be the set of vertices in $V(G) \setminus X$ with a neighbour in X_1 .

Suppose first that $\chi(X \cup M_1) \leq c_1 + a + b$. Let $X' = (X \cup M_1) \setminus B$; then $X \subseteq X'$, and so $\chi(G \setminus X') < \chi(G)$. From 2.6, some vertex $x' \in X'$ is d -equipped in $V(G) \setminus X'$. Now $x' \notin X \setminus X_1$, since no vertex in $X \setminus X_1$ is d -equipped in $V(G) \setminus X$. Also x' has at least d neighbours in $(V(G) \setminus (X \cup M_1)) \cup B$, and therefore at least one neighbour in $V(G) \setminus (X \cup M_1)$, since $|B| \leq b < d$; and no vertex in X_1

has any neighbour in $V(G) \setminus (X \cup M_1)$. Consequently $x' \notin X_1$, and so $x' \in M_1 \setminus B$. Choose $x \in X_1$ adjacent to x' . Then $N^1(x) \setminus X \subseteq M_1$, and so x' is d -equipped in $B \cup (V(G) \setminus (X \cup N^1(x)))$.

We may assume therefore that $\chi(X \cup M_1) > c_1 + a + b$, and so $\chi(M_1 \setminus B) > c_1$. Choose $Z \subseteq M_1 \setminus B$ minimal with $\chi(Z) > c_1$. Choose $x \in X_1$ with a neighbour in Z , and let Y be the set of neighbours of x in Z . Then $\chi(Y) \leq \tau$, and so by 2.6 applied to $G[Z], Y$, there exists $x' \in Y$ that is d -equipped in $Z \setminus Y$. Since x has no neighbours in $Z \setminus Y$, it follows that x, x' satisfy the theorem. This proves 2.7. ■

3 k -balls with large chromatic number

Let $k, d \geq 1$ be integers. A (k, d) -broom is a tree obtained from a path $v_0 - v_1 - \dots - v_k$ by adding d new vertices, each adjacent to v_k . We call v_0 the *root* of the broom. A (k, d) -bristle is obtained from a path $v_0 - v_1 - \dots - v_{k+d}$ by adding one new vertex adjacent to v_k . We call v_0 the *root* of the bristle. By a (k, d) -broom in G we mean a (k, d) -broom that is an induced subgraph of G , and we use similar language for other kinds of tree.

3.1 *Let $k, d, \tau \geq 1$; then there exists c with the following property. Let G be a graph with $\chi^1(G) \leq \tau$, and let $z \in V(G)$, such that $\chi(N^k[z]) > c$. Then there is a (k, d) -broom and a (k, d) -bristle in G , both with root z .*

Proof. Let $c_1 = 2\tau$, and inductively for $i \geq 2$, choose $c_i \geq 2dc_{i-1}$ such that 2.6 is satisfied with a, c replaced by $c_{i-1}, c_i/2$ respectively. We prove by induction on k (for the given value of d) that setting $c = c_k$ satisfies the theorem. Thus we may assume that either $k = 1$ or the claim holds for $k - 1$.

Let G satisfy $\chi^1(G) \leq \tau$, and let $z \in V(G)$, such that $\chi(N^k[z]) > c_k$. For each $s \geq 0$, let $L_s = N^s(z)$. Since $\chi(N^k[z]) > c_k$, there exists $s \geq 0$ with $s \leq k$ such that $\chi(L_s) > c_k/2$. Since $\chi(L_0), \chi(L_1) \leq \tau$, it follows that $s \geq 2$; and so $k \geq 2$, and the claim holds for $k - 1$.

Choose $S \subseteq L_s$ minimal such that $\chi(S) = \chi(L_s)$. Choose $u \in S$, and choose $v \in L_1$ joined to u by a path of length $s - 1$.

Let G' be the graph $G \setminus (L_0 \cup L_1 \setminus \{v\})$. If $\chi(N_{G'}^{k-1}[v]) > c_{k-1}$, then from the inductive hypothesis applied to G' , there is a $(k - 1, d)$ -broom and a $(k - 1, d)$ -bristle in G' with root v . But then adding the edge zv gives the desired (k, d) -broom and (k, d) -bristle in G with root z . We may therefore assume that $\chi(N_{G'}^{k-1}[v]) \leq c_{k-1}$.

Let X be the set of vertices in S that have G' -distance at most $k - 1$ from v . Thus $X \subseteq N_{G'}^{k-1}[v]$, and so $\chi(X) \leq c_{k-1}$. Now $u \in X$, since $s \leq k$, and so $X \neq \emptyset$. From the minimality of S , $\chi(S \setminus X) < \chi(S)$.

By 2.6, since $\chi(S) \geq c_k/2 \geq c_{k-1}d$, and $\chi(X) \leq c_{k-1}$, it follows that some vertex $x \in X$ is d -equipped in $S \setminus X$. Since $x \in X$, there is an induced path P of length at most $k - 1$ such that $V(P) \cap (L_0 \cup L_1) = \{v\}$. Since x has a neighbour in $S \setminus X$, it follows that the length of P is exactly $k - 1$, and no vertex of P different from x has a neighbour in $S \setminus X$. Also z has no neighbours in $S \setminus X$ since $s \geq 2$. But then P together with the edges zv and the various parts of the equipment of x gives a (k, d) -broom and a (k, d) -bristle, both with root z . This proves 3.1. ■

If A, B are disjoint subsets of $V(G)$, we say that A *covers* B if every vertex in B has a neighbour in A .

3.2 For all $c, \tau \geq 0$ and $d \geq 1$ and $k \geq r \geq 2$, there exists $c' \geq 0$ with the following property. Let G be a graph with $\chi^{r-1}(G) \leq \tau$, and let $z \in V(G)$ such that $\chi(N_G^r(z)) > c'$. Then there is a (k, d) -broom H_1 and a (k, d) -bristle H_2 in G , both with root z , and a subset W of $V(G)$, such that

- $z \in W$, and no other vertex of $H_1 \cup H_2$ belongs to W or has a neighbour in $W \setminus \{z\}$;
- $\chi(N_{G'}^r(z)) > c$, where G' is the graph $G[W]$.

Proof. Choose c_1 such that 3.1 is satisfied with c, k replaced by c_1, k . Let

$$c' = ((2k + 2d + 3)\tau + c + c_1)\tau.$$

Now let G, z be as in the theorem. For $u \in N^1(z)$ and $v \in N^i(z)$ where $i \geq 1$, we say that u is an *ancestor* of v and v is a *descendant* of u if there is a path of length $i - 1$ between u, v . Since $\chi(N^1(z)) \leq \tau$, there is a partition of $N^1(z)$ into τ stable sets; and since every vertex in $N^r(z)$ has an ancestor in $N^1(z)$, there is a stable set $L_1 \subseteq N^1(z)$ such that, if L_r denotes the set of vertices in $N^r(z)$ with an ancestor in L_1 , then

$$\chi(L_r) > c'/\tau = (2k + 2d + 3)\tau + c + c_1.$$

For $2 \leq i \leq r - 1$, let L_i be the set of descendants in $N^i(z)$ of members of L_1 .

Choose $B \subseteq L_1$ maximal such that the set of vertices in L_r with no ancestor in B has chromatic number at least c_1 . Let $L'_0 = \{z\}$ and for $1 \leq i \leq r$, let L'_i be the set of vertices in L_i with no ancestor in B . It follows that $L'_1 = L_1 \setminus B$, and $\chi(L'_r) > c_1$, and for $1 \leq i \leq r$, L'_{i-1} covers L'_i . Let

$$G_1 = G[L'_0 \cup \dots \cup L'_r].$$

Then $\chi(N_{G_1}^r(z)) > c_1$, and so $\chi(N_{G_1}^k[z]) > c_1$. It follows from 3.1 applied to G_1 , and from the choice of c_1 , that there is a (k, d) -broom H_1 and a (k, d) -bristle H_2 in G , both with root z , and both with vertex set a subset of $L'_0 \cup \dots \cup L'_r$.

In particular, there is a vertex in $L_1 \setminus B$, v say, and since the set of descendants of v in L_r has chromatic number at most τ , the maximality of B implies that $\chi(L'_r) \leq c_1 + \tau$. Consequently

$$\chi(L_r \setminus L'_r) > 2(k + d + 1)\tau + c.$$

Since

$$|V(H_1) \cup V(H_2)| \leq 2(k + d + 1),$$

the set of vertices with G -distance at most $r - 1$ from some vertex in $H_1 \cup H_2$ has chromatic number at most $2(k + d + 1)\tau$. Consequently there exists $L''_r \subseteq L_r \setminus L'_r$ with $\chi(L''_r) > c$ such that every vertex in L''_r has G -distance at least r from every vertex of $H_1 \cup H_2$. Let $L''_0 = \{z\}$, and for $1 \leq i \leq r - 1$, let L''_i be the set of vertices in $L_i \setminus L'_i$ that have G -distance at least i from every vertex of $H_1 \cup H_2$. We claim that L''_{i-1} covers L''_i for $1 \leq i \leq r$. Certainly L''_0 covers L''_1 , so we may assume that $i \geq 2$. Let $v \in L''_i$. Since $v \notin L'_i$, v has an ancestor in B , and since $i \geq 2$ it follows that v has a neighbour $u \in L_{i-1}$ with an ancestor in B . Consequently $u \in L_{i-1} \setminus L'_{i-1}$. But since the G -distance from v to $V(H_1 \cup H_2)$ is at least i , and u, v are adjacent, it follows that the G -distance from u to $V(H_1 \cup H_2)$ is at least $i - 1$, and so $u \in L''_{i-1}$. This proves that L''_{i-1} covers L''_i for $1 \leq i \leq r$. Let $W = L''_0 \cup L''_1 \cup \dots \cup L''_r$, and let $G' = G[W]$. Then $\chi(N_{G'}^r(z)) > c$. It remains to show that no vertex

of $W \setminus \{z\}$ is adjacent to a vertex of $H_1 \cup H_2$ different from z . Let $v \in L_i''$ say, where $i \geq 1$, and suppose v is adjacent to some vertex $u \in V(H_1 \cup H_2) \setminus \{z\}$. Since v has G -distance at least i from $V(H_1 \cup H_2)$, it follows that $i = 1$, and so $v \in B$ and $u \in L_1' \cup L_2'$. But $u \notin L_1$ since L_1 is stable, and $u \notin L_2'$ since no vertex of L_2' has an ancestor in B , a contradiction. Thus there is no such pair u, v . This proves 3.2. \blacksquare

Let $s \geq 1$, for $1 \leq i \leq s$ let $k_i, d_i \geq 1$, and let H_i be either a (k_i, d_i) -broom or a (k_i, d_i) -bristle; and let H be the rooted tree obtained from the disjoint union of H_1, \dots, H_s by identifying the roots of H_1, \dots, H_s to form the root of H . Let us call such a tree H the *rooted sum* of H_1, \dots, H_s .

3.3 *With notation as above, let H be the rooted sum of H_1, \dots, H_s ; and let $1 \leq r \leq \min(k_1, \dots, k_s)$. For all $\tau \geq 0$ and $r \geq 1$ there exists c with the following property. Let G be a graph with $\chi^1(G) \leq \tau$, and let $z \in V(G)$ be a vertex such that $\chi(N^r[z]) > c$. Then there is an induced subgraph of G isomorphic to H with root z .*

Proof. We may assume (by replacing d_1, \dots, d_s by their maximum) that $d_1 = \dots = d_s = d$ say; for fixed d , we proceed by induction on r ; and for fixed d, r , by induction on s . If $r = 1$ then the claim holds, setting $c = \tau$; so we may assume that $r > 1$, and (for the same values of d, s and k_1, \dots, k_s) the result holds with r, c replaced by $r - 1, c_1$. Let H' be the rooted sum of H_1, \dots, H_{s-1} (or if $s = 1$, let H' be a one-vertex graph, with that vertex as root). If $s = 1$ let $c_2 = 0$, and if $s > 1$ then by induction on s , we may choose c_2 such that the result holds (for the same values of d, r and k_1, \dots, k_{s-1}) with s, H, c replaced by $s - 1, H', c_2$. By 3.2 we may choose c_3 such that 3.2 holds with τ, c, k, r, c' replaced by c_1, c_2, k_s, r, c_3 . We claim that setting $c = c_1 + c_3$ satisfies the theorem.

For let G, z be as in the theorem, with $\chi(N^r[z]) > c_1 + c_3$. If there is a vertex z' with $\chi(N^{r-1}[z']) > c_1$ then the result follows from the choice of c_1 . Thus we may assume that $\chi^{r-1}(G) \leq c_1$; and in particular $\chi(N^r(z)) > c_3$. From 3.2 and the choice of c_3 , there is an induced subgraph J of G isomorphic to H_s , with root z , and a subset W of $V(G)$, such that

- $z \in W$, and no other vertex of J belongs to W or has a neighbour in $W \setminus \{z\}$; and
- $\chi(N_{G'}^r(z)) > c_2$, where G' is the graph $G[W]$.

From the choice of c_2 , there is an induced subgraph J' of G' , isomorphic to H' , with root z . But then the union of J and J' satisfies the theorem. This proves 3.3. \blacksquare

4 Spires and cathedrals

Let G be a graph, let P be an induced path of G , and $A, B \subseteq V(G)$, such that:

- $G[A]$ is connected;
- $A \cap B = \emptyset$;
- A covers B ;
- $V(P) \cap B = \emptyset$, and there is an end z of P in A such that $V(P) \cap A = \{z\}$; and

- no vertex in $V(P) \setminus \{z\}$ has any neighbours in $(A \cup B) \setminus \{z\}$.

In this situation we say that $\mathcal{S} = (P, A, B)$ is a *spire* of *height* d , where d is the length of P . We define $V(\mathcal{S}) = A \cup B \cup V(P)$. If $C \subseteq V(G)$, we say that the spire *dominates* C if

- C is disjoint from $V(\mathcal{S})$;
- there are no edges between $A \cup V(P)$ and C ; and
- B covers C .

A *cathedral* is a sequence of spires $(\mathcal{S}_1, \dots, \mathcal{S}_n)$, such that

- for $1 \leq i < j \leq n$, $V(\mathcal{S}_i) \cap V(\mathcal{S}_j) = \emptyset$; and
- for $1 \leq i < j \leq n$, if $u \in V(\mathcal{S}_i)$ and $v \in V(\mathcal{S}_j)$ are adjacent then $u \in B_i$ and $v \in A_j \cup B_j$

where $\mathcal{S}_i = (P_i, A_i, B_i)$ for $1 \leq i \leq n$. We say the cathedral is *free* if

- for $1 \leq i < j \leq n$, if $u \in V(\mathcal{S}_i)$ and $v \in V(\mathcal{S}_j)$ are adjacent then $u \in B_i$ and $v \in B_j$.

A cathedral has *height* d if each of its spires has height d , and *length* n if it has n spires. We say a cathedral *dominates* a set $C \subseteq V(G)$ if each of its spires dominates C .

4.1 *For all $\tau \geq 0$ and $k, d \geq 1$ there exist $c, n \geq 0$ with the following property. Let G be a graph with $\chi^k(G) \leq \tau$, and let $(\mathcal{S}_1, \dots, \mathcal{S}_n)$ be a free cathedral in G of height d , dominating a set C with $\chi(C) > c$. Then G is (k, d) -starry.*

Proof. Let $m = 2d^2$. Choose $n \geq 0$ such that every graph with n vertices has either a clique of cardinality $\tau + 1$ or a stable set of cardinality m . Choose c_0 such that 2.6 holds with a, c replaced by τ, c_0 . If $k \geq 2$, choose c such that 2.7 holds with a, b, c and d replaced by $(m + 1)\tau + c_0, 0, c$ and d respectively; and if $k = 1$, choose c such that 2.7 holds with a, b, c and d replaced by $(m + 1)\tau + c_0, mn, c$ and $d + mn$ respectively. Now let G and $(\mathcal{S}_1, \dots, \mathcal{S}_n)$ be as in the theorem, and let $\mathcal{S}_i = (P_i, A_i, B_i)$ for $1 \leq i \leq n$. We may assume that C is minimal such that $\chi(C) > c$, and in particular, $G[C]$ is connected. Choose $r_0 \in C$, and for $1 \leq i \leq n$ let b_i be a neighbour of r_0 in B_i . Since P_i has length d , and A_i covers B_i , and $G[A_i \cup V(P_i)]$ is connected, there is an induced path Q_i of $G[A_i \cup V(P_i) \cup \{b_i\}]$ of length $d - 1$ with one end b_i . By the choice of n , we may assume that b_1, \dots, b_m are pairwise nonadjacent.

Let R be the set of vertices in C that are adjacent to at least $2d$ of b_1, \dots, b_m ; thus $r_0 \in R$. Let S be the set of vertices in C that are adjacent to at least one and to at most $2d - 1$ of b_1, \dots, b_m . Let $L_0 = R$, and let L_1 be the set of all vertices in $C \setminus L_0$ with a neighbour in L_0 . Let L_2 be the set of all vertices in $C \setminus (L_0 \cup L_1)$ that either belong to S or have a neighbour in L_1 . For $i \geq 3$, inductively let L_i be the set of vertices in $C \setminus (L_0 \cup \dots \cup L_{i-1})$ with a neighbour in L_{i-1} . Let W be the set of vertices in C with G -distance at most k from a vertex in $\{r_0, b_1, \dots, b_m\}$.

(1) *If $\chi(L_k \setminus W) > c_0$ then the theorem holds.*

Suppose that $\chi(L_k \setminus W) > c_0$, and choose $Y \subseteq L_k \setminus W$ minimal with $\chi(Y) > c_0$. In particular $(R \cup S) \cap Y = \emptyset$, since $R \cup S \subseteq W$. There is no path of $G[C]$ of length at most $k - 2$ between Y and

S , since $Y \cap W = \emptyset$. Since $Y \subseteq L_k$, no vertex in Y has $G[C]$ -distance less than k from a vertex in R ; and since $y \cap W = \emptyset$, it follows that every vertex in Y has $G[C]$ -distance exactly k from some vertex in R . Since $Y \neq \emptyset$, there exists $r \in R$ such that $X \neq \emptyset$, where X denotes the set of vertices in Y with $G[C]$ -distance k from r . From 2.6 applied to X and $G[Y]$, there exists $x \in X$ that is d -equipped in $Y \setminus X$. Let P be a path of $G[C]$ of length k joining r, x . Since no vertex in Y has G -distance at most k from a vertex in $\{r_0, b_1, \dots, b_m\}$, it follows that no vertex of P belongs to S . Since $r \in R$, r is adjacent to at least d of b_1, \dots, b_m , say b_1, \dots, b_d . But then adding the parts of the equipment of x to the tree formed by the union of P , the paths Q_1, \dots, Q_d and the edges rb_1, \dots, rb_d , gives a (k, d) -binary star and a (k, d) -bristled star. This proves (1).

Thus we may assume that $\chi(L_k \setminus W) \leq c_0$. Let $X = L_0 \cup \dots \cup L_k$. Since $\chi(W) \leq (m+1)\tau$, and $L_0 \cup \dots \cup L_{k-1} \subseteq W$, it follows that $\chi(X) \leq (m+1)\tau + c_0$. By 2.7 applied to $G[C]$ and X , there exists $x \in X$ that is d -equipped in $C \setminus X$, and there exists $x' \in C \setminus X$ adjacent to x such that x' is d -equipped in $C \setminus (X \cup N^1(x))$. Since $x' \notin X$, it follows that $x' \in L_{k+1}$, and $x \in L_k$, and so there is an induced path T of $G[C]$ with ends x', u say, such that $x \in V(T)$, and either

- $k \geq 2$, and T has length $k - 1$ and $u \in S$, or
- T has length $k + 1$, and $u \in R$.

In either case no vertex of $T \setminus u$ is in R .

Suppose that the first bullet holds. It follows that no vertex of T different from u is in S ; and since $k \geq 2$ and so $S \subseteq X$, it follows that no vertex of the equipment of x belongs to S . Since $u \in L_2$, it follows that u is nonadjacent to r_0 , and since $u \in S$ we may assume that u is nonadjacent to b_1, \dots, b_d and adjacent to b_{d+1} . But then the union of the paths Q_1, \dots, Q_d , the edges r_0b_1, \dots, r_0b_d , the edge r_0b_{d+1} , the edge $b_{d+1}u$, and the path T can be extended to a (k, d) -binary star and to a (k, d) -bristled star by adding appropriate parts of the equipment of x .

We may therefore assume that the second bullet holds. Let v be the vertex of T adjacent to u , and w the other neighbour of v in T (this exists since T has length $k + 1 \geq 2$). Since $x' \in L_{k+1}$, no vertex of T different from v, w belongs to S .

(2) *If $S \subseteq X$ then the theorem holds.*

Since $S \subseteq X$, it follows that $x' \notin S$. Moreover, if $w \in S$ then $T \setminus \{u, v\}$ satisfies the first bullet above and we are done; so we may assume that $w \notin S$. We may assume that u is adjacent to b_1, \dots, b_{2d} . If v is nonadjacent to at least d of b_1, \dots, b_{2d} , say to b_1, \dots, b_d , then the union of the paths Q_1, \dots, Q_d , the edges ub_1, \dots, ub_d , the path $T \setminus x'$, and appropriate parts of the equipment of x , gives a (k, d) -binary star and a (k, d) -bristled star. Thus we may assume that v is adjacent to at least d of b_1, \dots, b_{2d} , say to b_1, \dots, b_d . But then the union of the paths Q_1, \dots, Q_d , the edges vb_1, \dots, vb_d , the path $T \setminus u$, and appropriate parts of the equipment of x' , gives a (k, d) -binary star and a (k, d) -bristled star. This proves (2).

In view of (2), we may assume that $S \not\subseteq X$, and consequently $k = 1$. Choose $s \in S \setminus X$; then s is nonadjacent to r_0 , and we may assume that s is nonadjacent to b_1, \dots, b_d and adjacent to b_{d+1} , and so the union of the paths Q_1, \dots, Q_{d+1} , the edges $r_0b_1, \dots, r_0b_{d+1}$ and the edge $b_{d+1}s$ gives a $(1, d)$ -bristled star. It remains to find a $(1, d)$ -binary star. To do so, we need to apply 2.7 more

carefully, using the “forbidden vertices” feature of 2.7. First we need:

(3) *We may assume that $|S \setminus L_1| < mn$.*

If not, then some vertex in $\{b_1, \dots, b_m\}$ (say b_{d+1}) is adjacent to more than n vertices in $S \setminus L_1$, and from the definition of n , at least d of these neighbours are pairwise nonadjacent, say y_1, \dots, y_d . For $1 \leq i \leq d$, y_i is adjacent to fewer than $2d$ of $\{b_1, \dots, b_m\}$, and since $m = 2d^2$, we may assume that y_1, \dots, y_d are all nonadjacent to all of b_1, \dots, b_d . But then the union of the paths Q_1, \dots, Q_d , the edges $r_0b_1, \dots, r_0b_d, r_0b_{d+1}$, and the edges $b_{d+1}y_1, \dots, b_{d+1}y_d$ gives a $(1, d)$ -binary star. This proves (3).

Now let us apply 2.7 again, to $X = L_0 \cup L_1$ and $G[C]$, setting $B = S \setminus L_1$ and replacing d by $mn + d$. We obtain a pair x, x' as before, where x is $(mn + d)$ -equipped (and hence d -equipped) in $C \setminus X$, and $x' \in L_2 \setminus S$, adjacent to x , such that x' is $(mn + d)$ -equipped in $S \cup (C \setminus (X \cup N^1(x)))$. In particular, since x has at least $mn + d$ pairwise nonadjacent neighbours in L_2 , and at most mn of them belong to S , it follows that x has d pairwise nonadjacent neighbours $y_1, \dots, y_d \in L_2 \setminus S$. Also, $x' \in L_2 \setminus S$, and by the same argument x' has d pairwise nonadjacent neighbours $y'_1, \dots, y'_d \in C \setminus (X \cup S \cup N^1(x))$. But now we finish the proof as in (2). More precisely, let T be the path $u-x-x'$, where $u \in R$. We may assume that u is adjacent to b_1, \dots, b_{2d} . If x is nonadjacent to at least d of b_1, \dots, b_{2d} , say to b_1, \dots, b_d , then the union of the paths Q_1, \dots, Q_d , the edges ub_i ($1 \leq i \leq d$), the edge ux , and the edges xy_1, \dots, xy_d gives a $(1, d)$ -binary star. If x is adjacent to at least d of b_1, \dots, b_{2d} , say to b_1, \dots, b_d , then the union of the paths Q_1, \dots, Q_d , the edges xb_i ($1 \leq i \leq d$), the edge xx' , and the edges $x'y'_1, \dots, x'y'_d$ gives a $(1, d)$ -binary star. This proves 4.1. \blacksquare

We can extend this result to cathedrals that are not free, as follows.

4.2 *For all $\tau \geq 0$ and $k, d \geq 1$ there exist $c, n \geq 0$ with the following property. Let G be a graph with $\chi^k(G) \leq \tau$, and let $(\mathcal{S}_1, \dots, \mathcal{S}_n)$ be a cathedral in G of height d , dominating a set C with $\chi(C) > c$. Then G is (k, d) -starry.*

Proof. Choose c_0, n_0 such that 4.1 holds with c, n replaced by c_0, n_0 . Let $n = dn_0$, and choose $c \geq 2^{n^2}c_0$ such that 2.6 holds with a, c replaced by τ, c . We claim that n, c satisfy 4.2. For let G be a graph with $\chi^k(G) \leq \tau$, and let $(\mathcal{S}_1, \dots, \mathcal{S}_n)$ be a cathedral in G of height d , dominating a set C with $\chi(C) > c$. Let $\mathcal{S}_i = (P_i, A_i, B_i)$ for $1 \leq i \leq n$. We may assume that C is minimal such that $\chi(C) > c$; and for $1 \leq i \leq n$, that every vertex in B_i has a neighbour in C (because any vertex in B_i with no neighbour in C can be removed).

(1) *We may assume that there exists i with $1 \leq i \leq n$, and a vertex $v \in B_i$ that has neighbours in at least d of A_{i+1}, \dots, A_n .*

Suppose not. For $1 \leq i \leq n$ and each $v \in B_i$, let $J(v)$ be the set of j with $i < j \leq n$ such that v has a neighbour in A_j . Thus each $|J(v)| \leq d - 1$. For each vertex $u \in C$, choose a neighbour $v_i \in B_i$ of u , and let $S(u)$ be the sequence $(J(v_1), \dots, J(v_n))$. There are at most 2^{n^2} possibilities for this sequence, and so there exists $C' \subseteq C$ with $\chi(C') \geq \chi(C)2^{-n^2} > c_0$ and a sequence S such that $S(v) = S$ for all $v \in C'$. Let $S = (J_1, \dots, J_n)$. Since there exists $v \in B_i$ with $J(v) = J_i$, it follows that $|J_i| < d$ for $1 \leq i \leq n$. Let H be the digraph with vertex set $\{1, \dots, n\}$ in which j is adjacent

from i if $i < j$ and $j \in J_i$. Since H has no directed cycles and all vertices have outdegree less than d , the graph underlying H has chromatic number at most d . Consequently it has a stable set I of cardinality n_0 , since $n = dn_0$. For each $i \in I$, let B'_i be the set of vertices $v \in B_i$ such that $J(v) = J_i$. It follows that B'_i covers C' . But then for each $i \in I$, (P_i, A_i, B'_i) is a spire dominating C' , and the sequence of these spires is a free cathedral of height d and length n_0 dominating C' , and the result follows from 4.1. This proves (1).

Choose v, i as in (1), where $v \in B_i$ has neighbours in at least d of A_{i+1}, \dots, A_n , say in A_{j_1}, \dots, A_{j_d} . If v has a neighbour in A_j for some $j > i$, then since v has no neighbour in $V(P_j) \setminus A_j$, there is an induced path Q_j of length d with one end v and with all other vertices in $A_j \cup V(P_j)$. Let R be the union of the paths Q_{j_1}, \dots, Q_{j_d} . Let S be the set of neighbours of v in C ; thus $S \neq \emptyset$ since every vertex of B_i has a neighbour in C . Let X be the set of vertices in C that can be joined to a vertex in S by a path of $G[C]$ of length at most $k - 1$. Thus $S \subseteq X$, and $\chi(X) \leq \tau$. By 2.6, there is a vertex $x \in X$ that is d -equipped in $C \setminus X$. Choose a path T of $G[C]$ between x and some $u \in S$, of length $k - 1$. Since x has a neighbour in C that does not belong to X , it follows that T has length $k - 1$ and no vertex of T except u belongs to S . But then the union of the paths Q_{j_1}, \dots, Q_{j_d} , the edge vu , the path T , and appropriate parts of the equipment of x , gives a (k, d) -binary star and a (k, d) -bristled star. This proves 4.2. ■

5 Building a cathedral

To apply 4.2 we need to prove that our graph contains an appropriate cathedral. First we need:

5.1 *For all $c, d, \tau \geq 0$ there exists $c' \geq 0$ with the following property. Let G be a graph with $\chi^1(G) \leq \tau$ and $\chi(G) > c'$. Then there is a spire (P, A, B) in G with height d , dominating a set C with $\chi(C) > c$.*

Proof. We may assume that $c \geq \tau$, by increasing c if necessary. Let $c' = 2c + d\tau + 1$; we claim that c' satisfies the theorem. For let G be as in the theorem, with $\chi(G) > c'$. We may assume that G is connected. Choose a vertex x_0 . Since $\chi(G) > c'$, there is a component of $G \setminus x_0$ with chromatic number at least c' , with vertex set C_1 say. By 2.3 applied to x_0 and C_1 , there is an induced path $x_0 \cdots x_d$ of G where $x_1, \dots, x_d \in C_1$, and a subset C_2 of C_1 , with the following properties:

- $x_0, \dots, x_d \notin C_2$;
- $G[C_2]$ is connected;
- x_d has a neighbour in C_2 , and x_0, \dots, x_{d-1} have no neighbours in C_2 ; and
- $\chi(C_2) \geq \chi(C_1) - d\chi^1(G) > 2c$.

Let P be the path $x_0 \cdots x_d$. For $i \geq 0$, let L_i be the set of vertices in C_2 with $G[C_2]$ -distance exactly i from x_d , and choose $i \geq 0$ such that $\chi(L_i) \geq \chi(C_2)/2 > c$. Since $c \geq \tau$ it follows that $i \geq 2$. Let $A = L_0 \cup \dots \cup L_{i-2}$, and $B = L_{i-1}$; then (P, A, B) is a spire of height d dominating L_i . This proves 5.1. ■

5.2 For all integers $d \geq 1$ and $n, c, \tau \geq 0$, there exists $c' \geq 0$ with the following property. Let G be a graph such that $\chi^2(G) \leq \tau$ and $\chi(G) > c'$. Then G admits a cathedral of length n and height d that dominates a set of chromatic number more than c .

Proof. For each integer $x \geq 0$, choose $\phi(x)$ such that 5.1 holds with c, c' replaced by $x, \phi(x)$. Let $c_n = c$, and inductively for $n - 1 \geq i \geq 0$ let $c_i = \phi(d\tau + c_{i+1})$. Let $c' = c_0$. Now let G be as in the theorem with $\chi(G) > c'$. For $1 \leq i \leq n$, choose a spire $\mathcal{S}_i = (P_i, A_i, B_i)$ and a set C_i inductively as follows. Let $C_0 = V(G)$. Suppose that for some i with $1 \leq i \leq n$ we have chosen C_{i-1} with $\chi(C_{i-1}) > c_{i-1}$. Since $c_{i-1} = \phi(d\tau + c_i)$, there is a spire (P_i, A_i, B_i) in $G[C_{i-1}]$, dominating a set $C'_i \subseteq C_{i-1}$ with $\chi(C'_i) > d\tau + c_i$. The set of vertices of C'_i with G -distance at most two from a vertex of $V(P_i) \setminus A_i$ has chromatic number at most $d\tau$, so there exists $C_i \subseteq C'_i$ with $\chi(C_i) > c_i$, such that every path in G between C_i and a vertex of $V(P_i) \setminus A_i$ has length at least three. This completes the inductive definition of $\mathcal{S}_i = (P_i, A_i, B_i)$ for $1 \leq i \leq n$. We see that:

- the spires \mathcal{S}_i ($1 \leq i \leq n$) are pairwise vertex-disjoint, and each of them dominates C_n ; and
- for $1 \leq i < j \leq n$, if $u \in V(\mathcal{S}_i)$ is adjacent to $v \in V(\mathcal{S}_j)$ then $u \in B_i$.

For $1 \leq i \leq n$, let B'_i be the set of vertices in B_i that have neighbours in C_n . Since for $1 \leq j \leq n$ every path in G between C_n and a vertex of $V(P_j) \setminus A_j$ has length at least three, it follows that for $1 \leq i < j \leq n$, no vertex in B'_i has a neighbour in $V(P_j) \setminus A_j$. Let $\mathcal{S}'_i = (P_i, A_i, B'_i)$ for $1 \leq i \leq n$; then $(\mathcal{S}'_1, \dots, \mathcal{S}'_n)$ is a cathedral in G of height d , dominating C_n , and $\chi(C_n) > c$. This proves 5.2. \blacksquare

Combining 4.2 and 5.2 we obtain:

5.3 For all $k, d \geq 1$ and $\tau \geq 0$ there exists c such that, if G is a graph with $\chi^2(G) \leq \tau$ and $\chi^k(G) \leq \tau$, and $\chi(G) > c$, then G is (k, d) -starry.

6 The case $k = 1$

It remains to prove 5.3 with the hypothesis that $\chi^2(G) \leq \tau$ omitted. (We may assume that it is not implied by the hypothesis $\chi^k(G) \leq \tau$, and so we only need to handle the case $k = 1$.) The proof of 5.2 no longer works, since in the notation of 5.2 we have no way to stop vertices in B'_i having neighbours in later paths P_j . The content of this section is our workaround.

If H is a d -superstar, we call its vertex of degree d its *root*. From repeated application of 3.2 with $r = 2$ and $k = d$ we deduce:

6.1 For all $c, d, \tau \geq 0$, there exists $c' \geq 0$ with the following property. Let G be a graph with $\chi^1(G) \leq \tau$, and let $z \in V(G)$ such that $\chi(N_G^2(z)) > c'$. Then there is a d -superstar H with root z , and a subset W of $V(G)$, such that

- $z \in W$, and no other vertex of H belongs to W or has a neighbour in $W \setminus \{z\}$; and
- $\chi(N_{G'}^2(z)) > c$, where G' is the graph $G[W]$.

Let us say a d -band in G is a triple (H, z, B) , where

- H is a d -superstar in G with root z ;
- $B \cap V(H) = \emptyset$, and z is adjacent to every vertex in B ; and
- no vertex in $H \setminus \{z\}$ has a neighbour in B .

(Thus, a band is like a spire, but with the path replaced by a subdivided star, and the set A is just a single vertex.) If $\mathcal{S} = (H, z, B)$ is a band, we write $V(\mathcal{S}) = V(H) \cup B$. A d -band (H, z, B) *dominates* C if $V(H) \cap C = \emptyset$, and B covers C , and there is no edge between $V(H)$ and C . We deduce:

6.2 *For all $d \geq 1$ and $\tau \geq 0$ there exists $c \geq 0$ with the following property. Let G be a graph with $\chi^1(G) \leq \tau$ and $\chi(G) > c$. Then G is $(1, d)$ -starry.*

Proof. Choose n_0, c' such that 4.1 is satisfied setting $k = 1$, $n = n_0$ and $c = c'$. Let $n = (2d + 1)n_0$. For each $x \geq 0$, choose $\phi(x)$ such that, if G is a graph with $\chi^2(G) \leq x$ and $\chi(G) > \phi(x)$, then G is $(1, d)$ -starry. (This is possible by 5.3.) For each $x \geq 0$, choose $\psi(x)$ such that 6.1 holds with c, c' replaced by $x, \psi(x)$.

Let $c_n = \max(c'2^{n^2}, d\tau)$, and for $n - 1 \geq i \geq 0$ let $c_i = \phi(\psi(c_{i+1}))$. Let $c = c_0$. Now let G be as in 6.2 with $\chi(G) > c$. For $1 \leq i \leq n$, choose a d -band $\mathcal{S}_i = (H_i, z_i, B_i)$ and a set C_i inductively as follows. Let $C_0 = V(G)$. Suppose that for some i with $1 \leq i \leq n$ we have chosen C_{i-1} with $\chi(C_{i-1}) > c_{i-1}$. Since $c_{i-1} \geq \phi(\psi(c_i))$, we may assume that $\chi^2(G[C_{i-1}]) > \psi(c_i)$, for otherwise the result follows from 5.3. By 6.1, there is a d -band (H_i, z_i, B_i) in $G[C_{i-1}]$, dominating a set $C'_i \subseteq C_{i-1}$ with $\chi(C'_i) > c_i$. This completes the inductive definition of $\mathcal{S}_i = (H_i, z_i, B_i)$ for $1 \leq i \leq n$.

We see that:

- the d -bands \mathcal{S}_i ($1 \leq i \leq n$) are pairwise vertex-disjoint, and each of them dominates C_n ; and
- for $1 \leq i < j \leq n$, if $u \in V(\mathcal{S}_i)$ is adjacent to $v \in V(\mathcal{S}_j)$ then $u \in B_i$.

We may assume that $G[C_n]$ is connected, and every vertex in B_i has a neighbour in C_n , for $1 \leq i \leq n$.

(1) *We may assume that there exists i with $1 \leq i \leq n$, and a vertex $v \in B_i$ that has neighbours in at least d of $V(H_{i+1}), \dots, V(H_n)$.*

Suppose not; then as in the proof of step (1) of 4.2, there exist $I \subseteq \{1, \dots, n\}$ with $|I| = n_0$, and $B'_i \subseteq B_i$ for each $i \in I$, and a subset $C' \subseteq C_n$ with $\chi(C') \geq \chi(C_n)2^{-n^2} > c'$, such that B'_i covers C' for each $i \in I$, and for $i, j \in I$ with $i < j$, there are no edges between B'_i and $V(H_j)$. For each $i \in I$, let P_i be a path of H_i of length d with one end z_i ; then $(P_i, \{z_i\}, B'_i)$ is a spire dominating C' , and the result follows from 4.1. This proves (1).

By (1), and by renumbering, we may assume that there exists $v \in B_1$ with neighbours in A_2, \dots, A_{d+1} . (We no longer need the other d -bands, so this renumbering is legitimate.) For $2 \leq i \leq d + 1$ let $y_i \in V(H_i)$ be adjacent to v . Then the union of H_1 , the edge z_1v , and the edges vy_2, \dots, vy_{d+1} forms a $(1, d)$ -binary star. Since v has a neighbour in C_n , and $\chi(C_n) > d\tau$, 2.3 implies that there is an induced path P of length d with one end v and all other vertices in C_n . But then the union of H_1 , the edges z_1v and vy_2 , and the path P , forms a $(1, d)$ -bristled star. This proves 6.2. ■

Let us put these pieces together to deduce 1.6.

Proof of 1.6. Let $k, d \geq 1$, and $\kappa \geq 0$; we need to prove that for all $\kappa \geq 0$, there exists c such that every graph G with $\omega(G) \leq \kappa$ and $\chi(G) > c$ is (k, d) -starry. We prove this by induction on κ . Let τ_1 be such that every graph G with $\omega(G) \leq \kappa - 1$ and $\chi(G) > \tau_1$ is (k, d) -starry. By 3.3 with $r = k$, there exists $\tau \geq \tau_1$ such that every graph G with $\chi^1(G) \leq \tau_1$ and $\chi^k(G) > \tau$ is (k, d) -starry. Choose c_1 such that 5.3 holds with c replaced by c_1 . Choose c_2 such that 6.2 holds with τ, c replaced by τ_1, c_2 . Let $c = \max(c_1, c_2)$, and let G be a graph with $\chi(G) > c$ and $\omega(G) \leq \kappa$. We may assume that $\chi^1(G) \leq \tau_1$, for otherwise the result follows from the induction on $\omega(G)$. We may assume that $\chi(G^k) \leq \tau$ for otherwise the result follows from 3.3. By 5.3 we may assume that $\chi^2(G) > \tau$ and so $k = 1$ because $\chi(G^k) \leq \tau$. But then the result follows from 6.2. This proves 1.6. \blacksquare

7 Two counterexamples

As we said in the beginning, our main tools are the lemmas of section 2. We proved there in particular that if G is a graph of very large chromatic number and with $\chi^1(G)$ bounded, and v is a vertex such that $\chi(G \setminus v) < \chi(G)$, then there is an induced d -star in G with centre v , and there is an induced $(d + 1)$ -edge path in G with second vertex v . What other trees containing v must be present? If we could find more, then the methods of this paper might allow us to prove the Gyárfás-Sumner conjecture for more types of trees. But there are not many more. For instance, Sophie Spirkl and the third author showed that there need not be an induced five-vertex path in G with middle vertex v . Here is the example.

Choose a large integer k , and take a minimal triangle-free graph with chromatic number more than k . Let I be the set of neighbours of some vertex u , and delete u . This produces a graph H say, and a stable subset I of $V(H)$, such that H is triangle-free, and k -colourable, but in every k -colouring all k colours occur in I .

For each subset $S \subseteq I$ with $|S| = k - 1$, make a gadget B^S as follows. Take some enumeration $\{s_1, \dots, s_{k-1}\}$ of S , take $2k$ new vertices $a_1^S, b_1^S, a_2^S, b_2^S, \dots, a_k^S, b_k^S$, and for $1 \leq i \leq k$ and $i \leq j \leq k - 1$ make a_i^S and b_i^S both adjacent to s_j . Also for $1 \leq i < j \leq k - 1$, make a_i^S adjacent to b_j^S and b_i^S adjacent to a_j^S . It is easy to see that every k -colouring of S can be extended to a k -colouring of the new vertices. Now add one more new vertex v^S adjacent to all of $a_1^S, b_1^S, \dots, a_k^S, b_k^S$. This defines B^S (that is, the set $\{a_1^S, b_1^S, \dots, a_k^S, b_k^S, v^S\}$, and the new edges incident with these new vertices). Let G_1 be the graph obtained from H by adding B^S for every choice of S . Now G_1 is not k -colourable; because in a k -colouring of H , some choice of S is coloured with all different colours, and this cannot be extended to a k -colouring of G_1 . Starting with H , let us add the gadgets B^S one by one until the chromatic number increases to $k + 1$, and then stop; let G be the graph just constructed and B^S the final gadget added. Let $v = v^S$; then $\chi(G \setminus v) < \chi(G)$, and there is no five-vertex induced path of G with middle vertex v .

One might also hope that the parts of the equipment of a vertex can be unified; say a vertex v is “properly d -equipped” in Y if there is an induced path of length d with first vertex v and all other vertices in Y , and d pairwise nonadjacent neighbours of v , all in Y , and such that none of them has any neighbours in $P \setminus v$. One might hope that 2.5 could be strengthened correspondingly. But this is false, even for $d = 2$ and for triangle-free graphs, as another counterexample (also due to Sophie

Spirkl and the third author) shows. Take H and I as before; and for each $S \subseteq I$ with $|S| = k$, let B^S be a gadget defined as follows. Let $S = \{s_1, \dots, s_k\}$, and take $k + 1$ new vertices a_1^S, \dots, a_k^S and v^S , and for all distinct i, j with $1 \leq i, j \leq k$, make a_i^S adjacent to s_j . Also make v^S adjacent to a_1^S, \dots, a_k^S . Adding all these gadgets increases the chromatic number, so, as before, add them one at a time until the chromatic number increases, let B^S be the last one, and let $v = v^S$. Then v is not properly 2-equipped in $V(G) \setminus \{v\}$.

References

- [1] M. Chudnovsky, A. Scott, P. Seymour and S. Spirkl, “Induced subgraphs of graphs with large chromatic number. VIII. Long odd holes”, submitted for publication, [arXiv:1701.07217](https://arxiv.org/abs/1701.07217).
- [2] A. Gyárfás, “On Ramsey covering-numbers”, *Coll. Math. Soc. János Bolyai*, in *Infinite and Finite Sets*, North Holland/American Elsevier, New York (1975), 10.
- [3] A. Gyárfás, “Problems from the world surrounding perfect graphs”, *Proceedings of the International Conference on Combinatorial Analysis and its Applications*, (Pokrzywna, 1985), *Zastos. Mat.* 19 (1987), 413–441.
- [4] A. Gyárfás, E. Szemerédi and Zs. Tuza, “Induced subtrees in graphs of large chromatic number”, *Discrete Math.* 30 (1980), 235–344.
- [5] H.A. Kierstead and S.G. Penrice, “Radius two trees specify χ -bounded classes”, *J. Graph Theory* 18 (1994), 119–129.
- [6] H.A. Kierstead and Y. Zhu, “Radius three trees in graphs with large chromatic number”, *SIAM J. Discrete Math.* 17 (2004), 571–581.
- [7] A.D. Scott, “Induced trees in graphs of large chromatic number”, *J. Graph Theory* 24 (1997), 297–311.
- [8] A. Scott and P. Seymour, “Induced subgraphs of graphs with large chromatic number. XIII. New brooms”, submitted for publication, [arXiv:1807.03768](https://arxiv.org/abs/1807.03768).
- [9] D.P. Sumner, “Subtrees of a graph and chromatic number”, in *The Theory and Applications of Graphs*, (G. Chartrand, ed.), John Wiley & Sons, New York (1981), 557–576.