

Errata for First Edition, 2003

We are grateful to Nick Higham (University of Manchester), Gisbert Stoyan (University of Budapest), Gerald Moore (Imperial College London), Goetz Alefeld and Jan Mayer (University of Karlsruhe), and Shahrokh Esmaili (University of Kurdistan) for helpful suggestions.

- p.11, Theorem 1.5, line 3: $\xi = g(\xi) \in [a, b] \longrightarrow \xi = g(\xi) \in (a, b)$
 p.55, in the first matrix on the r.h.s. of (2.25): $P^* \longrightarrow (P^*)^T$. Line below (2.25): $P^*P^* = I \longrightarrow (P^*)^T P^* = I$
 p.79, lines -7 and -8: $\|\mathbf{b}_r\| \longrightarrow \|\mathbf{b}_r\|_2$
 p.80, line -5: *Goldstein* should read *Goldstine*, and footnote 4 should read:
⁴ Herman Heine Goldstine (1913–) collaborated with John von Neumann on the development of the ENIAC (Electronic Numerical Integrator And Computer), the first electronic digital computer, which became operational in 1945.
 p.100, Theorem 3.6, line 1: insert *nonsingular* before *tridiagonal*
 p.103, Exercise 3.5, line 3: insert *with strict inequality for $j = n$* before *and*
 p.115, line -2: $D = [0, 1] \times [0, 3/5] \longrightarrow D = [0.5, 1] \times [0, 0.7]$
 p.135, footnote 1: $\lambda_{1/2} \longrightarrow \lambda_{1,2}$
 p.178, line 3, insert the missing term $-\lambda^{(0)}\delta\mathbf{x}$ on l.h.s.
 p.255, line 5: *form* \longrightarrow *from*
 p.264, line 10: insert *except for $m = n = 0$ when $\langle T_m, T_n \rangle = \pi$. after nonnegative integers m and n*
 p.298 (11.4): in the definition of $\varphi_0(x)$ replace h_0 by h_1
 p.313 (12.9), last sum in the chain: $j + 1 \longrightarrow j$ top and bottom
 p.331, Adams–Moulton method: $-9f_n \longrightarrow +f_n$
 p.334, line above (12.41): $r = 1, 2, \dots, n \longrightarrow r = 1, 2, \dots, k$
 p.335, line 4:
$$D_1 = \prod_{1 \leq r < s \leq \ell} (z_s - z_r)^{m_r \cdot m_s} \prod_{r=1}^{\ell} (m_r - 1)!! z_r^{m_r(m_r-1)/2}$$

 p.336, Example 12.5, line -6: $z_3 = -3.14 \longrightarrow z_3 \approx -3.14$
 p.365, Theorem 13.3, line 1: $j = 0, 1, \dots, n, \longrightarrow j = 1, \dots, n - 1,$
 p.369, insert after line 2: where $M_3 := \max_{x \in [a-h, b]} |y'''(x)|$ and $M_4 := \max_{x \in [a-h, b]} |y^{iv}(x)|$.
 p.377, line 12: $u_3(x, y) \longrightarrow u_3(x, t)$.
 p.378, lines 17–20: However, in the interval $(a, d]$, $\eta(x; t) > 0$ and $\partial f / \partial y \geq 0$, so that $\eta''(x; t) \geq 0$. Consequently, in the interval $[a, d]$, $\eta'(x; t) \geq \eta'(a; t) > 0$, and we have a contradiction. Thus $\eta(x; t) > 0$ for $a < x \leq b$. It then follows that $\eta''(x; t) \geq 0$, and hence also $\eta'(x; t) > 0$ on ...
 p.381, line -5:

$$-\frac{\delta^2 y_j}{h^2} + \beta_{-1} f_{j-1} y_{j-1} + \beta_0 f_j y_j + \beta_1 f_{j+1} y_{j+1} = \beta_{-1} g_{j-1} + \beta_0 g_j + \beta_1 g_{j+1}$$

- p.407, (14.37): $L^2(0, 1) \longrightarrow L^2(a, b)$
 p.410, line 3: $L^2(x_{i-1}^m, x_i^m) \longrightarrow L^2(x_{i-1}^{(m)}, x_i^{(m)})$

Errata for Second Printing, 2006

We are grateful to Christoph Ortner (University of Warwick), Niall Madden (National University of Ireland, Galway), Hakan Tor (Middle East Technical University, Ankara), Joseph F. Grcar (Castro Valley, California), Bart Vandereycken (Princeton University), Sandra Conor (University of Malaya), Nicolas Boumal (Princeton University), Taku Kaneda (Catholic University of Louvain), Shahrokh Esmaili (University of Kurdistan), and Michael Zellinger (University of Oxford) for helpful suggestions.

- p.ix line 3: copies of L^AT_EX files \longrightarrow the PDF file
 p.44, lines 1–5 of Section 2.2, and p.80 lines 13–18: Contrary to what is stated on pages 44 and 80, Gaussian elimination was not first published by Gauss in 1809, nor was the method extended to linear systems with general matrices by Jacobi in a posthumous publication from 1857. By the time Gauss was born, the elimination method was already an established lesson in algebra textbooks such as those of Bezout and Simpson. The details of the actual history of Gaussian elimination can be found in the article by Grcar, J.F., *How ordinary elimination became Gaussian elimination*. *Historia Mathematica* (2010), doi:10.1016/j.hm.2010.06.003.
 p.95, eq. (3.12): $PA = L^{(1)}U^{(1)}$ should read $PT = L^{(1)}U^{(1)}$
 p.151, Fig. 5.2: $H\mathbf{x}$ below the horizontal plane should be $H\mathbf{x}$
 p.159: 4 lines above Example 5.7: $p_{j-1}(\theta) \longrightarrow -p_{j-1}(\theta)$. The wording in Example 5.7 in lines 1–10 on p.160 should be modified accordingly, though the results remain the same.
 p.164, line below (5.35): Delete the sentence: “In fact, since A is tridiagonal ...”
 p.177, Exercise 5.7, line 3: Delete the words: “and tridiagonal”
 p.205, Eq. (7.7): $\frac{(b-a)^4}{196} M_3 \longrightarrow \frac{(b-a)^4}{192} M_3$
 p.281, last line should read: $A_0(\frac{1}{2} + \sqrt{\frac{1}{12}}) + A_1(\frac{1}{2} - \sqrt{\frac{1}{12}}) = \frac{1}{2}$
 p.288, Eq. (10.27): $x_k \longrightarrow x_k^*$
 p.299, lines -11 and -10: Theorem 3.4 \longrightarrow Exercise 3.5
 p.320, in Example 12.3, (12.20): $y' = y^2 + g(x) \longrightarrow y' = y^2 - g(x)$
 p.352: the last line of the second displayed Butcher tableax on the page should be $\frac{1}{6} \quad \frac{1}{3} \quad \frac{1}{3} \quad \frac{1}{6}$
 p.371: Lemma 3.1: $\frac{1}{24}h^4 \longrightarrow \frac{1}{24}h^2$
 p.382, line 2: $T_j = (\beta_{-1} + \beta_0 + \beta_1 - 1)y''(x_j) + Z_j^0 h;$ line 12: $\frac{1}{60480} M_8 \longrightarrow \frac{17}{60480} M_8$