

Multilevel Function Approximation

Filippo De Angelis, Mike Giles, Christoph Reisinger

Mathematical Institute, University of Oxford

Oden seminar

March 13, 2025

Objective

Want to construct an approximation for a scalar function

$$f : [0, 1]^d \rightarrow \mathbb{R}$$

with parametric dimension d in the range 1 – 8, where $f(\theta)$ is one of the following:

- a functional of the solution $u(\theta; x)$ of a PDE, with θ dependence in the PDE coefficients, the boundary data and/or the functional
- a parametric expectation $\mathbb{E}_\omega[g(\theta; \omega)]$, where $g(\theta; \omega)$ is a functional of the solution of an SDE

Problem: in either case we must approximate $f(\theta)$, and the more accurate the approximation, the greater the computational cost.

Objective: for given ε , lowest cost approximation \tilde{f} with

$$\|\tilde{f} - f\| < \varepsilon$$

Outline

- quick recap of key literature:
 - ▶ dense grid linear interpolation
 - ▶ sparse grid linear interpolation
 - ▶ convergence of PDE approximations
 - ▶ MLMC for SDEs, MIMC for SPDEs
 - ▶ MLMC for parametric integration (Heinrich)
- MLFA for PDEs
 - ▶ idea
 - ▶ dense grid linear interpolation
 - ▶ sparse grid linear interpolation
- MLFA for SDEs – extension of Heinrich's approach
 - ▶ randomised MLMC for SDE
 - ▶ randomised MLMC and sparse grids
 - ▶ MLMC decomposition for SDE
 - ▶ MLMC decomposition and sparse grids
- conclusions and references

Dense grid linear interpolation

For a 1-dimensional function, $f : [0, 1] \rightarrow \mathbb{R}$, if we use a uniform grid $\theta_j = j2^{-\ell}$, $j = 0, 1, \dots, 2^\ell$, then the piecewise linear interpolation of the values $f(\theta_j)$ has an error bound of the form

$$\|\tilde{f} - f\| < c(f)2^{-r\ell}$$

if $f \in C^r([0, 1])$ for $r \in \{1, 2\}$.

Using a tensor product grid in higher dimension d , this generalises to

$$\|\tilde{f} - f\| < c(f)2^{-r\ell}$$

if $f \in C^r([0, 1]^d)$, but now the number of evaluation points is $O(2^{d\ell})$ so the expense is much greater

Sparse grid linear interpolation

To avoid that “curse of dimensionality” as d increases, can instead use a Smolyak sparse grid interpolation based on piecewise multi-linear functions in each direction.

This has an error bound of the form

$$\|\tilde{f} - f\| < c(f) 2^{-r\ell} (\ell+1)^{d-1}$$



with the number of interpolation points being $O(2^\ell (\ell+1)^{d-1})$.

However, it needs more regularity in f , including mixed derivatives of degree up to r in each direction:

$$\frac{\partial^{\alpha_1 + \alpha_2 + \dots} f}{\partial \theta_1^{\alpha_1} \partial \theta_2^{\alpha_2} \dots}, \quad 0 \leq \alpha_j \leq r \leq 2.$$

Much better than dense grid interpolation for modest values of d , up to 8?

Dense versus sparse grid interpolation



Convergence of PDEs

If $f_h(\theta)$ is the functional which comes from the approximate solution of a PDE using a discretisation with spacing h , and input θ , then typically

$$\|f_h - f\| = O(h^q)$$

for some q , and the cost of evaluating $f_h(\theta)$ is $O(h^{-p})$ for some p .

Often, but not always, the θ derivatives of f_h will have the same rate of convergence.

MLMC for SDEs

When estimating $\mathbb{E}[P]$, with P a functional of the solution of an SDE, MLMC is based on the telescoping sum

$$\mathbb{E}[\widehat{P}_L] = \sum_{\ell=0}^L \mathbb{E}[\Delta \widehat{P}_\ell], \quad \Delta \widehat{P}_\ell \equiv \widehat{P}_\ell - \widehat{P}_{\ell-1}, \quad \widehat{P}_{-1} \equiv 0$$

where \widehat{P}_ℓ represents an approximation to output P on level ℓ using timestep $h_\ell = 2^{-\gamma\ell} h_0$. If there are also constants α, β such that


$$\mathbb{E}[\widehat{P}_\ell - P] = O(2^{-\alpha\ell}), \quad \mathbb{V}[\Delta \widehat{P}_\ell] = O(2^{-\beta\ell})$$

then the MLMC method chooses a near-optimal number of levels L , and number of samples $M_\ell, \ell = 0, 1, \dots, L$ to obtain a r.m.s. accuracy of ε at a cost of order

$$\begin{aligned} \varepsilon^{-2}, & \quad \beta > \gamma \\ \varepsilon^{-2} |\log \varepsilon|^2, & \quad \beta = \gamma \\ \varepsilon^{-2 - (\gamma - \beta)/\alpha}, & \quad \beta < \gamma \end{aligned}$$

MIMC for SPDEs

Haji-Ali, Nobile & Tempone (2015) developed an important extension, MIMC (Multi-Index Monte Carlo), incorporating sparse grid ideas to separately refine multiple parameters.



$$\mathbb{E}[P] \approx \sum_{l_1+l_2 \leq l} \mathbb{E}[\Delta_{l_1} \Delta_{l_2} \hat{P}_l]$$

This increases the range of applications with $O(\varepsilon^{-2})$ complexity.

Randomised MLMC for SDEs

In another important extension, in the “good” MLMC case, $\beta > \gamma$, Rhee & Glynn (2015) developed the randomised MLMC estimator

$$p_L^{-1} \Delta \hat{P}_L$$

where L is a random level, with $L = \ell$ with probability $p_\ell \propto 2^{-(\beta+\gamma)\ell/2}$.

Since

$$\begin{aligned} \mathbb{E}[p_L^{-1} \Delta \hat{P}_L] &= \sum_{\ell=0}^{\infty} \mathbb{P}[L = \ell] \mathbb{E}[p_L^{-1} \Delta \hat{P}_L \mid L = \ell] \\ &= \sum_{\ell=0}^{\infty} p_\ell \mathbb{E}[p_\ell^{-1} \Delta \hat{P}_\ell] = \sum_{\ell=0}^{\infty} \mathbb{E}[\Delta \hat{P}_\ell] = \mathbb{E}[P] \end{aligned}$$

it is an unbiased estimator, and it can be proved that the variance and expected cost are both finite if $\beta > \gamma$.


MLMC for parametric integration

Stefan Heinrich's MLMC research (2001) concerned the approximation of $f(\theta) = \mathbb{E}[g(\theta; \omega)]$, given exact sampling of $g(\theta; \omega)$ at unit cost.

In his formulation, the MLMC telescoping sum is

$$f \approx I_L[f] = I_0[f] + \sum_{\ell=1}^L I_\ell[f] - I_{\ell-1}[f]$$

where $I_\ell[f]$ represents a level ℓ interpolation.



MLMC for parametric integration

Heinrich then approximates $(I_\ell - I_{\ell-1})[f]$ through Monte Carlo sampling at required values of θ :

$$(I_\ell - I_{\ell-1})[f] \approx \frac{1}{M_\ell} \sum_{m=1}^{M_\ell} (I_\ell - I_{\ell-1})[g(\cdot; \omega^{\ell, m})]$$

As $\ell \rightarrow \infty$, $(I_\ell - I_{\ell-1})[f] \rightarrow 0$ and $\mathbb{V}[(I_\ell - I_{\ell-1})[g]] \rightarrow 0$, so fewer MC samples needed on finer levels.

Analysis assumes the number of θ points increases exponentially with dimension (as with dense tensor product grid), so the resulting complexity for linear interpolation is of order

$$\begin{aligned} \varepsilon^{-2}, & \quad d < 2r \\ \varepsilon^{-2} |\log \varepsilon|^2, & \quad d = 2r \\ \varepsilon^{-d/r}, & \quad d > 2r \end{aligned}$$

assuming $g(\theta; \omega)$ is sufficiently smooth w.r.t. θ

MLMC for parametric integration

Heinrich's work is the key starting point for our current work which extends it in several directions:

- PDEs with appropriate numerical approximation
- sparse grid interpolation to address curse of dimensionality
- weaker assumptions on smoothness of $g(\theta; \omega)$
- numerical approximation of $f(\theta) \equiv \mathbb{E}[g(\theta; \omega)]$ in cases without a finite variance, finite expected cost unbiased estimator

The PDE / sparse grid interpolation combination has been pioneered by Teckentrup *et al* (2015) for elliptic PDEs in stochastic collocation (parameter space is uncertainty space).

There is also an excellent review article by Tempone and Wolfers (2018).

MLFA for PDEs

The fundamental idea is very simple: building on Stefan Heinrich's approach, if the function f has an interpolation expansion

$$f = I_0[f] + \sum_{\ell=1}^{\infty} I_{\ell}[f] - I_{\ell-1}[f] = \sum_{\ell=0}^{\infty} \Delta I_{\ell}[f]$$

with $\Delta I_{\ell} \equiv I_{\ell} - I_{\ell-1}$, $I_{-1} \equiv 0$, and as $\ell \rightarrow \infty$, $\Delta I_{\ell}[f] \rightarrow 0$ since the number of interpolation points increases, then we will use an approximation

$$\tilde{f} = \sum_{\ell=0}^L \Delta I_{\ell}[f_{\ell}]$$

where f_{ℓ} is based on a PDE approximation with grid spacing h_{ℓ} and

- h_{ℓ} is small for small ℓ – a few expensive accurate PDE calculations
- h_{ℓ} is large for large ℓ – lots of cheap PDE calculations

MLFA for PDEs

It follows from the triangle inequality that

$$\|\tilde{f} - f\| \leq \|(I_L - I)[f]\| + \sum_{\ell=0}^L \|(I_\ell - I_{\ell-1})[f_\ell - f]\|.$$

If we assume second order accuracy in the interpolation so that

$$\|(I_L - I)[f]\| < c_1 2^{-2L}, \quad \|\Delta I_\ell[f_\ell - f]\| < c_2 2^{-2\ell} h_\ell^q$$

and the cost C_ℓ of constructing $(I_\ell - I_{\ell-1})[f_\ell]$ on level ℓ is bounded by

$$C_\ell < c_3 2^{d\ell} h_\ell^{-p}$$

then to achieve an accuracy of ε we can choose L s.t.

$$c_1 2^{-2L} \approx \varepsilon/2 \quad \implies \quad L = O(|\log \varepsilon|)$$

and ...

MLFA for PDEs

... choose h_ℓ to minimise

$$c_3 \sum_{\ell=0}^L 2^{d\ell} h_\ell^{-p}$$

subject to the requirement that

$$c_2 \sum_{\ell=0}^L 2^{-2\ell} h_\ell^q \approx \varepsilon/2.$$

Using a Lagrange multiplier gives the optimal h_ℓ as

$$h_\ell = 2^{(d+2)\ell/(p+q)} h_0$$

The accuracy requirement then becomes

$$c_2 h_0^q \sum_{\ell=0}^L 2^{-\nu\ell} \approx \varepsilon/2, \quad \nu \equiv (2p-dq)/(d+2)$$

MLFA for PDEs

$\nu > 0$ leads to $h_0 = O(\varepsilon^{1/q})$ and a total cost of $O(\varepsilon^{-p/q})$,

$\nu = 0$ leads to $h_0 = O(\varepsilon^{-1/q}L^{1/q})$ and a cost of $O(\varepsilon^{-p/q}|\log \varepsilon|^{1+p/q})$.

$\nu < 0$ leads to $h_0 = O(\varepsilon^{-1/q}2^{\nu L/q})$ and a cost of $O(\varepsilon^{-d/2})$.

Thus the total cost is of order

$$\begin{aligned} \varepsilon^{-p/q}, & \quad p/q > d/2 \\ \varepsilon^{-p/q} |\log \varepsilon|^{1+p/q}, & \quad p/q = d/2 \\ \varepsilon^{-d/2}, & \quad p/q < d/2 \end{aligned}$$

Note:

- $O(\varepsilon^{-p/q})$ is the cost of a single ε -accurate PDE calculation
- $O(\varepsilon^{-d/2})$ is the cost of an ε -accurate interpolation of unit cost data

In this sense the method has near-optimal asymptotic efficiency

MLFA for PDEs with sparse interpolation

With sparse interpolation the accuracy requirement becomes

$$c_2 \sum_{\ell=0}^L 2^{-2\ell} (\ell+1)^{d-1} h_\ell^q \approx \varepsilon/2.$$

and the cost bound becomes

$$C = c_3 \sum_{\ell=0}^L 2^\ell (\ell+1)^{d-1} h_\ell^{-p}$$

Optimising this results in the total cost being of order

$$\begin{aligned} \varepsilon^{-p/q}, & \quad p/q > 1/2 \\ \varepsilon^{-p/q} |\log \varepsilon|^{3d/2}, & \quad p/q = 1/2 \\ \varepsilon^{-1/2} |\log \varepsilon|^{3(d-1)/2}, & \quad p/q < 1/2 \end{aligned}$$

Note:

- $O(\varepsilon^{-p/q})$ is again the cost of a single ε -accurate PDE calculation
- $O(\varepsilon^{-1/2} |\log \varepsilon|^{3(d-1)/2})$ is the cost of an ε -accurate sparse interpolation of unit cost data

MLFA for PDEs

What's the catch? What can go wrong?

The method relies on the convergence of derivatives of the approximation


- second derivatives for regular grid when using linear interpolation
- higher degree mixed derivatives when using sparse grids
(up to second order in each direction)

Sometimes, can lose one order of convergence for each derivative.

MLFA for PDEs

A trivial example from mathematical finance: suppose we have “initial” boundary data $u(x) = H(x - K)$ where $H(\cdot)$ is the Heaviside step function.

Using discrete initial data $U_j = u(x_j)$ then as K varies the solution at a later time has the following form, so $O(\Delta x)$ convergence for solution, $O(1)$ error for derivative




MLFA for PDEs

It doesn't even have to be discontinuous to be a problem. If

$$u(K) = 1 - 0.2 K, \quad U(K) = 1 - 0.2 K + \Delta x \sin(K/\Delta x)$$

where Δx is the mesh size, then again

$$\|u - U\| = O(\Delta x), \quad \|u' - U'\| = O(1)$$




MLFA for PDEs

A solution to this kind of problem can be to use a finite element projection

e.g. define discrete initial data U_j as

$$U_j = \int_{x_{j-1}}^{x_{j+1}} \frac{1}{\Delta x^2} \min(x - x_{j-1}, x_{j+1} - x) u(x) dx$$



MLFA for SDEs

If $\beta > \gamma$ in the standard SDE discretisation sense, then randomised MLMC can be used to give an unbiased estimator Y , with $\mathbb{E}[Y(\theta; \omega)] = f(\theta)$ and finite variance and expected cost. If

$$\begin{aligned}\|(I_\ell - I)[f]\| &< c_1 2^{-r\ell} \\ \mathbb{V}[(I_\ell - I_{\ell-1})[Y]] &< c_2 2^{-s\ell}\end{aligned}$$

and the total expected cost is bounded by $c_3 \sum_0^L 2^{d\ell} M_\ell$, for M_ℓ samples per level, then ε r.m.s. accuracy can be achieved with cost of order

$$\varepsilon^{-2}, \quad s > d$$

$$\varepsilon^{-2} |\log \varepsilon|^2, \quad s = d$$

$$\varepsilon^{-2-(d-s)/r}, \quad s < d$$

MLFA for SDEs

The previous result is a slight generalisation of Heinrich's analysis which assumed $s = 2r$.

With sparse interpolation, the cost is reduced to order

$$\varepsilon^{-2}, \quad s > 1$$

$$\varepsilon^{-2} |\log \varepsilon|^{2+3(d-1)}, \quad s = 1$$

$$\varepsilon^{-2-(1-s)/r} |\log \varepsilon|^{(3+(1-s)/r)(d-1)}, \quad s < 1$$

MLFA for SDEs

If $\beta \leq \gamma$, then we can use a MIMC combination of path-based MLMC and Heinrich's MLMC. The starting point is the interpolation decomposition:

$$f \approx \sum_{\ell=0}^L (I_{\ell} - I_{\ell-1})[f], \quad I_{-1}[f] \equiv 0,$$

where I_{ℓ} uses a dense interpolation with spacing proportional to $2^{-\ell}$.

We then replace f with a timestep approximation expansion

$$f \approx \sum_{\ell=0}^L \sum_{\ell'=0}^{L'_{\ell}} \Delta I_{\ell}[\Delta f_{\ell'}], \quad \Delta I_{\ell}[\Delta f_{\ell'}] \equiv (I_{\ell} - I_{\ell-1})[f_{\ell'} - f_{\ell'-1}]$$

in which L'_{ℓ} is a decreasing function of ℓ , since $\Delta I_{\ell}[f]$ becomes smaller as ℓ increases and so less relative accuracy is required in its approximation.

MLFA for SDEs

The final step is to replace $\Delta I_\ell[\Delta f_{\ell'}]$ by a Monte Carlo estimate, giving the MIMC-style estimator

$$\tilde{f} = \sum_{\ell=0}^L \sum_{\ell'=0}^{L'_\ell} \left(\frac{1}{M_{\ell,\ell'}} \sum_{m=1}^{M_{\ell,\ell'}} \Delta I_\ell[\Delta g_{\ell'}(\cdot; \omega^{\ell,\ell',m})] \right)$$

We now need to choose $L, L'_\ell, M_{\ell,\ell'}$ to achieve the desired accuracy at the minimum cost.

$$\begin{aligned} \mathbb{E}[\tilde{f} - f] &= (I_L - I)[f] + \sum_{\ell=0}^L (I_\ell - I_{\ell-1})[f_{L'(\ell)} - f] \\ \implies \left\| \mathbb{E}[\tilde{f} - f] \right\| &\leq \|(I_L - I)[f]\| + \sum_{\ell=0}^L \|(I_\ell - I_{\ell-1})[f_{L'(\ell)} - f]\| \end{aligned}$$

and

$$\mathbb{V}[\tilde{f}] = \sum_{\ell=0}^L \sum_{\ell'=0}^{L'_\ell} \left(\frac{1}{M_{\ell,\ell'}} \mathbb{V}[\Delta I_\ell[\Delta g_{\ell'}(\cdot; \omega)]] \right)$$

MLFA for SDEs

If we have

$$\|\Delta I_\ell[\Delta f_{\ell'}]\| < c_1 2^{-r\ell - \alpha\ell'}$$

$$\mathbb{V}[\Delta I_\ell[\Delta g_{\ell'}]] < c_2 2^{-s\ell - \beta\ell'}$$

and the total cost is bounded by

$$c_3 \sum_{\ell=0}^L \sum_{\ell'=0}^{L'} 2^{d\ell + \gamma\ell'} M_{\ell,\ell'},$$

then ε RMS accuracy can be achieved at a computational cost of order

$$\varepsilon^{-2}, \quad \eta < 0$$

$$\varepsilon^{-2-\eta} |\log \varepsilon|^p, \quad \eta \geq 0$$

for some p (see MIMC analysis by [HNT16]), where

$$\eta = \max\left(\frac{\gamma - \beta}{\alpha}, \frac{d - s}{r}\right).$$

MLFA for SDEs

Note: in the best case when $\eta < 0$, the dominant contribution to the total cost comes from the base level $\ell = \ell' = 0$, which is why there are no log terms in its complexity.

With sparse interpolation the corresponding cost is of order

$$\begin{aligned} \varepsilon^{-2}, & \quad \eta < 0 \\ \varepsilon^{-2-\eta} |\log \varepsilon|^q, & \quad \eta \geq 0 \end{aligned}$$

for some q , where now

$$\eta = \max \left(\frac{\gamma - \beta}{\alpha}, \frac{1 - s}{r} \right).$$

Conclusions and future work

Conclusions:

- excellent asymptotic efficiency in approximating parametric functions arising from PDEs and SDEs – nearly optimal in some cases
- meta-theorems make various assumptions which need to be verified, especially for mixed derivatives when using sparse grid interpolation

On-going work:

- numerical results for PDEs and SDEs
- writing up numerical analysis for PDEs, to prove validity of mixed derivative assumptions for parabolic PDEs with non-smooth initial data, based on Carter, G (2007)
- writing up numerical analysis for SDEs, based on earlier work, G, Sheridan-Methven (2022) – in some cases the variance doesn't have the simple product form but the complexity can still be derived

References

- H.-J. Bungartz, M. Griebel. 'Sparse grids'. pp.147-269, Acta Numerica, 2004.
- R. Carter, M.B. Giles. 'Sharp error estimates for discretizations of the 1D convection–diffusion equation with Dirac initial data'. IMA Journal of Numerical Analysis, 27(2):406-425, 2007.
- F. De Angelis. PhD thesis, in preparation, Oxford University, 2025.
- M.B. Giles, O. Sheridan-Methven, 'Analysis of nested multilevel Monte Carlo using approximate Normal random variables'. SIAM/ASA Journal on Uncertainty Quantification 10(1):200-226, 2022.
- A.-L. Haji-Ali, F. Nobile, R. Tempone. 'Multi-index Monte Carlo: when sparsity meets sampling'. Numerische Mathematik, 132:767-806, 2016.
- S. Heinrich. 'Multilevel Monte Carlo methods'. Lecture Notes in Computer Science, 2179:58-67, 2001.
- C.-H. Rhee, P.W. Glynn. 'Unbiased estimation with square root convergence for SDE models'. Operations Research, 63(5):1026-1043, 2015.
- A.L. Teckentrup, P. Jantsch, C.G. Webster, M. Gunzburger. 'A Multilevel Stochastic Collocation Method for Partial Differential Equations with Random Input Data', SIAM/ASA Journal on Uncertainty Quantification, 3(1):1046-1074, 2015.
- R. Tempone, S. Wolfers. 'Smolyak's algorithm: a powerful black box for the acceleration of scientific computations'. pp.201-228 in Sparse Grids and Applications, Springer, 2018.