

A LOGARITHMIC BOUND FOR THE CHROMATIC NUMBER OF THE ASSOCIAHEDRON

LOUIGI ADDARIO-BERRY, BRUCE REED, ALEX SCOTT, AND DAVID R. WOOD

ABSTRACT. We show that the chromatic number of the n -dimensional associahedron grows at most logarithmically with n , improving a bound from and proving a conjecture of Fabila-Monroy et al. (2009).

1. INTRODUCTION

The associahedron \mathcal{A}_n is an $(n - 3)$ -dimensional convex polytope that arises in numerous branches of mathematics, including algebraic combinatorics [6, 12, 16, 29] and discrete geometry [2, 22, 23]. Associahedra are also called Stasheff polytopes after the work of Stasheff [29], following earlier work by Tamari [30]. We are only interested in the 1-skeleton of the associahedron, so we consider it as a graph, defined as follows.

The elements of the associahedron \mathcal{A}_n are triangulations T of the convex n -gon with vertices labeled by $\{0, \dots, n - 1\}$ in clockwise order. For any such triangulation T , we always denote triangles of T by the sequence ABC of their vertices, ordered so that $A < B < C$. We write $E(T)$ for the set of edges contained in T . Every triangulation T of \mathcal{A}_n contains the edges $01, 12, \dots, (n - 1)0$; we refer to these as *boundary edges*. For T in \mathcal{A}_n , each non-boundary edge $e \in E(T)$ is contained in a unique quadrilateral $Q = Q_T(e) = ABCD$ with $A < B < C < D$; we always list the vertices of quadrilaterals in increasing order. *Flipping* the edge e means replacing e by the other diagonal of Q ; see Figure 1. Two triangulations T, T' in \mathcal{A}_n are adjacent in \mathcal{A}_n if they may be obtained from one another by a single flip.

Graph-theoretic properties of associahedra have been well-studied. For example, it is easily seen that \mathcal{A}_n is $(n - 3)$ -regular. Lucas [17] showed that \mathcal{A}_n is Hamiltonian. Lee [15] determined the automorphism group of \mathcal{A}_n . Hurtado and Noy [13] showed that \mathcal{A}_n has connectivity $n - 3$, as well as giving simple proofs of the above results concerning the Hamiltonicity and the automorphism group of \mathcal{A}_n . Parlier and Petri [21] proved bounds on the genus of \mathcal{A}_n . The diameter of \mathcal{A}_n and several related questions have been studied extensively [1, 3–5, 7, 9, 18, 24–27]. Sleator et al. [27] proved that the diameter equals $2n - 10$ for sufficiently

Date: November 21, 2018; revised December 22, 2025.


2020 *Mathematics Subject Classification.* 52B05, 05C15.

Louigi Addario-Berry is supported by an NSERC Discovery Grant and by the Canada Research Chairs program.

Bruce Reed is supported by NSTC Grant 112-2115-M-001 -013 -MY3.

Alex Scott is supported by EPSRC grant EP/X013642/1.

David Wood is supported by the Australian Research Council and by NSERC..

FIGURE 1. Portions of two adjacent triangulations of an n -gon.

large n , and Pournin [24] showed that $2n - 10$ is the answer for $n > 12$. Several authors [10, 19, 20] studied random walks in \mathcal{A}_n .

This paper studies the chromatic number of \mathcal{A}_n , a quantity which was first considered by Fabila-Monroy et al. [11]. That work gave an explicit $\lceil \frac{n}{2} \rceil$ -colouring of \mathcal{A}_n , and observed that $\chi(\mathcal{A}_n) \in O(n/\log n)$, based on the result of Johansson [14] which says that every triangle-free graph with maximum degree Δ is $O(\Delta/\log \Delta)$ -colourable. No non-constant lower bound on $\chi(\mathcal{A}_n)$ is known. Indeed, the best known lower bound is $\chi(\mathcal{A}_{10}) \geq 4$ [private communication, Ruy Fabila-Monroy]. Fabila-Monroy et al. [11] conjectured a $O(\log n)$ upper bound. We prove this conjecture.

Theorem 1. $\chi(\mathcal{A}_n) \in O(\log n)$.

We prove Theorem 1 by tracking how several carefully chosen properties of triangulations change when an edge is flipped. To see how this yields a route to bounding the chromatic number of \mathcal{A}_n , first recall that if f is a graph homomorphism from \mathcal{A}_n to some graph G , which is to say that $f : V(\mathcal{A}_n) \rightarrow V(G)$ is adjacency-preserving, then $\chi(\mathcal{A}_n) \leq \chi(G)$. This fact may be generalized as follows. Suppose that $(G_i)_{i \in I}$ is a finite set of graphs and $(f_i : V(\mathcal{A}_n) \rightarrow V(G_i))_{i \in I}$ are functions such that for all adjacent triangulations T, T' in \mathcal{A}_n , there exists $i \in I$ for which $f_i(T)$ and $f_i(T')$ are adjacent in G_i . For each $i \in I$, let κ_i be a proper colouring of G_i with $\chi(G_i)$ colours, and colour each T in \mathcal{A}_n with the vector $(\kappa_i(f_i(T)))_{i \in I}$. If T and T' are adjacent in \mathcal{A}_n , then $(\kappa_i(f_i(T)))_{i \in I}$ and $(\kappa_i(f_i(T')))_{i \in I}$ differ in at least one coordinate. Thus this is a proper colouring of \mathcal{A}_n , and $\chi(\mathcal{A}_n) \leq \prod_{i \in I} \chi(G_i)$. The remainder of the paper is devoted to defining the five functions that we use (see (3)), and showing they have the requisite properties.

2. TYPES AND SCALES

Two fundamental notions that we use are the *type* of a quadrilateral and the *scale* of an edge. For a quadrilateral $Q = Q_T(e) = ABCD$ contained within triangulation T , we say Q is *type-1* if $e = AC$, and otherwise say Q is *type-2*; we say that an edge e is *type-1* or *type-2* according to the type of the quadrilateral $Q_T(e)$. For example, in Figure 1(A), $Q_T = ABCD$ is type-1 and $Q_T(BC) = ABEC$ is type-2, and in Figure 1(B), $Q_{T'}(BC) = BECD$ is type-1 and $Q_{T'}(BD) = ABCD$ is type-2.

Fix an integer $\alpha \geq 3$ to be chosen later (in fact we end up taking $\alpha = 3$). For an edge $e = UV$, define the *scale* of e to be

$$\sigma_e := \lceil \log_\alpha |U - V| \rceil \in \{0, 1, \dots, \lceil \log_\alpha(n-1) \rceil\}.$$

Note that $\sigma_e = 0$ if and only if e is a boundary edge. The scales of the edges incident to triangles within a fixed quadrilateral Q are a key input to the functions we define.

We first consider the effect of edge flips on triangles ABC , where two of the three incident edges have the same scale. If AB (resp. BC , AC) is the unique edge whose scale is different from the others, then we say ABC is a *type- ℓ* (resp. *type- m* , *type- r*) triangle. If all three edges have the same scale, then we say ABC is a *type- z* triangle. Let (ℓ_T, m_T, r_T, z_T) be the vector counting the number of type- ℓ , type- m , type- r and type- z triangles in T .

For the remainder of the paper, we fix a triangulation T , and consider the effect of flipping an edge $e = AC$ within a type-1 quadrilateral $Q_T(e) = ABCD$, to form another triangulation T' .

The next three propositions respectively consider the cases where $|\{\sigma_{AC}, \sigma_{BD}, \sigma_{BC}\}|$ is 3, 2 or 1.

Proposition 2. *If σ_{AC} , σ_{BD} and σ_{BC} are all different, then $(\ell_{T'}, m_{T'}, r_{T'}, z_{T'}) \neq (\ell_T, m_T, r_T, z_T)$.*

Proof. By assumption, $\sigma_{BD} \neq \sigma_{AC}$ and $\sigma_{BC} < \min(\sigma_{AC}, \sigma_{BD})$. We argue by contradiction. To this end, suppose that $(\ell_{T'}, m_{T'}, r_{T'}, z_{T'}) = (\ell_T, m_T, r_T, z_T)$. Since $\alpha \geq 3$,

$$\log_\alpha(D-A) \leq \log_\alpha(3 \max(B-A, C-B, D-C)) \leq 1 + \log_\alpha \max(B-A, C-B, D-C).$$

Taking ceilings, it follows that

$$\sigma_{AD} \leq 1 + \max(\sigma_{AB}, \sigma_{BC}, \sigma_{CD}). \quad (1)$$

The right-hand side of the preceding equation depends on three cases. We treat each possibility one at a time.

Suppose that $\sigma_{AB} = \max(\sigma_{AB}, \sigma_{BC}, \sigma_{CD})$. Using (1) and the fact that $\sigma_{AB} \leq \sigma_{AC} \leq \sigma_{AD}$, we find that either $\sigma_{AC} = \sigma_{AB}$ or $\sigma_{AC} = \sigma_{AB} + 1 = \sigma_{AD}$.

If $\sigma_{AC} = \sigma_{AB}$, as in Figure 2(A), then ABC is a type- m triangle so, since we assume the triangle type vector is unchanged by flipping edge AC , either ABD or BCD is also type- m . BCD is not type- m , as $\sigma_{BC} < \sigma_{BD}$ by assumption, so ABD is type- m and hence $\sigma_{AB} = \sigma_{AD}$. But then ABC and ACD are both type- m , which gives a contradiction as BCD is not.

If $\sigma_{AC} = \sigma_{AB} + 1$, as in Figure 2(B), then ACD is type- m or type- z , so either ABD or BCD is type- m or type- z . But BCD is neither, as $\sigma_{BC} < \sigma_{BD}$, and ABD is neither as $\sigma_{AB} \neq \sigma_{AD}$.

Next suppose that $\sigma_{BC} = \max(\sigma_{AB}, \sigma_{BC}, \sigma_{CD})$, as in Figure 2(C). Since $\sigma_{BC} \neq \sigma_{AC}$ and $\sigma_{BC} \neq \sigma_{BD}$ by assumption, and all scales are at most σ_{AD} , it must be that $\sigma_{AC} = \sigma_{BD} = \sigma_{AD}$; but this is ruled out by assumption.

Finally, suppose that $\sigma_{CD} = \max(\sigma_{AB}, \sigma_{BC}, \sigma_{CD})$. This case is symmetric to the first case, as we can apply the argument to a reversed copy of the associahedron (labelling the vertices counter-clockwise, which exchanges type- ℓ and type- m triangles, while leaving the other two types invariant). \square




FIGURE 2. Writing $\sigma = \max(\sigma_{AB}, \sigma_{BC}, \sigma_{CD})$, the subfigures correspond to possible configurations arising in the proof of Proposition 2.

Proposition 3. *If exactly two of σ_{AC} , σ_{BD} and σ_{BC} are equal, then $c(T') \neq c(T)$.*

Proof. First suppose that BC is not a boundary edge, and let V be the unique vertex of T with $B < V < C$ adjacent to both B and C . Note that $ABCD = Q_T(AC)$ is type-1 in T and $ABCD = Q_{T'}(BD)$ is type-2 in T' . Also, $Q_T(BC) = ABVC$ is type-2 in T and $Q_{T'}(BC) = BVCD$ is type-1 in T' . It is not hard to check that no other quadrilaterals change type when moving from T to T' . Thus

$$\begin{aligned} c(T') - c(T) &= 3\sigma_{BD} - 2\sigma_{AC} + 2\sigma_{BC} - 3\sigma_{BC} \quad \text{mod } (3\lceil \log_\alpha n \rceil) \\ &= 3\sigma_{BD} - 2\sigma_{AC} - \sigma_{BC} \quad \text{mod } (3\lceil \log_\alpha n \rceil). \end{aligned}$$

It follows that if $\sigma_{BC} < \sigma_{BD} = \sigma_{AC}$, then

$$c(T') - c(T) = \sigma_{BD} - \sigma_{BC} \quad \text{mod } (3\lceil \log_\alpha n \rceil) \neq 0;$$

the difference is non-zero modulo $(3\lceil \log_\alpha n \rceil)$ since all scales are at most $\lceil \log_\alpha n \rceil$. Similarly, if $\sigma_{BC} = \sigma_{BD} < \sigma_{AC}$ then

$$c(T') - c(T) = 2(\sigma_{AC} - \sigma_{BD}) \quad \text{mod } (3\lceil \log_\alpha n \rceil) \neq 0.$$

Finally, if $\sigma_{AC} = \sigma_{BC} < \sigma_{BD}$ then (since BD and AC are non-boundary edges)

$$c(T') - c(T) = 3(\sigma_{BD} - \sigma_{AC}) \quad \text{mod } (3\lceil \log_\alpha n \rceil) \neq 0$$

Since $\sigma_{BC} \leq \min(\sigma_{AC}, \sigma_{BD})$, these are the only possibilities. Here we use that $\sigma_{BD} \geq 1$ and $\sigma_{AC} \geq 1$ (since $D - B \geq 2$ and $C - A \geq 2$).

The case when BC is a boundary edge is very similar, but easier. In this case,

$$c(T') - c(T) = 3\sigma_{BD} - 2\sigma_{AC} \quad \text{mod } (3\lceil \log_\alpha n \rceil).$$

Since BC is a boundary edge, AC and BD are not, so $\sigma_{BC} = 0$ and $\sigma_{AC} \neq 0$ and $\sigma_{BD} \neq 0$. It follows by assumption that $\sigma_{AC} = \sigma_{BD}$, so

$$c(T') - c(T) = \sigma_{AC} \quad \text{mod } (3\lceil \log_\alpha n \rceil) \neq 0. \quad \square$$

Proposition 4. *If $\sigma_{AC} = \sigma_{BD} = \sigma_{BC}$ and $(\ell_{T'}, m_{T'}, r_{T'}, z_{T'}) = (\ell_T, m_T, r_T, z_T)$, then either $\sigma_{AD} = \sigma_{BC}$ or $\sigma_{AB} = \sigma_{BC} = \sigma_{CD}$.*

Proof. In this case ABC is type- ℓ or type- z , and BCD is type- m or type- z . If ABC is type- ℓ then since the triangle type vector does not change when flipping, it must be that ABD is type- ℓ , which implies that $\sigma_{AD} = \sigma_{BD}$, yielding the result in this case since $\sigma_{BD} = \sigma_{BC}$. Similarly, if BCD is type- m then it must be that

ACD is type- m , which implies that $\sigma_{AD} = \sigma_{AC}$. Otherwise, both ABC and BCD are type- z , in which case we indeed have $\sigma_{AB} = \sigma_{BC} = \sigma_{CD}$. \square

For each triangulation T in \mathcal{A}_n and $k \in \{1, 2\}$ and $i \in \{0, 1, \dots, \lceil \log_\alpha(n-1) \rceil\}$, let

$$s_i^k(T) := \#\{e \in E(T) : Q(e) \text{ is type-}k, \sigma_e = i\}.$$

Assign an integer label $c(T)$ to T given by

$$c(T) := \left(\sum_{i=0}^{\lceil \log_\alpha(n-1) \rceil} 2is_i^1(T) + \sum_{i=0}^{\lceil \log_\alpha(n-1) \rceil} 3is_i^2(T) \right) \pmod{3\lceil \log_\alpha n \rceil}.$$

The utility of such a labelling rule is explained by the following fact. We continue to work with triangulations T and T' related by an edge flip within quadrilateral $ABCD$ with $AC \in E(T)$ and $BD \in E(T')$, as above.

3. TWO MORE PARAMETERS

Propositions 2, 3 and 4 imply that the label $c(T)$ and the type vector (ℓ_T, m_T, r_T, z_T) together distinguish T from T' except in the following cases.

- (a) AC, BD, BC , and AD have the same scale and AB, CD have smaller scales.
- (b) AC, BD, BC, AD and AB have the same scale and CD has a smaller scale.
- (c) AC, BD, BC, AD and CD have the same scale and AB has a smaller scale.
- (d) AC, BD, BC, AB and CD have the same scale and AD has a larger scale.
- (e) All six edges AB, AC, AD, BC, BD and CD have the same scale.


To handle cases (a), (b) and (c) we track two additional parameters, and show that the parity of one or both parameters is different for T and T' . In cases (d) and (e) we again prove there is a change of parity, but of a third, more complicated parameter.

Figure 3(A) should make the following definitions clear. Orient the edges of the triangulation T so that the head of each edge has larger label. The *root edge* of T is the edge $\rho = (0, n-1)$. Now construct the following oriented tree \widehat{T} . First, augment T by adding a vertex v to the unbounded face, and join it to all vertices of the polygon. Let \widehat{T}_0 be the planar dual of the augmented graph; then \widehat{T} is formed from \widehat{T}_0 by removing all edges of \widehat{T}_0 lying entirely within the unbounded face of T . For each edge e of T there is a unique edge \hat{e} of \widehat{T} crossing e . Orient \hat{e} from the left to the right of e (when following e from tail to head). Root \widehat{T} at the edge $\hat{\rho}$, whose head is the unique vertex of \widehat{T} with out-degree 0. Note that \widehat{T} is a tree, which we call the *dual tree* of T .

Given an edge $e = UV$ of T with $e \neq \rho$, the triangle containing the head of \hat{e} is incident to both U and V ; let W be its third vertex. By the above choice of orientation for \hat{e} , either $W < \min(U, V)$ or $W > \max(U, V)$. In the first case, \hat{e} is a *left turn*, and in the second case it is a *right turn*.

Given a subgraph S of triangulation T , as illustrated in Figure 3(B), let \widehat{S} be the “dual” subgraph of \widehat{T} with the same vertex set as \widehat{T} and with edge set

$$E(\widehat{S}) := \{\hat{e} : e \in E(S)\}.$$



(A) The dual tree of a triangulation of an 8-gon. (B) The subgraph \widehat{S} of \widehat{T} corresponding to a subgraph S of a triangulation T .

FIGURE 3. The dual trees of an 8-gon and of a sub-triangulation of a 12-gon

A vertex of \widehat{S} is a *leaf* if it has degree 1. For each vertex v of \widehat{S} , let $L_S(v)$ and $R_S(v)$ be defined as follows. Write r for the root (the unique vertex of out-degree 0) of the tree component of \widehat{S} containing v . Then $L_S(v)$ and $R_S(v)$ are the number of left- and right-turns on the path from v to r , respectively; see Figure 4(A). (Figure 4(B) is used later in the section.)

Recall that T' is obtained from T by flipping edge AC within quadrilateral $ABCD$. For each $i \in \{1, 2, \dots, \lceil \log_\alpha n \rceil\}$, let S_i be the subgraph of T with edge set $E(S_i) = \{e \in E(T) : \sigma_e = i\}$, and let S'_i be the subgraph of T' with edge set $E(S'_i) = \{e \in E(T') : \sigma_e = i\}$. Define \widehat{S}_i and \widehat{S}'_i as in the preceding paragraph (so \widehat{S}_i is a subgraph of \widehat{T} and \widehat{S}'_i is a subgraph of \widehat{T}'). Let

$$L(T) := \sum_{i=1}^{\lceil \log_\alpha n \rceil} \sum_{v \in V(\widehat{S}_i)} L_{S_i}(v) \quad \text{and} \quad R(T) := \sum_{i=1}^{\lceil \log_\alpha n \rceil} \sum_{v \in V(\widehat{S}_i)} R_{S_i}(v).$$


The following proposition implies that in cases (a), (b) and (c), flipping edge AC yields a change in parity of at least one of G and D .

Proposition 5. *If the scales of the edges AB , AC , AD , BC , BD and CD are as in cases (a), (b) or (c) above, then $L(T') = L(T) - 1$ or $R(T') = R(T) + 1$ (or both).*

Proof. Let $\sigma = \sigma_{AC}$. We first claim that for all $i \neq \sigma$, the contributions to $L(T)$ and to $R(T)$ from scale- i vertices are unchanged by the edge flip operation; that is,

$$\sum_{i \neq \sigma} \sum_{v \in V(\widehat{S}_i)} L_{S_i}(v) = \sum_{i \neq \sigma} \sum_{v \in V(\widehat{S}'_i)} L_{S'_i}(v) \quad \text{and} \quad \sum_{i \neq \sigma} \sum_{v \in V(\widehat{S}_i)} R_{S_i}(v) = \sum_{i \neq \sigma} \sum_{v \in V(\widehat{S}'_i)} R_{S'_i}(v). \quad (2)$$

We prove these equalities in case (a); the other two cases are similar but easier.



(A) The left-turn and right-turn labelling of a component \hat{H} of \hat{S} with root vertex r . Labels are given in the form $(L(v), R(v))$ for all vertices v of \hat{S} . (B) The reduced tree \tilde{H} corresponding to the component \hat{H} , together with the triangulation of a polygon to which \tilde{H} is dual. The root edge of \tilde{H} is dashed.

FIGURE 4. In both subfigures, left-turn edges are red and right-turn edges are blue.

The triangle containing the head of the edge \hat{e}_{AB} dual to AB is ABC in T and is ABD in T' . The case (a) assumptions on the scales of the edges then imply that the head of \hat{e}_{AB} has out-degree 0 and in-degree 1 in $\hat{S}_{\sigma_{AB}}$. In particular, it is the root of its component of $\hat{S}_{\sigma_{AB}}$. Moreover, \hat{e}_{AB} is a right-turn edge in both T and T' , since C and D are both larger than A and B .

Similarly, the triangle containing the head of the edge \hat{e}_{CD} dual to CD is ACD in T and is BCD in T' . The assumptions on the scales of edges again imply that the head of \hat{e}_{CD} has in-degree 1 and out-degree 0 within $\hat{S}_{\sigma_{CD}}$, so is the root of its component of $\hat{S}_{\sigma_{CD}}$. Moreover, \hat{e}_{CD} is a left-turn edge in both T and T' , since A and B are both smaller than C and D .

Since the structures of T and of T' are unaffected outside of the quadrilateral $ABCD$, the equalities in (2) follow in case (a).

We now restrict our attention to the scale σ . We write $L(\cdot) = L_{S_\sigma}(\cdot)$ and $R(\cdot) = R_{S_\sigma}(\cdot)$, and likewise $L'(\cdot) = L_{S'_\sigma}(\cdot)$ and $R'(\cdot) = R_{S'_\sigma}(\cdot)$. Note that all vertices not lying within the quadrilateral $ABCD$ either belong to both S_σ and S'_σ or belong to neither of S_σ and S'_σ .

The remainder of the proof boils down to inspection of Figures 5, 6 and 7. In case (a), as illustrated in Figure 5, observe that $L(u) = L'(u) = a + 1$ and $R(u) = R'(u) = b + 1$, which implies that $(L(q), R(q)) = (L'(q), R'(q))$ for all vertices q not lying within $ABCD$. Since $L(v) + L(x) = 2a + 1 = L(p) + L(z) + 1$ and $R(v) + R(z) = 2b = R(p) + R(z) - 1$, it follows that $L(T) = L(T') + 1$ and $R(T') = R(T) - 1$.




FIGURE 5. The left-turn and right-turn labels near quadrilateral $ABCD$ in T and T' : case (a). Here $(L(x), R(x)) = (a, b)$, $(L(v), R(v)) = (a + 1, b)$ and $(L(u), R(u)) = (a + 1, b + 1)$.

Figure 6 depicts the situation in case (b). Here $R(u) = R'(u)$ and $R(w) = R'(w)$, which implies that $R(q) = R'(q)$ for all q not lying in $ABCD$. Since $R'(z) + R'(p) = R(v) + R(x) + 1$, it follows that $R(T') = R(T) + 1$.




FIGURE 6. The left-turn and right-turn labels near $ABCD$ in T and T' : case (b).

Finally, Figure 7 relates to case (c). In this case $L(u) = L'(u)$, $L(y) = L'(y)$, and $L(v) + L(x) = L'(z) + L(p) + 1$, so the above argument implies that $L(T') = L(T) - 1$. \square

We now turn our attention to cases (d) and (e). Consider any subgraph S of T , and let \hat{H} be a connected component of \hat{S} . Note that \hat{H} is a rooted sub-binary tree (that is, every vertex has degree at most three; see Figure 3(B)). Let \tilde{H} be the tree obtained from \hat{H} as follows (see Figure 4(A) and 4(B)). First, if the root r of \hat{H} has exactly two children then add a new vertex \tilde{r} incident only to r and reroot at \tilde{r} . Next, suppress each vertex of degree exactly two (that is, contract one edge incident to the vertex). We obtain a rooted binary tree \tilde{H} called the *reduced tree* of \hat{H} .




FIGURE 7. The left-turn and right-turn labels near $ABCD$ in T and T' : case (c).

Proposition 6. *For each $i \in \{1, 2, \dots, \lceil \log_\alpha n \rceil\}$ and for each component \hat{H} of \hat{S}_i , the reduced tree \tilde{H} has at most $2\alpha - 1$ leaves.*

Proof. Fix any vertex u of \tilde{H} with in-degree zero, and consider the edge uv incident to u in \tilde{H} . Then uv is dual to an edge AB with $\sigma_{AB} = i$ so with $\alpha^{i-1} < B - A \leq \alpha^i$. Now fix another vertex w of \tilde{H} with in-degree zero, write wx for the edge incident to w in \tilde{H} , and let CD be its dual edge. Then necessarily either $A < B \leq C < D$ or $C < D \leq B < A$.

Consider an oriented edge yr where r is the root of \hat{H} . Writing EF for the edge dual to yr , then the observation of the preceding paragraph implies that $F - E > \alpha^{i-1} \cdot \ell$, where ℓ is the number of vertices with in-degree zero in the subtree rooted at y . On the other hand, $\sigma_{EF} = i$ so $F - E \leq \alpha^i$; so $\ell \leq \alpha - 1$. If r has only one child (so is a leaf itself) this yields that \tilde{H} has at most α leaves. If r has two children then each of their subtrees contains at most $\alpha - 1$ leaves; in this case \tilde{r} is also a leaf, so the total number of leaves is at most $2(\alpha - 1) + 1$. \square

For any subgraph S of the triangulation T , the embedding of \hat{T} in the plane induces a total order of the connected components of \hat{S} , given by the order their roots are visited by a clockwise tour around the contour of \hat{T} starting from the head of the root edge ρ . For each $1 \leq i \leq \lceil \log_\alpha n \rceil$, list the components of \hat{S}_i in the order just described as $H_{i,1}, \dots, H_{i,\ell}$, where $\ell = \ell(T, i)$ is the number of such components. Then, for $1 \leq j \leq \ell(T, i)$ let $\tilde{H}_{i,j}$ be the reduced tree of $H_{i,j}$.

Each tree from $(\tilde{H}_{i,j}, i \leq \lceil \log_\alpha n \rceil, j \leq \ell(T, i))$ is a dual to a unique triangulation $\tilde{T}_{i,j}$ of a polygon, as in Figure 4(B). Proposition 6 implies that $\tilde{T}_{i,j}$ belongs to an associahedron \mathcal{A}_k for some $k \leq 2\alpha - 1$. Let ϕ be a proper colouring of the disjoint union of $(\mathcal{A}_k, k \leq 2\alpha - 1)$, with colours $\{1, \dots, \chi(\mathcal{A}_{2\alpha-1})\}$, and define

$$I(T) := \left(\sum_{i=0}^{\lceil \log_\alpha n \rceil} \sum_{j=1}^{\ell(T,i)} \phi(\tilde{T}_{i,j}) \right) \pmod{\chi(\mathcal{A}_{2\alpha-1})}.$$

Proposition 7. *In cases (d) and (e), we have $I(T) \neq I(T')$.*

Proof. Write $vx = \hat{e}$ for the dual edge of AC in T , and $zp = \hat{e}_{BD}$ for the dual edge of BD in T' . Figures 8 and 9 illustrate cases (d) and (e) respectively.




FIGURE 8. The structure near the quadrilateral $ABCD$ in T and T' case (d). The dashed edges have scale $\sigma_{AD} > \sigma$, all other edges of the triangulations shown in the figure have scale σ .




FIGURE 9. The structure near the quadrilateral $ABCD$ in T and T' in case (e). All edges of the triangulations shown in the figure have scale σ .

The clockwise contour exploration of a rooted plane tree is a walk around the outside of the tree which starts and finishes at the root, keeping the unbounded face to its left at all times. This walk traverses each edge exactly twice, and records the vertices it visits in sequence, with repetition. In cases (d) and (e), for the clockwise contour explorations of \hat{T} and of \hat{T}' , there are (possibly empty) strings P_1, \dots, P_5 so that the sequences recorded by the contour explorations of \hat{T} and of \hat{T}' are respectively of the form

$$P_1 s x v w P_2 w v u P_3 u v x y P_4 y x s P_5 \quad \text{and} \quad P_1 s p w P_2 w p z u P_3 u z y P_4 y z p s P_5 ;$$

see Figures 8 and 9.

The contour explorations of \widehat{T} and \widehat{T}' agree until they visit dual vertices lying within $ABCD$. It follows that if $H_{\sigma,j}$ is the component of S_σ containing \hat{e}_{AC} , then the component of S'_σ containing \hat{e}_{BD} is $H'_{\sigma,j}$.

In case (d), since x has two children in $H_{\sigma,j}$, by construction it is the unique child of the root of $\tilde{H}_{\sigma,j}$. It is thus natural to identify s with the root of $\tilde{H}_{\sigma,j}$. We may likewise identify s with the root of $\tilde{H}'_{\sigma,j}$, since p has two children in $H'_{\sigma,j}$. After the addition of s as a root, the vertices v, x, p and z all have degree 3, so none of these vertices are suppressed when constructing $\tilde{H}_{\sigma,j}$ and $\tilde{H}'_{\sigma,j}$ from $H_{\sigma,j}$ and $H'_{\sigma,j}$.

In case (e), vertices v, x, p and z all have degree 3, and the edges xs and ps belong to $H_{\sigma,j}$ and $H'_{\sigma,j}$, respectively.

It follows from the two preceding paragraphs that in cases (d) and (e), we may view v and x as vertices of both $H_{\sigma,j}$ and $\tilde{H}_{\sigma,j}$, and p and z as vertices of both $H'_{\sigma,j}$ and $\tilde{H}'_{\sigma,j}$. It is then clear that flipping the edge AC in the triangulation T corresponds precisely to flipping the corresponding edge in $\tilde{T}_{\sigma,j}$ to form $\tilde{T}'_{\sigma,j}$.

Since $\tilde{T}_{\sigma,j}$ and $\tilde{T}'_{\sigma,j}$ are related by a single edge flip, and ϕ is a proper colouring, it follows that $\phi(\tilde{T}_{\sigma,j}) \neq \phi(\tilde{T}'_{\sigma,j})$. Since all other components of S_σ , and more generally of each $(S_i, 1 \leq i \leq \lceil \log_\alpha n \rceil)$, are unchanged when moving from T to T' , the result follows. \square

4. FINISHING THE PROOF

Proof of Theorem 1. Consider the graph G with vertex set \mathbb{N}^4 where distinct vertices (ℓ, m, r, z) and (ℓ', m', r', z') are adjacent if $|\ell' - \ell| + |m' - m| + |r' - r| + |z' - z| \leq 4$. Define $\kappa(\ell, m, r, z) := (\ell \bmod 5, m \bmod 5, r \bmod 5, z \bmod 5)$ for each vertex (ℓ, m, r, z) of G . Consider distinct vertices (ℓ, m, r, z) and (ℓ', m', r', z') in G with $\kappa(\ell, m, r, z) := \kappa(\ell', m', r', z')$. Without loss of generality, $\ell \neq \ell'$. Thus $|\ell - \ell'| \geq 5$, implying that (ℓ, m, r, z) and (ℓ', m', r', z') are not adjacent in G . Hence κ is a 625-colouring of G . (We are not optimizing constants.)

For each triangulation T of \mathcal{A}_n , colour T by the 5-tuple

$$\psi(T) := (\kappa((\ell_T, m_T, r_T, z_T)), c(T), L(T) \bmod 2, R(T) \bmod 2, I(T)). \quad (3)$$

We now show that ψ is a proper colouring of \mathcal{A}_n .

Consider adjacent vertices T and T' in \mathcal{A}_n . Since T and T' agree on all but four triangles, $|\ell_T - \ell_{T'}| + |m_T - m_{T'}| + |r_T - r_{T'}| + |z_T - z_{T'}| \leq 4$. In all situations covered by Proposition 2, the vectors $(\ell_{T'}, m_{T'}, r_{T'}, z_{T'})$ and (ℓ_T, m_T, r_T, z_T) are distinct and thus adjacent in G . Thus, $\kappa((\ell_{T'}, m_{T'}, r_{T'}, z_{T'})) \neq \kappa((\ell_T, m_T, r_T, z_T))$. In the situations covered by Proposition 3, we have $c(T) \neq c(T')$. By Proposition 5, in cases (a)–(c), either $L(T) \not\equiv L(T') \pmod{2}$ or $R(T) \not\equiv R(T') \pmod{2}$ or both. Finally, in cases (d) and (e), by Proposition 7 we have $I(T) \neq I(T')$.

Thus $\psi(T) \neq \psi(T')$, implying for any integer $\alpha \geq 3$,

$$\chi(\mathcal{A}_n) \leq 625 \cdot \lceil 3 \log_\alpha n \rceil \cdot 2 \cdot 2 \cdot \chi(\mathcal{A}_{2\alpha-1}).$$

Taking $\alpha = 3$ yields $\chi(\mathcal{A}_{2\alpha-1}) = \chi(\mathcal{A}_5) = 3$, since \mathcal{A}_5 is a 5-cycle. It follows that

$$\chi(\mathcal{A}_n) \leq 625 \cdot \lceil 3 \log_3 n \rceil \cdot 12 \in O(\log n). \quad \square$$

Acknowledgements. This research was initiated at the 2018 Barbados Workshop on Graph Theory, held at the Bellairs Research Institute of McGill University. Thanks to the referees for several helpful comments.

REFERENCES

- [1] OSWIN AICHHOLZER, WOLFGANG MULZER, AND ALEXANDER PILZ. [Flip distance between triangulations of a simple polygon is NP-complete](#). *Discrete Comput. Geom.*, 54(2):368–389, 2015. MR: [3372115](#).
- [2] LOUIS J. BILLERA, PAUL FILLIMAN, AND BERND STURMFELS. [Constructions and complexity of secondary polytopes](#). *Adv. Math.*, 83(2):155–179, 1990. MR: [1074022](#).
- [3] PROSENJIT BOSE, ANNA LUBIW, VINAYAK PATHAK, AND SANDER VERDONSCHOT. [Flipping edge-labelled triangulations](#). *Comput. Geom.*, 68:309–326, 2018. MR: [3715060](#).
- [4] JEAN CARDINAL, STEFAN LANGERMAN, AND PABLO PÉREZ-LANTERO. [On the diameter of tree associahedra](#). *Electron. J. Combin.*, 25(4):#4.18, 2018. MR: [3874284](#).
- [5] CESAR CEBALLOS AND VINCENT PILAUD. [The diameter of type \$D\$ associahedra and the non-leaving-face property](#). *European J. Combin.*, 51:109–124, 2016. MR: [3398843](#).
- [6] FRÉDÉRIC CHAPOTON, SERGEY FOMIN, AND ANDREI ZELEVINSKY. [Polytopal realizations of generalized associahedra](#). *Canad. Math. Bull.*, 45(4):537–566, 2002. MR: [1941227](#).
- [7] SEAN CLEARY AND ROLAND MAIO. [Edge conflicts do not determine geodesics in the associahedron](#). *SIAM J. Discrete Math.*, 32(2):1003–1015, 2018. MR: [3796362](#).
- [8] NICOLAAS G. DE BRUIJN AND PAUL ERDŐS. On a combinatorial problem. *Nederl. Akad. Wetensch., Proc.*, 51:1277–1279 = *Indagationes Math.* 10, 421–423 (1948), 1948. MR: [0028289 \(10,424a\)](#).
- [9] PATRICK DEHORNOY. [On the rotation distance between binary trees](#). *Adv. Math.*, 223(4):1316–1355, 2010. MR: [2581373](#).
- [10] DAVID EPPSTEIN AND DANIEL FRISHBERG. [Improved mixing for the convex polygon triangulation flip walk](#). In KOUSHA ETESSAMI, URIEL FEIGE, AND GABRIELE PUPPIS, eds., *Proc. 50th Int’l Colloquium on Automata, Languages, and Programming (ICALP 2023)*, vol. 261 of *LIPICs*, pp. 56:1–56:17. Schloss Dagstuhl, 2023.
- [11] RUY FABILA-MONROY, DAVID FLORES-PEÑALOZA, CLEMENS HUEMER, FERRAN HURTADO, JORGE URRUTIA, AND DAVID R. WOOD. [On the chromatic number of some flip graphs](#). *Discrete Math. Theor. Comput. Sci.*, 11(2):47–56, 2009. MR: [2535071](#).
- [12] CHRISTOPHE HOHLWEG, CARSTEN E. M. C. LANGE, AND HUGH THOMAS. [Permutahedra and generalized associahedra](#). *Adv. Math.*, 226(1):608–640, 2011. MR: [2735770](#).
- [13] FERRAN HURTADO AND MARC NOY. [Graph of triangulations of a convex polygon and tree of triangulations](#). *Comput. Geom. Theory Appl.*, 13(3):179–188, 1999. MR: [1723053](#).
- [14] ANDERS JOHANSSON. Some results on colourings of graphs. Ph.D. thesis, University of Umeå, 1994.

- [15] CARL W. LEE. [The associahedron and triangulations of the \$n\$ -gon.](#) *European J. Combin.*, 10(6):551–560, 1989. MR: 1022776.
- [16] JEAN-LOUIS LODAY AND MARÍA O. RONCO. [Hopf algebra of the planar binary trees.](#) *Adv. Math.*, 139(2):293–309, 1998. MR: 1654173.
- [17] JOAN M. LUCAS. [The rotation graph of binary trees is Hamiltonian.](#) *J. Algorithms*, 8(4):503–535, 1987. MR: 920505.
- [18] THIBAUT MANNEVILLE AND VINCENT PILAUD. [Graph properties of graph associahedra.](#) *Sém. Lothar. Combin.*, B73d, 2014–2016. MR: 3383157.
- [19] LISA MCSHINE AND PRASAD TETALI. [On the mixing time of the triangulation walk and other Catalan structures.](#) In PANOS M. PARDALOS, SANGUTHEVAR RAJASEKARAN, AND JOSÉ ROLIM, eds., *Randomization Methods in Algorithm Design*, vol. 43 of *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.*, pp. 147–160. Amer. Math. Soc., 1997.
- [20] MICHAEL MOLLOY, BRUCE REED, AND WILLIAM STEIGER. [On the mixing rate of the triangulation walk.](#) In *Randomization methods in algorithm design*, vol. 43 of *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.*, pp. 179–190. Amer. Math. Soc., 1999.
- [21] HUGO PARLIER AND BRAM PETRI. [The genus of curve, pants and flip graphs.](#) *Discrete Comput. Geom.*, 59(1):1–30, 2018. MR: 3738334.
- [22] VINCENT PILAUD AND FRANCISCO SANTOS. [The brick polytope of a sorting network.](#) *European J. Combin.*, 33(4):632–662, 2012. MR: 2864447.
- [23] VINCENT PILAUD AND CHRISTIAN STUMP. [Brick polytopes of spherical subword complexes and generalized associahedra.](#) *Adv. Math.*, 276:1–61, 2015. MR: 3327085.
- [24] LIONEL POURNIN. [The diameter of associahedra.](#) *Adv. Math.*, 259:13–42, 2014. MR: 3197650.
- [25] LIONEL POURNIN. [The asymptotic diameter of cyclohedra.](#) *Israel J. Math.*, 219(2):609–635, 2017. MR: 3649601.
- [26] LIONEL POURNIN. [Eccentricities in the flip-graphs of convex polygons.](#) *J. Graph Theory*, 92(2):111–129, 2019. MR: 3994734.
- [27] DANIEL D. SLEATOR, ROBERT E. TARJAN, AND WILLIAM P. THURSTON. [Rotation distance, triangulations, and hyperbolic geometry.](#) *J. Amer. Math. Soc.*, 1(3):647–681, 1988. MR: 928904.
- [28] N. J. A. SLOANE. [Coordination sequence for 4-dimensional cubic lattice.](#) *The On-Line Encyclopedia of Integer Sequences*, A008412, 2025.
- [29] JAMES DILLON STASHEFF. [Homotopy associativity of \$H\$ -spaces. I, II.](#) *Trans. Amer. Math. Soc.*, 108:293–312, 1963. MR: 0158400.
- [30] DOV TAMARI. [Monoïdes préordonnés et chaînes de Malcev.](#) Thèse, Université de Paris, 1951.

(L. Addario-Berry) DEPARTMENT OF MATHEMATICS AND STATISTICS, MCGILL UNIVERSITY,
MONTRÉAL, QUÉBEC, CANADA
Email address: `louigi@gmail.com`

(B. Reed) INSTITUTE OF MATHEMATICS, ACADEMIA SINICA, TAIWAN
Email address: `bruce.al.reed@gmail.com`

(A. Scott) MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, OXFORD OX2 6GG, UNITED
KINGDOM
Email address: `scott@maths.ox.ac.uk`

(D. R. Wood) SCHOOL OF MATHEMATICS, MONASH UNIVERSITY, MELBOURNE, AUSTRALIA
Email address: `david.wood@monash.edu`